

An Overview of Common Pediatric Malocclusions and Treatment Modalities

Alexander Schmidt*

School of Dentistry, University of Munich, Munich, Bavaria, Germany

DESCRIPTION

Malocclusion, which refers to the misalignment of teeth and jaws, is a common dental concern in pediatric patients and can significantly impact both oral health and self-esteem. It can range from minor misalignments to more severe jaw discrepancies, affecting a child's ability to chew, speak, or maintain proper oral hygiene. Malocclusions can be classified into different types, including Class I, Class II, Class III, open bite, and crossbite, each with its distinct characteristics and treatment approaches. Class I malocclusion is considered the most common, where the upper teeth slightly overlap the lower teeth, but the bite remains functional. Treatment for Class I malocclusion generally focuses on improving alignment through braces or clear aligners, often with the addition of retainers or expanders to guide proper eruption of the teeth. Class II malocclusion, also known as an overbite, is characterized by the upper teeth overlapping significantly over the lower teeth [1-2].

This condition can be further divided into two subcategories: Class II Division 1, where the upper teeth protrude outward, and Class II Division 2, where the upper teeth tilt inward. Early intervention is crucial for Class II malocclusions, and treatment typically involves using appliances like headgear or functional appliances to realign the jaw, followed by braces or clear aligners to refine tooth placement. In more severe cases, surgical intervention may be necessary once the child's growth has slowed or stopped. Class III malocclusion, or an underbite, occurs when the lower jaw protrudes forward, causing the lower teeth to extend beyond the upper teeth. This condition often requires a combination of functional appliances and, in certain cases, jaw surgery to reposition the lower jaw. Early orthodontic treatment using appliances like the Herbst appliance or Frankel appliance can help guide the jaw into a more favorable position. An open bite, which occurs when the upper and lower teeth do not meet when the mouth is closed, often results from habits like thumb-sucking or tongue thrusting [3-6].

Treatment for open bite usually involves a combination of orthodontic appliances, tongue training exercises, and in severe cases, surgical intervention. Similarly, a crossbite occurs when the upper teeth are positioned inside the lower teeth, either in the front or back of the mouth. Crossbites can lead to tooth

wear, gum recession, and asymmetric jaw growth if not addressed early. Treatment typically involves using palatal expanders to widen the upper jaw, along with braces or functional appliances to align the teeth properly. The primary treatment modality for pediatric malocclusions is orthodontic care, which includes options such as traditional braces, clear aligners, space maintainers, and in more severe cases, jaw surgery. Braces, the most common orthodontic treatment, consist of brackets and wires that gradually shift teeth into the correct position. Clear aligners offer a less visible alternative, particularly for mild to moderate misalignments, and are popular among older children and adolescents [7-8].

Space maintainers are used when children lose baby teeth prematurely, preventing neighboring teeth from shifting and ensuring space for the permanent teeth to emerge. In more complex cases, jaw surgery may be required once the child's growth has stabilized to correct severe misalignment and optimize function. Early diagnosis and intervention are crucial in managing malocclusions to ensure proper oral development and avoid long-term dental and functional issues. Pediatric dentists play a vital role in diagnosing these conditions and guiding children through the treatment process to ensure that they achieve optimal dental health and a confident smile [9-10].

CONCLUSION

In conclusion, pediatric malocclusions are a common but significant concern in children's dental health, affecting not only the alignment of teeth but also their overall oral function and self-esteem. Early diagnosis and timely intervention are key to preventing the progression of these conditions and ensuring that children grow into adulthood with healthy, properly aligned teeth. From mild issues like Class I malocclusions to more complex cases like Class II or Class III misalignments, various treatment options are available, including braces, clear aligners, functional appliances, and, in some cases, surgery. The role of pediatric dentists is crucial in identifying these conditions early and guiding children through the most appropriate treatment modalities. By addressing malocclusions during the critical stages of dental development, it is possible to improve a child's oral health, function, and appearance, setting them on a path to a

Correspondence to: Alexander Schmidt, School of Dentistry, University of Munich, Munich, Bavaria, Germany, E-mail: alexander.schmidt@dent.um.edu

Received: 27-Jun-2025, Manuscript No. AEDJ-25-40344; **Editor assigned:** 30-Jun-2025, PreQC No. AEDJ-25-40344 (PQ); **Reviewed:** 15-Jul-2025, QC No. AEDJ-25-40344; **Revised:** 22-Jul-2025, Manuscript No. AEDJ-25-40344 (R); **Published:** 28-Jul-2025, DOI: 10.35248/0976-156X.25.17.336

Citation: Schmidt A (2025) An Overview of Common Pediatric Malocclusions and Treatment Modalities. Ann Essence Dent. 17:336.

Copyright: © 2025 Schmidt A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

lifetime of optimal dental well-being. Ultimately, the goal of pediatric orthodontics is not just to achieve a beautiful smile, but also to ensure proper jaw alignment and overall oral health, enhancing the child's quality of life both now and in the future.

REFERENCES

1. Liu Y, Liu Y, Guo J, Jin F, Cao Y, Jiang L. et al. Expert consensus on pediatric orthodontic therapies of malocclusions in children. *Int J Oral Sci.* 2024;16(1):32.
2. Achmad H, Auliya N. Management of Malocclusion in Children using Myobrace Appliance: a systematic review. *Ann Rom Soc Cell Biol.* 2021;25(2):2120-2136.
3. Zou J, Meng M, Law CS, Rao Y, Zhou X. Common dental diseases in children and malocclusion. *Int. J Oral Sci* 2018;10(1):7.
4. Kanas R, Carapezza L, Kanas S. Treatment classification of Class III malocclusion. *J Clin Pediatr Dent.* 2008;33(2):175-186.
5. Stomatologic S. Worldwide prevalence of malocclusion in the different stages of dentition. *Eur J Paediatr Dent.* 2020;21(2): 115-122.
6. Bondemark L, Kallunki J, Paulsson L. An updated systematic review regarding early Class II malocclusion correction. *JWFO.* 2019;8(3): 89-94.
7. Alqahtani AA, Alkahtani LA, Abushiqa AA, Alasmari AS, Al-Jalooud AH. Dental occlusion and malocclusion: Prevalence, types and treatment. *EC Dental Science.* 2019;1777.
8. Lin LH, Huang GW, Chen CS. Etiology and treatment modalities of anterior open bite malocclusion. *J Exp Clin Med* 2013;5(1):1-4.
9. Dolker T, Shadani K, Galhotra V, Rathod P, Gandham R, Mukhtar U. et al. Interceptive treatment modalities in class 2 malocclusion-A review of literature. *JAMDSR.* 2023;11(9):14-18.
10. Andresen T, Bahr C, Ciranna-Raab C. Efficacy of osteopathy and other manual treatment approaches for malocclusion-A systematic review of evidence. *Int J Osteopath Med.* 2013;16(2):99-113.