OPEN ACCESS Freely available online

Advances in Medical Research

Perspective

Cardiovascular Disease in Women: Gender-Specific Risk Factors and Management

Benjamin James*

Department of Cardiology, New York Medical College, Valhalla, New York, United States of America

DESCRIPTION

Cardiovascular Disease (CVD) remains one of the leading causes of morbidity and mortality among women globally. However, despite its prominence, heart disease in women has historically been under-recognized, with much of the research and clinical focus being centered on male populations. As awareness grows, it has become increasingly clear that women experience CVD differently than men, with unique risk factors, symptoms, and outcomes. Understanding these gender-specific nuances is important for improving diagnosis, prevention, and treatment for women. This article explores the distinctive risk factors that contribute to CVD in women and the evolving approach to managing heart disease with a gender-specific lens.

Gender-specific risk factors

While traditional cardiovascular risk factors such as high blood pressure, high cholesterol, diabetes, smoking, and obesity affect both men and women, several gender-specific factors influence cardiovascular health in women. These factors include hormonal influences, pregnancy-related conditions, and psychosocial elements that uniquely impact women's cardiovascular risk.

Hormonal factors: Estrogen, a hormone that predominates in women before menopause, has a protective effect on cardiovascular health by helping to maintain healthy cholesterol levels and promote arterial flexibility. However, after menopause, estrogen levels decline significantly, and the protective benefit is lost. As a result, women's risk for heart disease increases significantly post-menopause, often surpassing that of men at similar ages. This shift highlights the importance of considering age and hormonal status when assessing cardiovascular risk in women.

Pregnancy-related conditions: Pregnancy is an essential phase in a woman's life, but it also serves as a unique window into a woman's future cardiovascular health. Conditions such as preeclampsia, gestational diabetes, and gestational hypertension are all known to increase the risk of developing CVD later in life. Women who experience preeclampsia, for example, have a higher risk of developing hypertension and heart disease within a decade of giving birth. Furthermore, gestational diabetes can increase the likelihood of type 2 diabetes, which in turn raises the risk of CVD. These pregnancy-related complications provide critical signals that clinicians should monitor over a woman's lifetime to mitigate long-term cardiovascular risks.

Psychosocial factors: Women are more likely than men to experience mental health issues such as depression and anxiety, which can adversely affect cardiovascular health. The link between mental health and heart disease is particularly strong in women, with depression being identified as a major risk factor for cardiovascular events. Stress, caregiving responsibilities, and social isolation also disproportionately affect women and can contribute to increased cardiovascular risk. These factors, when not addressed, can lead to poor health behaviors (e.g., poor diet, lack of exercise) and physiological changes (e.g., increased inflammation, hypertension) that heighten the risk of heart disease.

Gender-specific management and treatment

The management of CVD in women requires a customized, gender-sensitive approach. Treatment plans should address not only traditional cardiovascular risk factors but also the unique challenges women face, including hormonal changes, pregnancy history, and mental health concerns.

Prevention and early detection: Preventive strategies in women should begin early, especially for those with pregnancy-related complications such as preeclampsia or gestational diabetes. Lifestyle interventions including regular exercise, a balanced diet, smoking cessation, and stress management should be emphasized. Screening for hypertension, diabetes, and lipid disorders is essential, particularly in women with risk factors like a history of preeclampsia or gestational diabetes. Healthcare providers should also be vigilant in monitoring women for mental health issues, providing support for conditions like depression and anxiety, which are linked to poorer cardiovascular outcomes.

Correspondence to: Benjamin James, Department of Cardiology, New York Medical College, Valhalla, New York, United States of America; E-mail: benjaminj@gmail.com

Received: 23-Dec-2024, Manuscript No. IDAMR-24-36172; **Editor assigned:** 25-Dec-2024, PreQC No. IDAMR-24-3672 (PQ); **Reviewed:** 08-Jan-2024, QC No. IDAMR-24-36172; **Revised:** 24-Sep-2025, Manuscript No. IDAMR-24-36172 (R); **Published:** 01-Oct-2025, DOI: 10.12715/2564-8942.25.8.071

Citation: James B (2025) Cardiovascular Disease in Women: Gender-Specific Risk Factors and Management. J Adv Med Res. 8:071.

Copyright: © 2025 James B. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Pharmacological interventions: The pharmacological treatment of CVD in women is often similar to that in men, with the use of statins, antihypertensive medications, antiplatelet agents, and diabetes management medications. However, women may respond differently to certain medications, which can affect drug efficacy and side effects. For example, women are more likely to experience adverse effects from blood thinners like aspirin, which can complicate management. The use of Hormone Replacement Therapy (HRT) in postmenopausal women also remains controversial. While HRT may offer benefits for symptom relief, it can also increase the risk of heart disease in certain women, necessitating careful risk assessment before initiation.

CONCLUSION

CVD in women is a complex issue that requires a gender-specific approach to risk assessment, diagnosis, and management. By recognizing the unique risk factors that affect women such as hormonal changes, pregnancy-related conditions, and psychosocial stressors healthcare providers can deliver more effective care. The differences in symptoms and outcomes also necessitate a more nuanced understanding of how heart disease presents and progresses in women. With early detection, personalized management plans, and an emphasis on mental health, we can improve cardiovascular outcomes for women, ultimately saving lives and improving quality of life. As awareness continues to grow, the integration of gender-specific strategies in cardiovascular care will be important in reducing the burden of heart disease among women worldwide.