

# On the space of coefficients in a statistical type metric space

**Bilalov B.T.**

Institute of Mathematics and Mechanics of Azerbaijan National Academy of Sciences  
9, B.Vahabzade Str., Az-1141, Baku, Azerbaijan  
bilalov.bilal@gmail.com

**Guliyeva F.A.**

Institute of Mathematics and Mechanics of Azerbaijan National Academy of Sciences  
9, B.Vahabzade Str., Az-1141, Baku, Azerbaijan  
guliyeva-fatima@mail.ru

## Abstract

Fuzzy Banach space is considered. The concepts of fuzzy completeness, fuzzy minimality, fuzzy biorthogonality, fuzzy basicity and fuzzy space of coefficients are introduced. Weakly completeness of fuzzy space of coefficients with regard to fuzzy norm and weakly basicity of canonical system in this space are proved. Weakly basicity criterion in fuzzy Banach space is presented in terms of coefficient operator.

**Mathematics Subject Classification:** 46A35, 46S40, 26E50

**Keywords:** fuzzy basicity, fuzzy completeness, fuzzy minimality, fuzzy biorthogonality, fuzzy space of coefficients

## 1 Introduction

The concept of the space of coefficients belongs to the theory of bases. As is known, every basis in a Banach space has a Banach space of coefficients which is isomorphic to an initial one (see, e.g., [1;2]). Every nondegenerate system (to be defined later) in a Banach space generates the corresponding Banach space of coefficients with canonical basis (see, e.g., [2;3]). Therefore, space of coefficients plays an important role in the study of approximative properties of systems. It has very important applications in various fields of science, such as solid body physics, molecular physics, multiple production of particles, aviation, medicine, biology, data compression, etc (see, e.g., [4;5] and references within). All these applications are closely related to wavelet analysis, and there arose a great interest in them lately [see, e.g., 5]. It is well

known that many topological spaces are nonnormable. Therefore, the study of various properties of the space of coefficients in topological spaces is of special scientific interest.

Applications in various branches of mathematics and natural sciences have lately induced a strong interest toward the study of different research problems in terms of fuzzy structures. More details on this topic can be found in [6-9] and references therein. A large number of research works is appearing these days which deal with the concept of fuzzy set-numbers, and fuzzification of many classical theories has also been made. The concept of Schauder basis in *intuitionistic fuzzy normed space* and some results related to this concept have recently been studied in [10-12;23-27]. These works introduced the concepts of strongly and weakly intuitionistic fuzzy (Schauder) basis in *intuitionistic fuzzy banach spaces* (IFBS in short). Some of their properties are revealed. The concepts of *strongly* and *weakly intuitionistic fuzzy approximation properties* (*sif-AP* and *wif-AP* in short, respectively) are also introduced in these works. It is proved that if the intuitionistic fuzzy space has a *wif-basis*, then it has a *wif-AP*. All the results in these works are obtained on condition that IFBS admits equivalent topology using the family of norms generated by *t-norm* and *t-conorm* (we will define them later).

In our work, we define the basic concepts of classical basis theory in *intuitionistic fuzzy normed spaces* (IFNS in short). Concepts of *weakly* and *strongly fuzzy spaces of coefficients* are introduced. *Weakly completeness* of these spaces with regard to *fuzzy norm* and *weakly basicity* of canonical system in them are proved. *Weakly basicity* criterion in fuzzy Banach space is presented in terms of coefficient operator.

In Section 2, we recall some notations and concepts. In Section 3, we state main results. We first define *fuzzy space of coefficients* and then introduce the corresponding *fuzzy norms*. We prove that for nondegenerate system the corresponding *fuzzy space of coefficients* is *weakly fuzzy complete*. Moreover, we show that the canonical system forms a *weakly basis* for this space.

## 2 Some preliminary notations and concepts

We will use the usual notations:  $N$  will denote the set of all positive integers,  $R$  will be the set of all real numbers,  $C$  will be the set of complex numbers and  $K$  will denote a field of scalars ( $K \equiv R$ , or  $K \equiv C$ ),  $R_+ \equiv (0, +\infty)$ . We state some concepts and facts from IFNS theory to be used later.

**Definition 2.1.** *Let  $X$  be a linear space over a field  $K$ . Functions  $\mu; \nu : X \times R \rightarrow [0, 1]$  are called fuzzy norms on  $X$  if the following conditions hold:*

1.  $\mu(x; t) = 0, \forall t \leq 0, \forall x \in X;$

2.  $\mu(x; t) = 1, \forall t > 0 \Rightarrow x = 0$ ;
3.  $\mu(cx; t) = \mu\left(x; \frac{t}{|c|}\right), \forall c \neq 0$ ;
4.  $\mu(x; \cdot) : R \rightarrow [0, 1]$  is a non-decreasing function of  $t$  for  $\forall x \in X$  and  $\lim_{t \rightarrow \infty} \mu(x; t) = 1, \forall x \in X$ ;
5.  $\mu(x + y; s + t) \geq \min\{\mu(x; s); \mu(y; t)\}, \forall x, y \in X, \forall s, t \in R$ ;
6.  $\nu(x; t) = 1, \forall t \leq 0, \forall x \in X$ ;
7.  $\nu(x; t) = 0, \forall t < 0 \Rightarrow x = 0$ ;
8.  $\nu(cx; t) = \nu\left(x; \frac{t}{|c|}\right), \forall c \neq 0$ ;
9.  $\nu(x; \cdot) : R \rightarrow [0, 1]$  is a non-increasing function of  $t$  for  $\forall x \in X$  and  $\lim_{t \rightarrow \infty} \nu(x; t) = 0, \forall x \in X$ ;
10.  $\nu(x + y; s + t) \leq \max\{\nu(x; s); \nu(y; t)\}, \forall x, y \in X, \forall s, t \in R$ ;
11.  $\mu(x; t) + \nu(x; t) \leq 1, \forall x \in X, \forall t \in R$ .

Then the triplet  $(X; \mu; \nu)$  is called an intuitionistic fuzzy normed space.

The above concepts allow to introduce the following kinds of convergence (or topology) in IFNS:

**Definition 2.2.** Let  $(X; \mu; \nu)$  be a fuzzy normed space and let  $\{x_n\}_{n \in N} \subset X$  be some sequence. Then it is said to be strongly intuitionistic fuzzy convergent to  $x \in X$  (denoted by  $x_n \xrightarrow{s} x, n \rightarrow \infty$  or  $s\text{-}\lim_{n \rightarrow \infty} x_n = x$  in short) if and only if for  $\forall \varepsilon > 0, \exists n_0 = n_0(\varepsilon) : \mu(x_n - x; t) \geq 1 - \varepsilon, \nu(x_n - x; t) \leq \varepsilon, \forall n \geq n_0, \forall t \in R$ .

**Definition 2.3.** Let  $(X; \mu; \nu)$  be a fuzzy normed space and let  $\{x_n\}_{n \in N} \subset X$  be some sequence. Then it is said to be weakly intuitionistic fuzzy convergent to  $x \in X$  (denoted by  $x_n \xrightarrow{w} x, n \rightarrow \infty$ , or  $w\text{-}\lim_{n \rightarrow \infty} x_n = x$  in short) if and only if for  $\forall t \in R_+, \forall \varepsilon > 0, \exists n_0 = n_0(\varepsilon; t) : \mu(x_n - x; t) \geq 1 - \varepsilon, \nu(x_n - x; t) \leq \varepsilon, \forall n \geq n_0$ .

More details on these concepts can be found in [12-22].

Let  $(X; \mu; \nu)$  be an IFNS, and let  $M \subset X$  be some set. By  $L[M]$  we denote the linear span of  $M$  in  $X$ . The weakly (strongly) intuitionistic fuzzy convergent closure of  $L[M]$  will be denoted by  $\overline{L_w[M]} (\overline{L_s[M]})$ . If  $X$  is complete with respect to the weakly (strongly) intuitionistic fuzzy convergence, then we will call it intuitionistic fuzzy weakly (strongly) Banach space ( $IFB_wS$  or  $X_w$

( $IFB_sS$  or  $X_s$ ) in short). Let  $X$  be an  $IFB_wS$  ( $IFB_sS$ ). We denote by  $X_w^*$  ( $X_s^*$ ) the linear space of linear and continuous in  $IFB_wS$  ( $IFB_sS$ ) functional over the same field  $K$ .

Now we define the corresponding concepts of basis theory for IFNS. Let  $\{x_n\}_{n \in N} \subset X$  be some system.

**Definition 2.4.** System  $\{x_n\}_{n \in N}$  is said  $w$ -complete ( $s$ -complete) in  $X_w$  (in  $X_s$ ), if  $\overline{L_w[\{x_n\}_{n \in N}]} \equiv X_w$  ( $\overline{L_s[\{x_n\}_{n \in N}]} \equiv X_s$ ).

**Definition 2.5.** System  $\{x_n^*\}_{n \in N} \subset X_w^*$  ( $\{x_n^*\}_{n \in N} \subset X_s^*$ ) is called  $w$ -biorthogonal ( $s$ -biorthogonal) to the system  $\{x_n\}_{n \in N}$ , if  $x_n^*(x_k) = \delta_{nk}$ ,  $\forall n, k \in N$ , where  $\delta_{nk}$  is the Kronecker symbol.

**Definition 2.6.** System  $\{x_n\}_{n \in N} \subset X_w$  ( $\{x_n\}_{n \in N} \subset X_s$ ) is called  $w$ -linearly ( $s$ -linearly) independent in  $X$ , if  $\sum_{n=1}^{\infty} \lambda_n x_n = 0$  in  $X_w$  (in  $X_s$ ) implies  $\lambda_n = 0$ ,  $\forall n \in N$ .

**Definition 2.7.** System  $\{x_n\}_{n \in N} \subset X_w$  ( $\{x_n\}_{n \in N} \subset X_s$ ) is called  $w$ -basis ( $s$ -basis) for  $X_w$  (for  $X_s$ ) if  $\forall x \in X$ ,  $\exists! \{\lambda_n\}_{n \in N} \subset K : \sum_{n=1}^{\infty} \lambda_n x_n = x$  in  $X_w$  (in  $X_s$ ).

We will also need the following concept.

**Definition 2.8.** System  $\{x_n\}_{n \in N} \subset X$  is called nondegenerate, if  $x_n \neq 0$ ,  $\forall n \in N$ .

### 3 Main results

**3.1. Space of coefficients.** Let  $X$  be an IFNS and let  $\{x_n\}_{n \in N} \subset X$  be some system. Assume that

$$\mathcal{H}_{\bar{x}}^w \equiv \left\{ \{\lambda_n\}_{n \in N} \subset C : \sum_{n=1}^{\infty} \lambda_n x_n \text{ converges in } X_w \right\};$$

$$\mathcal{H}_{\bar{x}}^s \equiv \left\{ \{\lambda_n\}_{n \in N} \subset C : \sum_{n=1}^{\infty} \lambda_n x_n \text{ converges in } X_s \right\}.$$

It is not difficult to see that,  $\mathcal{H}_{\bar{x}}^w$  and  $\mathcal{H}_{\bar{x}}^s$  are linear spaces with regard to component-specific summation and component-specific multiplication by a scalar. Take  $\forall \lambda \equiv \{\lambda_n\}_{n \in N} \in \mathcal{H}_{\bar{x}}^w$  and assume

$$\mu_K(\bar{\lambda}; t) = \inf_m \mu \left( \sum_{n=1}^m \lambda_n x_n; t \right); \nu_K(\bar{\lambda}; t) = \sup_m \nu \left( \sum_{n=1}^m \lambda_n x_n; t \right).$$

Let's show that  $\mu_K$  and  $\nu_K$  satisfy the conditions 1)-11).

1) It is clear that  $\mu_K(\bar{\lambda}; t) = 0, \forall t \leq 0$ .

2) Let  $\mu_K(\bar{\lambda}; t) = 1, \forall t > 0$ . Hence,  $\mu(\sum_{n=1}^m \lambda_n x_n; t) = 1, \forall m \in N, \forall t > 0$ .

Suppose that the system  $\{x_n\}_{n \in N}$  is nondegenerate. It follows from the above-stated relations that for  $m = 1$  we have  $\mu(\lambda_1 x_1; t) = 1, \forall t > 0$ . Hence,  $\lambda_1 x_1 = 0 \Rightarrow \lambda_1 = 0$ . Continuing this process, we get at the end of this process that  $\lambda_n = 0, \forall n \in N$ , i.e.  $\bar{\lambda} = 0$ .

3) The validity of relation  $\mu_K(A\bar{\lambda}; t) = \mu_K(\bar{\lambda}; \frac{t}{|c|}), \forall c \neq 0$ , is beyond any doubt.

4) As  $\mu(x; \cdot)$  is a non-degenerate function on  $R$ , it is not difficult to see that  $\mu_K(\bar{\lambda}; \cdot)$  has the same property. Let's show that  $\lim_{t \rightarrow \infty} \mu_K(\bar{\lambda}; t) = 1$ . Take  $\forall \varepsilon > 0$ . It is clear that  $\exists t_0 > 0 : \mu(S; t_0) \geq 1 - \varepsilon$ . Let  $S_m = \sum_{n=1}^m \lambda_n x_n$  and  $w\text{-}\lim_{m \rightarrow \infty} S_m = S \in X_w$ . Then it follows from the definition of  $\lim_w$  that  $\exists m_0(\varepsilon; t_0) : \mu(S_m - S; t_0) \geq 1 - \varepsilon, \forall m \geq m_0(\varepsilon; t_0)$ . Property 4. implies

$$\mu(S_m; 2t_0) = \mu(S_m - S + S; t_0 + t_0) \geq \min\{\mu(S_m - S; t_0); \mu(S; t_0)\}.$$

As a result we get

$$\mu(S_m; t_0) \geq 1 - \varepsilon, \forall m \geq m_0(\varepsilon; t_0). \tag{1}$$

As  $\mu(x; \cdot)$  is a non-decreasing function of  $t$ , it follows from (1) that

$$\mu(S_m; t) \geq 1 - \varepsilon, \forall m \geq m_0(\varepsilon; t_0), \forall t \geq t_0. \tag{2}$$

We have

$$\mu_K(\bar{\lambda}; t) = \inf_m \mu(S_m; t) = \min\left\{\mu(S_1; t); \dots; \mu(S_{m_0-1}; t); \inf_{m \geq m_0} \mu(S_m; t)\right\}, \tag{3}$$

where  $m_0 = m_0(\varepsilon; t_0)$ . As  $\lim_{t \rightarrow \infty} \mu(S_k; t) = 1$  for  $\forall k \in N, \exists t_k(\varepsilon); \forall t \geq t_k(\varepsilon) : \mu(S_k; t) \geq 1 - \varepsilon, k = \overline{1, m_0 - 1}$ . Let  $t_\varepsilon^0 = \max\{t_k(\varepsilon), k = \overline{1, m_0 - 1}\}$ . Then it is clear that

$$\mu(S_k; t) \geq 1 - \varepsilon, \forall t \geq t_\varepsilon^0. \tag{4}$$

It follows from (2) and (3) that

$$\inf_{m \geq m_0} \mu(S_m; t) \geq 1 - \varepsilon, \forall t \geq t_0.$$

Let  $t_\varepsilon = \max\{t_0; t_\varepsilon^0\}$ . Hence we obtain from (3) and (4)

$$\mu_K(\bar{\lambda}; t) \geq 1 - \varepsilon, \forall t \geq t_\varepsilon.$$

Thus,  $\lim_{t \rightarrow \infty} \mu_K(\bar{\lambda}; t) = 1, \forall \bar{\lambda} \in \mathcal{K}_{\bar{x}}^w$ .

5) Let  $\bar{\lambda}, \bar{\mu} \in \mathcal{K}_{\bar{x}}^w$  ( $\bar{\lambda} \equiv \{\lambda_n\}_{n \in N}; \bar{\mu} \equiv \{\mu_n\}_{n \in N}$ ) and  $s, t \in R$ . We have

$$\begin{aligned} \mu_K(\bar{\lambda} + \bar{\mu}; s + t) &= \inf_m \mu \left( \sum_{n=1}^m (\lambda_n + \mu_n) x_n; s + t \right) = \inf_m \mu \left( \sum_{n=1}^m \lambda_n x_n + \right. \\ &\quad \left. + \sum_{n=1}^m \mu_n x_n; s + t \right) \geq \inf_m \min \left\{ \mu \left( \sum_{n=1}^m \lambda_n x_n; s \right); \mu \left( \sum_{n=1}^m \mu_n x_n; t \right) \right\} = \\ &= \min \left\{ \inf_m \mu \left( \sum_{n=1}^m \lambda_n x_n; s \right); \inf_m \mu \left( \sum_{n=1}^m \mu_n x_n; t \right) \right\} = \\ &= \min \{ \mu(\bar{\lambda}; s); \mu(\bar{\mu}; t) \}. \end{aligned}$$

6) As  $\nu(x; t) = 1, \forall t \leq 0$ , it is clear that  $\nu_w(\bar{\lambda}; t) = 1, \forall t \leq 0, \forall \bar{\lambda} \in \mathcal{K}_{\bar{x}}^w$ .

7) Let the system  $\{x_n\}_{n \in N}$  be nondegenerate. Assume that  $\nu_K(\bar{\lambda}; t) = 0, \forall t > 0$ . Then  $\nu(\sum_{n=1}^m \lambda_n x_n; t) = 0, \forall t > 0, \forall m \in N$ . For  $m = 1$  we have  $\nu(\lambda_1 x_1; t) = 0, \forall t > 0 \Rightarrow \lambda_1 x_1 = 0 \Rightarrow \lambda_1 = 0$ . Continuing this process, we get  $\lambda_n = 0, \forall n \in N \Rightarrow \bar{\lambda} = 0$ .

8) Clearly,  $\nu_K(c\bar{\lambda}; t) = \nu_K(\bar{\lambda}; \frac{t}{|c|}), \forall c \neq 0$ .

9) It follows from the property 9. that  $\nu(x; \cdot)$  is a non-increasing function on  $R$ . Therefore,  $\nu_K(\bar{\lambda}; \cdot)$  is a non-increasing function on  $R$ . Let us show that  $\lim_{t \rightarrow \infty} \nu_K(\bar{\lambda}; t) = 0$ . Let  $S_m = \sum_{n=1}^m \lambda_n x_n$  and  $w\text{-}\lim_{m \rightarrow \infty} S_m = S \in X$ . Take  $\forall \varepsilon > 0$ . It is clear that  $\exists t_0 > 0 : \nu(S; t_0) \leq \varepsilon$ . Then it follows from the definition of  $\lim_w$  that  $\exists m_0 = m_0(\varepsilon; t_0) : \nu(S_m - S; t_0) \leq \varepsilon, \forall m \geq m_0$ . We have

$$\begin{aligned} \nu(S_m; t_0) &= \nu(S_m - S + S; t_0 + t_0) \leq \\ &\leq \max \{ \nu(S_m - S; t_0); \nu(S; t_0) \} \leq \varepsilon, \forall m \geq m_0. \end{aligned}$$

As  $\nu(x; \cdot)$  is a non-increasing function, it is clear that

$$\nu(S_m; t) \leq \varepsilon, \forall m \geq m_0, \forall t \geq t_0. \quad (5)$$

We have

$$\nu_K(\bar{\lambda}; t) = \sup_m \nu(S_m; t) = \max \left\{ \nu(S_1; t); \dots; \nu(S_{m_0-1}; t); \sup_{m \geq m_0} \nu(S_m; t) \right\}.$$

As  $\lim_{t \rightarrow \infty} \nu(S_k; t) = 0$  for  $\forall k \in N$ , we have  $\exists t_k(\varepsilon); \forall t \geq t_k(\varepsilon) : \nu(S_k; t) \leq \varepsilon, k = \overline{1, m_0 - 1}$ . Let  $t_\varepsilon^0 = \max \{ t_k(\varepsilon), k = \overline{1, m_0 - 1} \}$ . It is clear that  $\nu(S_k; t) \leq \varepsilon, \forall t \geq t_\varepsilon^0$ . It follows from (5) that  $\sup_{m \geq m_0} \nu(S_m; t) \leq \varepsilon, \forall t \geq t_0$ . Let

$t_\varepsilon = \max \{t_0; t_\varepsilon^0\}$ . Then it is clear that  $\nu_K(\bar{\lambda}; t) \leq \varepsilon, \forall t \geq t_\varepsilon \Rightarrow \lim_{t \rightarrow \infty} \nu_K(\bar{\lambda}; t) = 0$ .

10) Let  $\bar{\lambda}, \bar{\mu} \in \mathcal{K}_{\bar{x}}^w$  ( $\bar{\lambda} \equiv \{\lambda_n\}_{n \in N}; \bar{\mu} \equiv \{\mu_n\}_{n \in N}$ ) and  $s, t \in R$ . We have

$$\begin{aligned} \nu_K(\bar{\lambda} + \bar{\mu}; s + t) &= \sup_m \nu \left( \sum_{n=1}^m (\lambda_n + \mu_n) x_n; s + t \right) \leq \\ &\leq \sup_m \max \left\{ \nu \left( \sum_{n=1}^m \lambda_n x_n; s \right); \nu \left( \sum_{n=1}^m \mu_n x_n; t \right) \right\} = \\ &= \max \left\{ \sup_m \nu \left( \sum_{n=1}^m \lambda_n x_n; s \right); \sup_m \nu \left( \sum_{n=1}^m \mu_n x_n; t \right) \right\} = \\ &= \max \{ \nu_K(\bar{\lambda}; s); \nu_K(\bar{\mu}; t) \} . \end{aligned}$$

11)  $\mu_K(\bar{\lambda}; t) + \nu_K(\bar{\lambda}; t) = \inf_m \mu(\sum_{n=1}^m \lambda_n x_n; t) + \sup_m \nu(\sum_{n=1}^m \lambda_n x_n; t) \leq$

$$\leq \sup_m \left[ \mu \left( \sum_{n=1}^m \lambda_n x_n; t \right) + \nu \left( \sum_{n=1}^m \lambda_n x_n; t \right) \right] \leq 1, \forall \bar{\lambda} \in \mathcal{K}_{\bar{x}}^w, \forall \lambda \in R.$$

Thus, we have proved the validity of the following

**Theorem 3.1.** *Let  $(X; \mu; \nu)$  be a fuzzy normed space and let  $\{x_n\}_{n \in N} \subset X$  be a nondegenerate system. Then the space of coefficients  $(\mathcal{K}_{\bar{x}}^w; \mu_K; \nu_K)$  is also weakly fuzzy normed space.*

The following theorem is proved in absolutely the same way.

**Theorem 3.2.** *Let  $(X; \mu; \nu)$  be a fuzzy normed space and let  $\{x_n\}_{n \in N} \subset X$  be a nondegenerate system. Then the space of coefficients  $(\mathcal{K}_{\bar{x}}^s; \mu_K; \nu_K)$  is also strongly fuzzy normed space.*

**3.2. Completeness of the space of coefficients.** Subsequently, we assume that  $(X; \mu; \nu)$  is IFBS. Let us show that  $(\mathcal{K}_{\bar{x}}^w; \mu_K; \nu_K)$  is a strongly fuzzy complete normed space. First we prove the following

**Lemma 3.3.** *Let  $x_0 \neq 0, x_0 \in X$ , and let  $\{\lambda_n\}_{n \in N} \subset R$  be some sequence. If  $w\text{-}\lim_{n \rightarrow \infty} (\lambda_n x_0) = 0$ , i.e. for  $\forall \varepsilon > 0, \exists n_0 = n_0(\varepsilon; t): \mu(\lambda_n x_0; t) > 1 - \varepsilon$  ( $\nu(\lambda_n x_0; t) < \varepsilon$ ),  $\forall n \geq n_0$ ; then  $\lambda_n \rightarrow 0, n \rightarrow \infty$ .*

**Proof.** As  $x_0 \neq 0$ , it is clear that  $\exists t_0 > 0 : \mu(x_0; t_0) < 1$  (it follows from the property 1). If  $\lambda_n \neq 0$  we have  $\mu(\lambda_n x_0; t) = \mu\left(x_0; \frac{t}{|\lambda_n|}\right)$ , for  $\forall t > 0$ . Let the sequence  $\{\lambda_n\}_{n \in N}$  not convergent to zero. Then  $\exists \{\lambda_{n_k}\}_{k \in N}$  and  $\exists \delta > 0 :$

$|\lambda_{n_k}| \geq \delta, \forall k \in N \Rightarrow \frac{t}{|\lambda_{n_k}|} \leq \frac{t}{\delta}$ . As  $\mu(x_0; \cdot)$  is a non-decreasing function of  $t$ , then  $\mu\left(x_0; \frac{t}{|\lambda_{n_k}|}\right) \leq \mu\left(x_0; \frac{t}{\delta}\right), \forall t \in R_+$ . Take  $\tilde{t} = \delta t_0$ . We have  $\mu(\lambda_{n_k} x_0; \tilde{t}) \leq \mu(x_0; t_0) < 1, \forall k \in N$ . So we came upon a contradiction which proves the Lemma.

In the further we assume that the following condition is also fulfilled.

12) The functions  $\mu(x; \cdot), \nu(x; \cdot) : R_+ \rightarrow [0, 1]$  are continuous for  $\forall x \in X$ .

Take  $w$ -fundamental sequence  $\{\bar{\lambda}_n\}_{n \in N} \subset \mathcal{H}_{\bar{x}}^w, \bar{\lambda}_n \equiv \left\{ \lambda_k^{(n)} \right\}_{k \in N}$ . Then,

$\lim_{n, m \rightarrow \infty} \mu_K(\bar{\lambda}_n - \bar{\lambda}_m; t) = 1, \forall t \in R$ , i.e.

$$\lim_{n, m \rightarrow \infty} \inf_r \mu\left(\sum_{k=1}^m (\lambda_k^{(n)} - \lambda_k^{(m)}) x_k; t\right) = 1, \forall t \in R.$$

Take  $\forall k_0 \in N$  and fix it. We have

$$\left(\lambda_{k_0}^{(n)} - \lambda_{k_0}^{(m)}\right) x_{k_0} = \sum_{k=1}^{k_0} \left(\lambda_k^{(n)} - \lambda_k^{(m)}\right) x_k - \sum_{k=1}^{k_0-1} \left(\lambda_k^{(n)} - \lambda_k^{(m)}\right) x_k.$$

Then from property 5) we get

$$\mu\left(\left(\lambda_{k_0}^{(n)} - \lambda_{k_0}^{(m)}\right) x_{k_0}; t\right) \geq \min \left\{ \mu\left(\sum_{k=1}^{k_0} \left(\lambda_k^{(n)} - \lambda_k^{(m)}\right) x_k; \frac{t}{2}\right); \mu\left(\sum_{k=1}^{k_0-1} \left(\lambda_k^{(n)} - \lambda_k^{(m)}\right) x_k; \frac{t}{2}\right) \right\}.$$

It follows directly from this relation that

$$\lim_{n, m \rightarrow \infty} \mu\left(\left(\lambda_{k_0}^{(n)} - \lambda_{k_0}^{(m)}\right) x_{k_0}; t\right) = 1, \forall t \in R.$$

As  $x_{k_0} \neq 0$ , Lemma 3.3 implies  $\lim_{n, m \rightarrow \infty} \left| \lambda_{k_0}^{(n)} - \lambda_{k_0}^{(m)} \right| = 0$ , i.e. the sequence  $\left\{ \lambda_{k_0}^{(n)} \right\}_{n \in N}$  is fundamental in  $R$ . Let  $\lambda_{k_0}^{(n)} \rightarrow \lambda_{k_0}$ , as  $n \rightarrow \infty$ . Denote  $\bar{\lambda} \equiv \left\{ \lambda_n \right\}_{n \in N}$ . Let us show that  $\lim_{n \rightarrow \infty} \mu_K(\bar{\lambda}_n - \bar{\lambda}; t) = 1, \forall t \in R$ . Take  $\forall \varepsilon > 0, \forall t > 0$ . It is clear that

$$\exists n_0 = n_0(\varepsilon; t) : \mu_K(\bar{\lambda}_n - \bar{\lambda}_{n+p}; t) > 1 - \varepsilon, \forall n \geq n_0, \forall p \in N.$$

Consequently

$$\inf_r \mu\left(\sum_{k=1}^r \left(\lambda_k^{(n)} - \lambda_k^{(n+p)}\right) x_k; t\right) > 1 - \varepsilon, \forall n \geq n_0, \forall p \in N.$$



Hence

$$\mu \left( \sum_{k=1}^r \left( \lambda_k^{(n)} - \lambda_k^{(n+p)} \right) x_k; t \right) > 1 - \varepsilon, \forall n \geq n_0, \forall r, p \in N. \quad (6)$$

As shown above,  $\lim_{n, m \rightarrow \infty} \mu \left( \left( \lambda_k^{(n)} - \lambda_k^{(m)} \right) x_k; t \right) = 1, \forall t \in R$ . Let us show that  $\lim_{m \rightarrow \infty} \mu \left( \lambda_k^{(m)} x_k; t \right) = \mu \left( \lambda_k x_k; t \right), \forall t \in R_+$ . Indeed, if  $\lambda_k = 0$ , then  $\mu \left( 0; t \right) = 1, \forall t \in R_+$ , and, clearly,  $\lim_{m \rightarrow \infty} \mu \left( \lambda_k^{(m)} x_k; t \right) = 1$ , for  $\forall t \in R_+$ . If  $\lambda_k \neq 0$ , then for sufficiently large values of  $m$  we have  $\lambda_k^{(m)} \neq 0$ , and as a result

$$\mu \left( \lambda_k^{(m)} x_k; t \right) = \mu \left( x_k; \frac{t}{|\lambda_k^{(m)}|} \right) \xrightarrow{m \rightarrow \infty} \mu \left( x_k; \frac{t}{|\lambda_k|} \right) = \mu \left( \lambda_k x_k; t \right), \forall t \in R_+.$$

Passage to the limit in the inequality (6) as  $p \rightarrow \infty$  yields

$$\mu \left( \sum_{k=1}^r \left( \lambda_k^{(n)} - \lambda_k \right) x_k; t \right) \geq 1 - \varepsilon, \forall n \geq n_0, \forall r \in N. \quad (7)$$

We have

$$\begin{aligned} \mu \left( \sum_{k=r}^{r+p} \left( \lambda_k^{(n)} - \lambda_k \right) x_k; t \right) &= \mu \left( \sum_{k=1}^{r+p} \left( \lambda_k^{(n)} - \lambda_k \right) x_k - \sum_{k=1}^{r-1} \left( \lambda_k^{(n)} - \lambda_k \right) x_k; t \right) \geq \\ &\geq \min \left\{ \mu \left( \sum_{k=1}^{r+p} \left( \lambda_k^{(n)} - \lambda_k \right) x_k; \frac{t}{2} \right); \mu \left( \sum_{k=1}^{r-1} \left( \lambda_k^{(n)} - \lambda_k \right) x_k; \frac{t}{2} \right) \right\} \geq \\ &\geq 1 - \varepsilon, \forall n \geq n_0, \forall r, p \in N. \end{aligned}$$

As  $\bar{\lambda}_n \in \mathcal{K}_{\bar{x}}^w$ , it is clear that  $\exists m_0^{(n)} : \forall m \geq m_0^{(n)}, \forall p \in N$ :

$$\mu \left( \sum_{k=m}^{m+p} \lambda_k^{(n)} x_k; t \right) > 1 - \varepsilon.$$

Consequently

$$\begin{aligned} \mu \left( \sum_{k=m}^{m+p} \lambda_k x_k; t \right) &= \mu \left( \sum_{k=m}^{m+p} \left( \lambda_k - \lambda_k^{(n)} \right) x_k + \sum_{k=m}^{m+p} \lambda_k^{(n)} x_k; t \right) \geq \\ &\geq \min \left\{ \mu \left( \sum_{k=m}^{m+p} \left( \lambda_k - \lambda_k^{(n)} \right) x_k; \frac{t}{2} \right); \mu \left( \sum_{k=m}^{m+p} \lambda_k^{(n)} x_k; \frac{t}{2} \right) \right\} \geq \end{aligned}$$

$$\geq 1 - \varepsilon, \forall m \geq m_0^{(n)}, \forall p \in N.$$

It follows that the series  $\sum_{k=1}^{\infty} \lambda_k x_k$  is weakly fuzzy convergent in  $X_w$ , i.e.  $\exists w\text{-}\lim_{m \rightarrow \infty} \sum_{k=1}^m \lambda_k x_k$ . Consequently,  $\bar{\lambda} \in \mathcal{K}_{\bar{x}}^w$  and the relation (7) implies that  $\lim_{n \rightarrow \infty} \mu_K(\bar{\lambda}_n - \bar{\lambda}; t) = 1, \forall t \in R_+$ . It can be proved in similar way that  $\lim_{n \rightarrow \infty} \nu_K(\bar{\lambda}_n - \bar{\lambda}; t) = 0, \forall t \in R_+$ . As a result we obtain that the space  $(\mathcal{K}_{\bar{x}}^w; \mu_K; \nu_K)$  is weakly fuzzy complete. Thus, we have proved the following

**Theorem 3.4.** *Let  $(X; \mu; \nu)$  be a fuzzy Banach space with condition 12) and let  $\{x_n\}_{n \in N} \subset X$  be a nondegenerate system. Then the space of coefficients  $(\mathcal{K}_{\bar{x}}^w; \mu_K; \nu_K)$  is a weakly fuzzy complete normed space.*

Consider operator  $T : \mathcal{K}_{\bar{x}}^w \rightarrow X$  defined by

$$T\bar{\lambda} = \sum_{n=1}^{\infty} \lambda_n x_n, \bar{\lambda} \equiv \{\lambda_n\}_{n \in N} \in \mathcal{K}_{\bar{x}}^w.$$

Let  $w\text{-}\lim_{n \rightarrow \infty} \bar{\lambda}_n = \bar{\lambda}$  in  $\mathcal{K}_{\bar{x}}^w$ , where  $\bar{\lambda}_n \equiv \{\lambda_k^{(n)}\}_{k \in N} \in \mathcal{K}_{\bar{x}}^w$ . We have

$$\begin{aligned} \mu(T\bar{\lambda}_n - T\bar{\lambda}; t) &= \mu\left(\sum_{k=1}^{\infty} (\lambda_k^{(n)} - \lambda_k) x_k; t\right) \geq \\ &\geq \inf_m \mu\left(\sum_{k=1}^m (\lambda_k^{(n)} - \lambda_k) x_k; t\right) = \mu(\bar{\lambda}_n - \bar{\lambda}; t). \end{aligned}$$

It follows directly that  $w\text{-}\lim_{n \rightarrow \infty} T\bar{\lambda}_n = T\bar{\lambda}$ , i.e. the operator  $T$  is weakly fuzzy continuous. Let  $\bar{\lambda} \in \text{Ker}T$ , i.e.  $T\bar{\lambda} = 0 \Rightarrow \sum_{n=1}^{\infty} \lambda_n x_n = 0$ , where  $\bar{\lambda} \equiv \{\lambda_n\}_{n \in N} \in \mathcal{K}_{\bar{x}}^w$ . It is clear that if the system  $\{x_n\}_{n \in N}$  is *w-linearly independent*, then  $\lambda_n = 0, \forall n \in N$ , and, as a result,  $\text{Ker}T = \{0\}$ . In this case  $\exists T^{-1} : \text{Im}T \rightarrow \mathcal{K}_{\bar{x}}^w$ . If, in addition,  $\text{Im}T$  is *w-closed* in  $X$ , then  $T^{-1}$  is also continuous.

Denote by  $\{\bar{e}_n\}_{n \in N} \subset \mathcal{K}_{\bar{x}}^w$  a canonical system in  $\mathcal{K}_{\bar{x}}^w$ , where  $\bar{e}_n = \{\delta_{nk}\}_{k \in N} \in \mathcal{K}_{\bar{x}}^w$ . Obviously,  $T\bar{e}_n = x_n, \forall n \in N$ . Let us prove that  $\{\bar{e}_n\}_{n \in N}$  forms an *w-basis* in  $\mathcal{K}_{\bar{x}}^w$ . Take  $\forall \bar{\lambda} \equiv \{\lambda_n\}_{n \in N} \in \mathcal{K}_{\bar{x}}^w$  and show that the series  $\sum_{n=1}^{\infty} \lambda_n \bar{e}_n$  is weakly fuzzy convergent in  $\mathcal{K}_{\bar{x}}^w$ . In fact, the existence of  $w\text{-}\lim_{m \rightarrow \infty} \sum_{n=1}^m \lambda_n x_n$  in  $X_w$  implies that for  $\forall \varepsilon > 0$ , and  $\forall t > 0, \exists m_0 = m_0(\varepsilon; t) \in N$

$$\mu\left(\sum_{n=m}^{m+p} \lambda_n x_n; t\right) > 1 - \varepsilon, \forall m \geq m_0, \forall p \in N.$$

We have

$$\mu_K\left(\sum_{n=m}^{m+p} \lambda_n \bar{e}_n; t\right) = \inf_r \left(\sum_{n=m}^r \lambda_n x_n; t\right) \geq 1 - \varepsilon, \forall m \geq m_0, \forall p \in N.$$

It follows that the series  $\sum_{n=1}^{\infty} \lambda_n \bar{e}_n$  is weakly fuzzy convergent in  $\mathcal{K}_{\bar{x}}^w$ . Moreover

$$\begin{aligned} \mu_K \left( \bar{\lambda} - \sum_{n=1}^m \lambda_n \bar{e}_n; t \right) &= \mu_K (\{0; \dots; 0; \lambda_{m+1}; \dots\}; t) = \inf_r \mu \left( \sum_{n=m+1}^r \lambda_n x_n; t \right) \geq \\ &\geq 1 - \varepsilon, \forall m \geq m_0, \forall t \in R_+. \end{aligned}$$

Consequently,  $w\text{-}\lim_{m \rightarrow \infty} \sum_{n=1}^m \lambda_n \bar{e}_n = \bar{\lambda}$ , i.e.  $\bar{\lambda} \stackrel{w}{=} \sum_{n=1}^{\infty} \lambda_n \bar{e}_n$ . Consider the functionals  $e_n^*(\bar{\lambda}) = \lambda_n, \forall n \in N$ . Let us show that they are *w-continuous*. Let  $w\text{-}\lim_{n \rightarrow \infty} \bar{\lambda}_n = \bar{\lambda}$ , where  $\bar{\lambda}_n \equiv \left\{ \lambda_k^{(n)} \right\}_{k \in N} \in \mathcal{K}_{\bar{x}}^w$ . As established in the proof of Theorem 3, we have  $\lambda_k^{(n)} \rightarrow \lambda_k, n \rightarrow \infty$ , i.e.  $e_k^*(\bar{\lambda}_n) \rightarrow e_k^*(\bar{\lambda})$ , as  $n \rightarrow \infty$ , for  $\forall k \in N$ . Thus,  $e_k^*$  is *w-continuous* in  $\mathcal{K}_{\bar{x}}^w$  for  $\forall k \in N$ . On the other hand, it is easy to see that  $e_n^*(\bar{e}_k) = \delta_{nk}, \forall n, k \in N$ , i.e.  $\{e_n^*\}_{n \in N}$  is *w-biorthogonal* to  $\{\bar{e}_n\}_{n \in N}$ . As a result we obtain that the system  $\{\bar{e}_n\}_{n \in N}$  forms an *w-basis* in  $\mathcal{K}_{\bar{x}}^w$ . So we get the validity of the following

**Theorem 3.5.** *Let  $(X; \mu; \nu)$  be a fuzzy Banach space with condition 12) and let  $\{x_n\}_{n \in N} \subset X$  be a nondegenerate system. Then the corresponding space of coefficients  $(\mathcal{K}_{\bar{x}}^w; \mu_K; \nu_K)$  is weakly fuzzy complete with canonical *w-basis*  $\{\bar{e}_n\}_{n \in N}$ .*

Suppose that the system  $\{x_n\}_{n \in N}$  is *w-linearly independent* and  $ImT$  is closed. Then it is easily seen that  $\{x_n\}_{n \in N}$  forms an *w-basis* in  $ImT$ , and, in case of its *w-completeness* in  $X_w$ , it forms an *w-basis* for  $X_w$ . In this case,  $\mathcal{K}_{\bar{x}}^w$  and  $X_w$  are isomorphic, and  $T$  is an isomorphism between them. The opposite of it is also true, i.e. if the above-defined operator  $T$  is an isomorphism between  $\mathcal{K}_{\bar{x}}^w$  and  $X_w$ , then the system  $\{x_n\}_{n \in N}$  forms an *w-basis* in  $X_w$ . We will call  $T$  a coefficient operator. Thus, the following theorem holds.

**Theorem 3.6.** *Let  $(X; \mu; \nu)$  be a fuzzy Banach space with condition 12),  $\{x_n\}_{n \in N} \subset X$  be a nondegenerate system,  $(\mathcal{K}_{\bar{x}}^w; \mu_K; \nu_K)$  be a corresponding weakly fuzzy complete normed space and  $T : \mathcal{K}_{\bar{x}}^w \rightarrow X_w$ , be a coefficient operator. System  $\{x_n\}_{n \in N}$  forms an *w-basis* for  $X_w$  if and only if the operator  $T$  is an isomorphism between  $\mathcal{K}_{\bar{x}}^w$  and  $X_w$ .*

## 4 Conclusion

Thus we arrive at the following conclusion: Let 5-tuple  $(X; \mu; \nu; *; \diamond)$  be an IFB<sub>w</sub>S. Then:

1. this structure generates the corresponding concepts for the theory of approximation;

2. every nondegenerate system  $\{x_n\}_{n \in N} \subset X$  generates a corresponding weakly fuzzy complete normed space  $(\mathcal{K}_{\bar{x}}^w; \mu_K; \nu_K)$  of coefficients;
3. canonical system  $\{\bar{e}_n\}_{n \in N}$  forms a weak basis for  $\mathcal{K}_{\bar{x}}^w$  ;
4. system  $\{x_n\}_{n \in N}$  generates a coefficient operator  $T : \mathcal{K}_{\bar{x}}^w \rightarrow X$  ;
5. system  $\{x_n\}_{n \in N}$  forms a weak basis for  $X$  if and only if  $T$  is an isomorphism between  $\mathcal{K}_{\bar{x}}^w$  and  $X$  .

Note that many results of this work are new in classical case, too.

## References

- [1] Singer I. Bases in Banach spaces. v.1, *Springer*, 1970, 673 p.
- [2] Heil Ch. A Basis Theory Primer. *Springer*, 2011, 536 p.
- [3] Bilalov B.T., Najafov T.I. On the approximation properties of some systems in Banach spaces. *Int. Journal of Math. Analysis*, v.5, 2011, No19, pp.923-933.
- [4] Dremin I.M., Ivanov O.V., Nechitailo V.A. Wavelets and their uses, *Usp. physics. nauk*, 2001, v.171, No 5, pp. 465-501.
- [5] Charles K. Chui. An introduction to wavelets. *Academic Press*, Jan 3, 1992, 266 p.
- [6] El Naschie M.S . On the uncertainty of Cantorian geometry and the two slit experiment, *Chaos, Solitons and Fractals*, 9(3), 1998, pp. 517-529.
- [7] El Naschie M.S. On fuzzy Kahler-like manifold which is consistent with the two-slit experiment, *Int. J. Nonlinear Sci. Numer. Simul.* 6 (2), 2005, pp.95-98.
- [8] El Naschie M.S. A review of E-infinity theory and the mass spectrum of high energy particle physics, *Chaos, Solitons & Fractals*, 19 (1), 2004, pp. 209–236.
- [9] Tanaka Y., Mizuno Y., Kado T. Chaotic dynamics in the Friedman equation, *Chaos, Solitons & Fractals*, 24 ,2005, pp. 407–422.
- [10] Mursaleen M., Karakaya V., Mohiuddine S.A. Schauder Basis, Separability, and Approximation Property in Intuitionistic Fuzzy Normed Space, *Abstract and Applied Analysis*, 2010, Article ID 131868, 14 pages.

- [11] Yılmaz Y. Schauder bases and approximation property in fuzzy normed spaces, *Computers & Mathematics with Applications*, vol. 59, no. 6, 2010, pp. 1957–1964.
- [12] Debnath P., Sen M. A Remark on the Separability and Approximation Property in Intuitionistic Fuzzy  $n$ -normed Linear Spaces, *International Mathematical Forum*, vol.6, No.39, 2011, pp. 1933-1940.
- [13] Bag T., Samanta S.K. Fuzzy bounded linear operators. *Fuzzy Sets and Systems*, vol. 151, 2005, pp. 513-547.
- [14] Schweizer B., Sklar A. Statistical metric spaces. *Pacific J. Math.* 10, 1960, pp. 313–334.
- [15] Saadati R., Park J.H. On the intuitionistic fuzzy topological spaces. *Chaos, Solitons & Fractals*, 27, 2006, pp.331-344.
- [16] Kramosil I., Michelek J. Fuzzy metric and statistical metric spaces. *Kybernetika*, vol. 11, 1975, pp. 326-334.
- [17] Deng Z.K. Fuzzy pseudo metric spaces. *J. Math. Anal. Appl.*, vol. 86, No. 1, 1982, pp. 74-95.
- [18] Erceg M.A. Metric spaces in fuzzy set theory. *J. Math. Anal. Appl.*, 69 1979, pp. 205–230.
- [19] Park J. H. Intuitionistic fuzzy metric spaces. *Chaos, Solitons and Fractals*, 22, 2004, pp. 1039– 1046.
- [20] Saadati R., Vaezpour S.M. Some results on fuzzy Banach spaces. *J. Appl. Math. Comput.*, 17 , 2005, pp. 475–484.
- [21] Lowen R. Fuzzy Set Theory: Basic Concepts, Techniques and Bibliography. *Kluwer Academic Publishers*, 1996, 408 p.
- [22] Megginson R.E. An Introduction to Banach Space Theory. Graduate Texts in Mathematics, 183, *Springer, New York*, 1998, XIX + 596 pp.
- [23] Bilalov B.T., Farahani S.M., Guliyeva F.A. The Intuitionistic Fuzzy Normed Space of Coefficients. *Abstract and Applied Analysis, Hindawi Publishing Corporation*, Volume 2012, Article ID 969313, 11 pages doi:10.1155/2012/969313
- [24] Bilalov B.T., Guliyeva F.A. On the approximation property system in statistical type metric space. *Pure Mathematical Sciences*, Vol 1, 2012, no 4, 151-164

- [25] Guliyeva F.A. On the space of coefficients in the intuitionistic fuzzy normed space. *Caspian Journal of Applied Mathematics, Ecology and Economics*, vol 1, no 2 (2013), pp.61-73
- [26] Bilalov B.T., Guliyeva F.A. On the basicity property system in the intuitionistic Fuzzy metric space. *Azerbaijan Journal of Mathematics*, VOL 4, NO 1 (2014), pp.136-149
- [27] Bilalov B.T., Guliyeva F.A. The Weakly Basicity of System in the Intuitionistic Fuzzy Metric Space. *Advances in Fuzzy Mathematics*. ISSN 0973-533X Volume 9, Number 1 (2014), pp. 33-45

**Received: April, 2015**