



Baseline Study on the Occurrence of *Cryptosporidium Spp* from Streams Water, After Torrential Rains in Bamenda, Cameroon

¹Polycarp N Chia*, ²Prof. Chinyere Ukaga N, ³Prof. Kenneth A Yongabi, ⁴Prof. BEB Nwoke, & ⁵Prof. PM Tih

^{1a}Department of Parasitology, Imo State University Owerri, Nigeria. PO Box, 2000 Owerri, Imo state Nigeria

^{1b}Department of Biochemistry Catholic University of Cameroon (CATUC) Bamenda P.O Box, 782 Mankon Bamenda - Cameroon. Website: www.catuc.org

²Department of Animal and Environmental Biology PO Box, 2000 Owerri, Imo State University Nigeria.

^{3a}Phyto-Biotechnology Research Foundation Institute, P.O Box 921 Bamenda North West Region-Cameroon

^{3b} Catholic University of Cameroon, Bamenda, Cameroon(CATUC) P.O Box, 782 Mankon Bamenda, North west Region Cameroon

⁴Department of Animal and Environmental Biology, PO Box, 2000 Owerri, Imo State University Nigeria

⁵Cameroon Baptist Convention Health Board, Bamenda Cameroon PO Box, 01.Nkwen Bamenda Cameroon. Website: www.cbhealthservice.org

Abstract

Oocysts of Cryptosporidia are zoonotic infective stages of *Cryptosporidia spp* that can easily contaminate the environment if not properly disposed of. Torrential rains would usually wash them into the nearby streams and likely be consumed by humans when the water is used. Cryptosporidiosis in people with suppressed immune systems have no cure as the robust oocyst are not destroyed by chlorination. The aim of this study is to determine the presence of Cryptosporidia oocysts in stream water after torrential rains

60 samples of stream water were collected immediately after rain storm in June and July 2014, from 5 streams flowing through Bamenda town. 2 samples of 1.5L were collected from each stream, on 6 different days. *Cryptosporidia spp* were present in all the streams but there was no significant difference between their occurrence and the stream's location, Chi square = 2.605, >0.05

The streams flowing through Bamenda town are polluted

Key words: storm-water, zoonotic, Cryptosporidiosis, chlorination, oocysts

1. Introduction

Zoonotic pathogens of enteric origin are likely sources of contamination of the environment, with transmission via soil, agriculture, water and sediments (Bonetta *et al.*, 2011). Proceedings of the meeting entitled "*Cryptosporidium: from Molecules to Disease*" (2001) observed that there are many reports of this parasite being acquired from public water supplies. However, these may well represent cases of cryptosporidiosis transmitted to humans by domestic animals such as kittens and puppies, from ruminants on farms, or by contact with other humans. Food grown on soil fertilized with manure could also be considered a potential source of infection; however, contaminated water represents the major source of infections for humans. *Cryptosporidium* oocysts may remain viable in water for over 140 days (Hooda, *et al.*, 2000) and are very resistant to the most common disinfectants (Campbell *et al.*, 1982) making them difficult to destroy by conventional chlorination.

The public health importance of cryptosporidiosis has not been widely reported despite the growing interest of the slang "water is life" as there is no on-going monitoring of our water sources for contaminants. The Bamenda highlands is actually a water shed and our water is not safe as it is likely being contaminated with environmentally resistant Oocysts of *Cryptosporidia spp*. The method use in purifying our water(Chlorination) does not eliminated these Oocysts, and their infective dose is very low as only 1 to 10 Oocysts(Pereira *et al.*, 2000) can elicit an infection (cryptosporidiosis) which has adverse effects on people with weakened immune systems like HIV/AIDS patients, children, the elderly pregnant women and people on chemotherapy. At the moment in Bamenda the only dam supplying the Bamenda metropolis has no water and denizens move out of town to get water.

Cattle and sheep are grazing free range in this water catchment and during torrential rains, all cattle dung and sheep dropping are washed into the dam. Zero grazing is not practiced. To better understand the gravity of the contamination of our water, it is necessary to analyze faecal samples from animals grazing in water catchment areas as well as the water taking its rise from the water catchment area. Rural to urban migration is putting a lot of pressure on the scarce portable water. Some residents who have built closed to stream in areas that are not accessible have built their toilets close to stream. Pipes are connected to the toilets and when there are heavy rains, they will open pipes to let out faeces into the nearby streams figure 2Some gardeners down stream use this water to irrigate their vegetables. Previous studies in Cameroon by Ajeegah *et al.*, (2007) and Ajeegah, *et al.*, (2005) identified Cryptosporidia Oocysts in municipal lake in Yaounde-Cameroon. There have been a lot of studies on the biodynamic of the resistant forms of these emerging pathogens in water (Smith and Thompson, (2001), Ajeegah, *et al.*, 2005). Most of these analyses have been carried out on streams not on humans. Ajeegah, *et al.*, (2005) asserted that quantitative data on the identification of Oocysts in lake ecosystems and their survival in a lentic environment is still unavailable in developing countries in general and Cameroon in particular. The hydrosystem of the municipal lake in Yaounde was considered as a significant reservoir of Oocysts, that may be resuspended in the water column and give rise to sporadic increases in the Oocysts in the whole river basin in which is situated the lake, resulting in the outbreaks of Cryptosporidiosis as many gardeners along the streams emanating from this lake use the water for irrigation.

streams but the others from Up-station reach the foothill zone through a series of small waterfalls over bare hill slopes which, together with the hills, form the most striking feature of the Bamenda urban area. Since the streams are youthful, flow is rapid. Thus, debris and watershed runoff from the slope sub-system become part of the input of the stream channels. During the rainy season, the linear-dendritic streaming system (Figure 1), combined with various erosional processes such as river valleys, causes spillage, subjecting houses and properties, particularly those along stream courses, to periodic flooding during torrential rainstorms. It is during this rainstorm that some inhabitants near the streams open their toilets to discharge faeces into the streams.



Figure 2.0. Pipes projecting from a toilet into a nearby stream in New layout Nkwen . Picture by Polycarp Ndikvu July 2015 (Figure 2) and thick film, of stool settles after the rains (figure 3).



Figure 3.0. Film on the water below the pipes projecting from the toilet into the stream in New layout Nkwen. Picture by Polycarp Ndikvu, July 2015

Pig styles are also built along the stream banks and when rains fall the faeces are washed into the stream water.



Figure 4. Pigsty the along New Layout stream. Picture by Polycarp Ndikvu July 2015

Some gardeners use the water down stream to irrigate their vegetables (figure 5)



Garden along the Ayaba stream, Picture by Polycarp Ndikvu July 2015

2.1.3 Sample collection

A reconnaissance visit was made to human-water points along each of the streams flowing through Bamenda Metropolis, a week before sampling started.

Stream water was collected in duplicates of 1.5L sterile containers, after torrential rains from 5 streams running through Bamenda town. The stream water was collected immediately after 6 different storm events. The samples were transported immediately to the Phyto-Biotechnology Research Foundation (PRF) laboratory for analysis. A total of 60 samples were collected for the study

2.1.4 Laboratory analysis

Water samples were filtered using Sieve (strainer) with small holes, of 400–450 μ m in size. The small inexpensive nylon tea or coffee strainer was used to remove large particles. The method used to recovered oocysts and/or cysts was formol-ether concentration method as per (Ukaga *et al.*, 2002; Cheesbrough 2006).

A small amount of about 750 μ l of the concentrated suspension of each water source was drawn out. 2 smears were made on slides, air dried and fixed with methanol (Kinyoun) and the other observed under X10 and X40 (wet smear (Cheesbrough, 2009). Lugol's iodine was added for the identification cysts and ova.

The smears were stained with Modified Kinyoun's Acid-fast Stain (Cold) as described by Garcia, (2001)

Microscopic identification of the parasites in each was under magnification 10X and 40X objectives. Standard textbooks and pictures provided (Lindsay *et al.*, 2000) by the Phyto-Biotechnology Research Foundation Institute Bamenda, Cameroon, were used to established the identity of the parasites.

2.1.5 Statistical analysis

Data was entered into Ms Excel[®] 2003 (Microsoft corporation, USA) and analysis were conducted using SPSS for Windows version 12. Prevalence was calculated as a percentage of d/n where d is the number of animals infected and n = Total number of water samples analysed. Chi square test was used to explore the association between the occurrence of *Cryptosporidium* oocysts in a stream and the stream's location. Also a p- value of <0.05 was considered to be statistically significant.

3. Results

Of the 60 samples collected from the 5 streams, 12 samples were collected from each stream. The prevalence *Cryptosporidium spp* and *Giardia spp* was determined according to the different streams sampled. *Giardia spp* was included only because of its zoonotic potentials.

The occurrence of *Cryptosporidium spp* in Sisia Stream 3(25%)table 1a and 1b, New layout stream 4(33.33%) table 2a and 2b, Progressive Comprehensive high School stream(PCHS) 3(25%)table 3a and 3b, Ayaba stream 1(25%) table 4a and 4band Lourdes stream 4(33%)table 5a and 5b. New Layout stream was highly polluted 75% while Ayaba stream was the least polluted 25% table 4a. table 6 show the overall prevalence in all the streams.

The distribution of cryptosporidium across sample1 is the same as the distribution across sample 2 implying that the results for occurrence of cryptosporidium is not affected by the sample the collected.(results cannot be biased by sample). There is no significant association between the occurrence of *Cryptosporidium* oocysts in a stream and its location, Chi square = 2.605, >0.05

Table 1. Occurrence of parasites in Sisia stream

DAY	<i>Cryptosporidium spp</i>	<i>Giardia spp</i>
1	1	0
2	0	1
3	1	0
4	0	0
5	1	0
6	0	0
Total occurrence	3	1

Legend: 1= occurrence; 0 = absences

Table 1b. Prevalence of parasites in Sisia stream

Parasite	Occurrence	Prevalence (%)	Overall Prevalence (%)
<i>Cryptosporidium spp</i>	3	25	33.33%
<i>Giardia spp</i>	1	8.33	

Table 2. Occurrence of parasites in New Layout stream

DAY	<i>Cryptosporidium spp</i>	<i>Giardia spp</i>
1	0	1
2	1	0
3	1	1
4	1	1
5	1	1
6	0	1
Total occurrence	4	5

Table 2b Prevalence of parasites in PCHS stream

Parasite	Occurrence	Prevalence (%)	Overall Prevalence (%)
<i>Cryptosporidium spp</i>	4	33.33	75%
<i>Giardia spp</i>	5	41.67	

Table 3a. Occurrence of parasites in PCHS stream

DAY	<i>Cryptosporidium spp</i>	<i>Giardia spp</i>
1	0	1
2	0	0
3	1	1
4	0	1
5	1	1
6	1	0
Total occurrence	3	4

Table 3b Prevalence of parasites in PCHS stream

Parasite	Occurrence	Prevalence (%)	Overall Prevalence (%)
<i>Cryptosporidium spp</i>	3	25	58
<i>Giardia spp</i>	4	33	

Table 4. Occurrence of parasites in Ayaba stream

DAY	<i>Cryptosporidium spp</i>	<i>Giardia spp</i>
1	0	0
2	0	0
3	0	0
4	1	0
5	0	0
6	0	0
Total occurrence		

Table 4b Prevalence of parasites in Ayaba stream

Parasite	Occurrence	Prevalence (%)	Overall Prevalence (%)
<i>Cryptosporidium spp</i>	1	25	8.33%
<i>Giardia spp</i>	0	0	

Table 5. Occurrence of parasites in Lourdes Stream

DAY	<i>Cryptosporidium spp</i>	<i>Giardia spp</i>
1	1	0
2	1	0
3	1	1
4	0	0
5	1	1
6	0	1
Total occurrence		

Table 5b Prevalence of parasites in Lourdes stream

Parasite	Occurrence	Prevalence (%)	Overall Prevalence (%)
<i>Cryptosporidium spp</i>	4	33	57
<i>Giardia spp</i>	3	25	

Table .6 Occurrences and Prevalence of parasites in streams in Bamenda Metropolis

<i>Cryptosporidium spp</i> / <i>Giardia spp</i> positivity in Stream water in BAMENDA Town					
DAYS OF SAMPLE COLLECTION	SISIA STREAM	NEW LAYOUT STREAM	PCHS STREAM	AYABA STREAM	LOURDES STREAM
D1	1/0	0/1	0/1	0/0	1/0
D2	0/1	1/0	0/0	0/0	1/0
D3	1/0	1/1	1/1	0/0	1/1
D4	0/0	1/1	0/1	1/0	0/0
D5	1/0	1/1	1/1	0/0	1/1
D6	0/0	0/1	1/0	0/0	0/1
SUB-TOTAL	3/1	4/5	3/4	1/0	4/3
PREVALENCE	25%/8.33%	33.33%/41.67%	25%/33%	25%/0.0%	33%/25%
TOTAL	33.33%	75%	58%	8.33%	57%

Legend: The first figure in each stream represents sample 1 as well as *Cryptosporidium spp* while the second figure represents sample 2 as well as *Giardia spp*. 1= occurrence, 0= absence

4. Discussion /Conclusion

Epidemiological study carried out by Ntonifor *et al*, (2013) to determine the prevalence, intensity of infection gastrointestinal (GIT) parasites in grazing ruminants in Jakiri, Bui Division, North West Region of Cameroon, did not consider Cryptosporidiosis as being of any importance. Previous studies in Cameroon by Ajeegah *et al*, (2007) and Ajeegah, *et al*, (2005) identified Cryptosporidia Oocysts in municipal lake in Yaounde-Cameroon. However, the source of contamination was never traced. The present study is to determine the presence of oocysts in stream water and if the location of the streams has an influence in their occurrence; or the distribution of oocyst in the first sample is different from the second sample.

In conclusion there is urgent need fro an integrated approach to the detection of *Cryptosporidia spp* by veterinarians, molecular epidemiologists, and parasitologists in order to establish the dynamics of this parasite in Cameroon as this would lead to better understanding of zoonotic potentials of this parasite.

4.1 Recommendation:

All the flood plains in town could be reclaimed converted into public gardens and parks. Sanitary inspectors should move along the streams everyday to see that the people should not build along the streams or through sewage.

Reference

- Adekunle, B. Ayinmode and Benjamin, O. Fagbemi, (2010) "Prevalence of Cryptosporidium infection in cattle from South Western Nigeria," in VETERINARSKI ARHIV 80 (6), 723-731.
- Bennema, S.C., Vercruvse, J., Morgan, E., Stafford, K., Hoglund, J., Demeler, J., Von Samson-Himmelstierna, G., Charlier, J. (2010) "Epidemiology and risk factors for exposure to gastrointestinal nematodes in dairy herds in Northwestern Europe," in Vet. Parasitol. 173: 247-254,
- Biu A. A., A. Maimunatu A. F. Salamatu and E. T. Agbadu, (2009) "A faecal survey of gastrointestinal parasites of ruminants on the University of Maiduguri Research Farm," in *International J. of Biomedical and Health Science*, Vol.5 No.4.
- Bonetta, S., Borelli, E., Bonetta, S., Conio, O., Palumbo, F. & Carraro, E. (2011). Development of a PCR protocol for the detection of *Escherichia coli* O157:H7 and *Salmonella* spp. in surface water. *Environ Monit Assess* 177, 493–503.
- Campbell, I., A.S. Tzipori, G. Hutchison, K.W. Angus, (1982) "Effect of disinfectants on survival of cryptosporidium oocysts," in Vet. Rec. 111, 414–415.
- Current, W.L and Garcia L. S, (1991) "Cryptosporidiosis," in *Clin. Microbiol. Rev.* 4: 325–358,
- De Graaf D.C, Vanopdenbosch E, Ortega-Mora L.M, Abbassi H, and Peeters J.E. (1999) "A review of the importance of cryptosporidiosis in farm animals, in *Int J Parasitol.* 29(8):1269-87.
- Dorny, P., Stoliar, V., Charlier, J., Meas, S., Sorn, S., Chea, B., Holl, D., Van Aken, D., Vercruvse, J. (2011). Infections with gastrointestinal nematodes, *Fasciola* and *Paramphistomum* in cattle in Cambodia and their association with morbidity parameters," in Vet. Parasitol., 175: 293-199.
- Fayer, R., M. Santin and D. Macarasin, (2012) "Detection of concurrent infection of dairy cattle with *Blastocystis*, *Cryptosporidium*, *Giardia*, and *Enterocytozoon* by molecular and microscopic methods," in *Parasitol. Res.* 111(3): 1349-55,
- Fayer, R., M. Santín and J.M. Trout, (2008) "*Cryptosporidium ryanae* n. sp. (Apicomplexa: Cryptosporidiidae) in cattle (*Bos taurus*)," in Vet. Parasitol. 156(3-4): 191-8,
- Fayer, R., M. Santín, J.M. Trout and E. Greiner (2006) "Prevalence of species and genotypes of *Cryptosporidium* found in 1-2-year-old dairy cattle in the eastern United States," in *Vet. Parasitol.* 135(2): 105-12.
- Gountié Dedzo M. and Nono A. (2013) "Natural hazards along the Bamenda escarpment and its environs: The case of landslide, rock fall and flood risks (Cameroon volcanic line, North-West Region)," in *Glo. Adv. Res. J. Geol. Min.* Vol. 2(1) pp. 015-026..
- Guedjeo C.S., Kagou Dongmo A, Ngapgue F, Nkouathio D. G., Zangmo Tefogoum G, Hooda, P. S Edwards, A. C.; Anderson, H. A, Miller, A, (2011) "A review of water Inpankaew T., Phasuk C., Pinyopanuwat N., Chimnoi W., Kengradomkij C., Arunwipat P., and Anakewith T, "Prevalence of Gastro-Intestinal Parasites of Dairy Cows in Thailand. Kasetsart" in *J. Nat. Sci.*, 45: 40-45,
- Hooda P.S., A.C Edwards, H.A. Anderson, A. Miller, (2000) "A review of water quality concerns in livestock farming areas," in *Sci. Total Environ.* 250: 143–167, 2000 quality concerns in livestock farming areas," in *Science of total Environmental.* 250:143-167,
- Jongwutiwes, S., R. Tiangtip, S. Yentakarm, N. Chantachum, (2002) "Simple method for long-term copro-preservation of *Cryptosporidium* oocysts for morphometric and molecular analysis," in *Trop. Med. Int. Health* 7, 257-264
- Le Chevallier, M. W.; W. D. Norton; R. G Lee (1991) "Occurrence of *Giardia* and *Cryptosporidium* spp. in surface water supplies," *Applied & Environmental Microbiology*, vol. 57, pp. 2610-2616.
- Lindsay, D.S., S.J. Upton, D.S. Owens, U.M. Morgan, J.R. Mead and B.L. Blagburn, (2000) "*Cryptosporidium andersoni* n. sp. (Apicomplexa: Cryptosporidiidae) from cattle, *Bos Taurus*" in *J. Eukaryot. Microbiol.* 47(1): 91-5,
- MacKenzie, W.R.; N. J. Hoxie; M. E. Proctor; M. S. Gradus; K. A. Blair; D. F. Peterson; J. J. Kazmierczak; D. G. Addiss; K. R. Fox, J. B. Rose; J. P. Davis (1994) "A massive outbreak of *Cryptosporidium* infection transmitted through the public water supply," *New England Journal of Medicine*, vol. 331, pp. 161-167.
- Mallinath, R. H. K., P. G. Chikkachowdappa, A. K. J. Gowda, P. E.D'Souza, (2009) "Studies on the prevalence of cryptosporidiosis in bovines in organized dairy farms in and around Bangalore, South India" in Vet. arhiv 79:461-470,
- Monica Cheesbrough in: District Laboratory Practice in Tropical Countries Cambridge University Press. 2nd Edition 2009 available from www.cambridge.org/9780521676304
- Ntonifor H. N, Shei S. J, Ndaleh N. W, and Mbunkur G. N, (2013). "Epidemiological studies of gastrointestinal parasitic infections in ruminants in Jakiri, Bui Division, North West Region of Cameroon," in *Journal of Veterinary Medicine and Animal Health.* Vol. 5(12), pp. 344 -352,
- Pereira S.J., N.E. Ramirez, L. Xiao, L.A, (2002) "Ward, Pathogenesis of human and bovine *Cryptosporidium parvum* in gnotobiotic pigs," in *J. Infect. Dis.* 186: 715–718,
- Report of Infectious Disease Epidemiology Section Office of Public Health Louisiana Dept of Health & Hospitals (2011). Available from www.infectiousdisease.dhh.louisiana.gov
- Rinaldi, M., Dreesen, L., Hoorens, P.R., Li R.W., Claerebout, E., Goddeeris, B., Vercruvse, J. Van Den Broek, W., Geldhof, P, (2011) "Infection with gastrointestinal nematode *Ostertagia ostertagi* in cattle affects mucus biosynthesis in the abomasums" in Vet. Res., 42: 61,
- Singh, B.B., R. Sharma, H. Kumar, H.S. Banga, R.S. Aulakh, J.P. Gill and J.K. Sharma, (2006) "Prevalence of *Cryptosporidium parvum* infection in Punjab (India) and its association with diarrhea in neonatal dairy calves," in Vet. Parasitol. 140(1-2):162-5,
- Stromberg, B.E., Gasbarre, L.C., Waite, A., Bechtol, D.T., Brown, M.S., Robinson, N.A., Olson, E.J., Newcomb, H, (2012) "*Cooperia punctata*: effect on cattle productivity?" in Vet. Parasitol, 183: 284-291,
- Sunday Shende Kometa1 & Mathias Ashu Tambe Ebot (2012). Watershed Degradation in the Bamendjin Area of the North West Region of Cameroon and Its Implication for Development. *Journal of Sustainable Development*; Vol. 5, No. 9.
- Sunday Shende Kometa1 & Mathias Ashu Tambe Ebot, (2012) "Watershed Degradation in the Bamendjin Area of the North West Region of Cameroon and Its Implication for Development," in *Journal of Sustainable Development*; Vol. 5, No. 9,
- Pereira S.J., N.E. Ramirez, L. Xiao, L.A, (2002) "Ward, Pathogenesis of human and bovine *Cryptosporidium parvum* in gnotobiotic pigs," in *J. Infect. Dis.* 186: 715–718,
- M A. Tita1, D. Kuitcha, A. Magha1, J. Ndjama, Veronique KK. Beyala; (2012) Occurrence of macrophytes in the Mezam river system in Bamenda (Cameroon) and their role in nutrient retention :Syllabus Review, Sci. Ser. 3, 1 - 10
- R. A. B. Nichols, B. M. Campbell, and H. V. Smith (2003), *Applied and Environmental Microbiology*, Vol. 69 No. 7, pp 4183–4189

Acknowledgment

The Phyto-Biotechnology Research Foundation (PRF) for the Laboratory space, the laboratory staffs for their skilled in crossing check each sample analysed, Prof. Kenneth Yongabi A, who kept an eye everyday on the progress report as well as the Prof. Ukaga, Prof Nwoke and Prof. Tih for their proof reading with an eagles eyes. This is part of a PhD thesis non-funded.

Author Bibliography



Polycarp Ndikvu CHIA

I am a PhD student of Medical Parasitology and hold a first degree in Medical laboratory Technology and a Master in Medical Parasitology.

I lecture in the Catholic University of Cameroon (CATUC) Bamenda and the acting Head of Department of Biochemistry.

After working with the Cameroon Baptist Convention Health Board hospital laboratories for 5 years, I became interested in infectious diseases especially Zoonotics. Cryptosporidiosis is a challenging disease without a cure and the method use in disinfecting water does not destroy the oocyst. I am interested in investigating the dynamics of this pathogen thoroughly both in my PhD and after.

Email: polycarpc@gmail.com