

# SOME RESULTS ON JANOWSKI CLOSE TO CONVEX MAPPINGS

Melike Aydoğın

Department of Mathematics  
Isik University- Sile Kampusu  
Mesrutiyet Koyu Universite Sokak, D Kapi No:2  
34980, Istanbul, Turkey

## Abstract

Let  $J_C(A, B)$  denote the class of functions  $\phi(z) = z + \sum_{k=2}^{\infty} a_k z^k$  are analytic in the open unit disc  $D$  such that

$$\frac{z\phi'(z)}{s(z)} \prec \frac{1 + Az}{1 + Bz}, -1 \leq B < A \leq 1 \quad (1)$$

where  $s(z) = z + \sum_{k=2}^{\infty} b_k z^k$  is convex in  $D$ . In this paper we determine the coefficient estimates distortion theorems for this class.

**Mathematics Subject Classification:** 30C45, 30C55

**Keywords:** subordination, convex function, close to convex function, janowski function

## 1 Introduction

Let  $H(D)$  be the linear space of all analytic functions defined in the open unit disc  $D$ . Let  $w(z) = \sum_{k=1}^{\infty} c_k z^k$  be an analytic function in the open unit disc and satisfying the conditions  $w(0) = 0$  and  $|w(z)| < 1, z \in D$ .

Let  $C$  denote the class of functions such as

$$s(z) = z + \sum_{k=2}^{\infty} b_k z^k \quad (2)$$

analytic and convex in  $D$ .

Let  $J_C$  denote the class of functions such as

$$\psi(z) = \int_0^z \frac{s(z)}{z} \quad (3)$$

is Janowski and convex in  $D$ .

Let  $J_C(A, B)$  denote the class of functions such as

$$\phi(z) = z + \sum_{k=2}^{\infty} a_k z^k \quad (4)$$

analytic in  $D$  and satisfying the conditions;

$$\frac{z\phi'(z)}{s(z)} \prec \frac{1 + Az}{1 + Bz}, -1 \leq B < A \leq 1, z \in D, s(z) \in C \quad (5)$$

If we use the definition of subordination principle for  $\phi(z) \in J_C(A, B)$  if and only if  $\phi(z)$  can be represented in the form:

$$\frac{z\phi'(z)}{s(z)} = \frac{1 + Aw(z)}{1 + Bw(z)}, w(z) \in H(D), -1 \leq B < A \leq 1, z \in D \quad (6)$$

We study the  $J_C(A, B)$  and obtain coefficient estimates, distortion theorems.

## 2 Some Preliminary Lemmas

We need the following lemmas.

**Lemma 2.1** Let  $\frac{z\phi'(z)}{s(z)} = p(z) = 1 + \sum_{k=1}^{\infty} p_k z^k$  then  $|p_n| \leq (A - B)$ ,  $n \geq 1$ . The bounds are sharp, being attained for the functions

$$p_n(z) = \frac{1 + A\delta z^n}{1 + B\delta z^n}, |\delta| = 1 \quad (7)$$

[Goel and Mehrok].

**Lemma 2.2** If  $w(z) \in H(D)$  then for  $|z| = r < 1$

$$|zw'(z) - w(z)| \leq \frac{r^2 - |w(z)|^2}{1 - r^2} \quad (8)$$

[Singh and Goel].

**Lemma 2.3** Let  $p(z) = \frac{1+Bw(z)}{1+Aw(z)}$ ,  $w(z) \in H(D)$ , then for  $|z| = r < 1$

$$\operatorname{Re}\left[Ap(z) + \frac{B}{p(z)}\right] + \frac{r^2 |Ap(z) - B|^2 - |1 - p(z)|^2}{(1 - r^2) |p(z)|} \leq \frac{AB(A + B)r^2 - 4ABr + (A + B)}{(1 - Ar)(1 - Br)}; R_1 \leq R_0,$$

$$\frac{2}{(1-r^2)}(1-ABr^2) - [(1-A)(1-B)(1+Ar^2)(1+Br^2)]^{\frac{1}{2}}; R_1 \geq R_0.$$

where  $A \neq 1$ ,  $R_1 = \frac{1-Br}{1-Ar}$  and

$$R_0^2 = \frac{(1-B)(1+Br^2)}{(1-A)(1+Ar^2)}$$

The bounds are sharp.[Goel and Mehrotra].

### 3 Coefficient Inequalities

**Theorem 3.1** If  $\phi(z) \in J_C(A, B)$  then,

$$|a_n| \leq \frac{1}{n} + \frac{(n-1)(A-B)}{n}, n \geq 2 \quad (9)$$

The bounds are sharp.

Proof: Using (1.2) and (1.4) in (2.1) we get;

$$z(1 + \sum_{k=2}^{\infty} ka_k z^{k-1}) = (z + \sum_{k=2}^{\infty} b_k z^k)(1 + \sum_{k=1}^{\infty} p_k z^k) \quad (10)$$

Equating the coefficient of  $z^n$  in (3.2) we have;

$$na_n = b_n + p_1 b_{n-1} + p_2 b_{n-2} + \dots + p_{n-1} \quad (11)$$

Therefore using (3.2),

$$n|a_n| \leq |b_n| + (A-B)[|b_{n-1}| + |b_{n-2}| + \dots + |b_2| + 1] \quad (12)$$

Also it is well known that  $|b_n| \leq 1$ ,  $n \geq 2$ . Hence  $|a_n| \leq \frac{1}{n} + \frac{(n-1)(A-B)}{n}$ . For  $n = 2$  equality signs in (3.1) holds for the function  $\phi(z)$

$$\phi'_n(z) = \frac{1}{(1-\delta_1 z)} \frac{1+A\delta_2 z^{n-1}}{1+B\delta_2 z^{n-1}}, |\delta_1| = 1, |\delta_2| = 1 \quad (13)$$

On putting  $A = 1$ ,  $B = -1$  in above theorem, we get the following result due to Grawod and Thomas.

**Corollary 3.2** Let  $\phi(z) \in J$ , then  $|a_n| \leq 2 - \frac{1}{n}$

## 4 Main Results

**Theorem 4.1** *If  $\phi(z) \in J_C(A, B)$  then for  $|z| = r$ ,  $0 < r < 1$*

$$\frac{1 - Ar}{(1 - Br)^{\frac{2B-A}{B}}} \leq |\phi'(z)| \leq \frac{1 + Ar}{(1 - Br)^{\frac{2B-A}{B}}}; B \neq 0, \quad (14)$$

$$\frac{e^{-Ar}(1 - Ar)}{r(1 - Br)} \leq |\phi'(z)| \leq \frac{e^{Ar}(1 + Ar)}{r(1 - Br)}; B = 0. \quad (15)$$

$$\int_0^r \frac{1 - At}{(1 - Bt)^{\frac{2B-A}{B}}} dt \leq |\phi(z)| \leq \int_0^r \frac{1 + At}{(1 - Bt)^{\frac{2B-A}{B}}} dt; B \neq 0, \quad (16)$$

$$\int_0^r \frac{e^{-At}(1 - At)}{t(1 - Bt)} dt \leq |\phi(z)| \leq \int_0^r \frac{e^{At}(1 + At)}{t(1 - Bt)} dt; B = 0. \quad (17)$$

*Estimates are sharp.*

Proof: Let  $s(z) = z + b_2z^2 + \dots$  analytic and convex;

$$\operatorname{Re}\left(1 + z \frac{s''(z)}{s'(z)}\right) > 0$$

If  $\frac{\phi'(z)}{s'(z)} > 0$

$$\operatorname{Re} \frac{\phi'(z)}{s'(z)} = \frac{1 + A\phi(z)}{1 + B\phi(z)}$$

$$z \frac{s'(z)}{s(z)} = \frac{1 + A\phi(z)}{1 + B\phi(z)}$$

Let  $\psi(z)$  Janowski and convex  $\in J_C$ ;  $\psi'(z) = \frac{s(z)}{z}$ ,

$$\log \psi'(z) = \log s(z) - \log z$$

$$\frac{\psi''(z)}{\psi'(z)} = \frac{1}{z} + \frac{s'(z)}{s(z)} \Rightarrow z \frac{\psi''(z)}{\psi'(z)} = -1 + z \frac{s'(z)}{s(z)} \Rightarrow$$

$$1 + z \frac{\psi''(z)}{\psi'(z)} = z \frac{s'(z)}{s(z)} \Rightarrow \psi(z) \text{ is convex}$$

$$1 + z \frac{\psi''(z)}{\psi'(z)} = z \frac{s'(z)}{s(z)} = \frac{1 + A\phi(z)}{1 + B\phi(z)}$$

Therefore we can write the following inequalities;

$$r(1 - Br)^{\frac{A-B}{B}} \leq |s(z)| \leq r(1 + Br)^{\frac{A-B}{B}}; B \neq 0, \quad (18)$$

$$e^{-Ar} \leq |s(z)| \leq e^{Ar}; B = 0. \quad (19)$$



