

27th European

Ophthalmology Congress

November 26-28, 2018 | Dublin, Ireland

Effectiveness of the modified method of surgical treatment of terminal glaucoma

Bakhritdinova Fazilat Arifovna and **B E Bilalov**
Tashkent Medical Academy, Uzbekistan

Introduction: Terminal glaucoma is one of the most severe forms of glaucomatous process and is usually accompanied by pain syndrome, which practically is not amenable to drug therapy.

Purpose: The purpose is to improve the results of surgical treatment of terminal glaucoma by preventing excessive scarring by auto drainage of the filtration zone.

Material & Methods: Depending on the conducted surgical treatment the patients were divided into 2 groups. The first (control) group consisted of patients who underwent deep sclerectomy with preliminary posterior trepanation of the sclera - 30 patients (30 eyes). The second group consisted of 30 (30 eyes) patients who underwent an antiglaucomatous operation with auto drainage of the filtration zone.

Results: A week after the operation, the tonometric intraocular pressure (IOP) in the patients of the main group decreased 2.3 times relative to the preoperative indices and averaged 20.3 ± 1.8 mmHg. In the patients of the control group, it decreased 2.1 times relative to the operative indices and averaged 22.5 ± 1.7 mmHg. According to UBM data, in the early postoperative period (1 month) in the patients of the main group the filtration pillow (FP) was significantly expressed in the patients of the main group, and in the patients of the control group it was less expressed and more flat. The thickness of scleral flap (SF) in the patients of the main group decreased somewhat in comparison with the previous periods, averaging 0.28 ± 0.002 mm, and the acoustic density of the structure remained in the range of $65 \pm 5\%$. In patients of the control group, the thickness and acoustic density of the SF continued to increase and averaged 0.38 ± 0.01 mm and $90 \pm 5\%$, respectively. In this phenomenon of developing subatrophy of the eyeball for up to one year, no patient was registered.

Conclusion: The efficiency of using the method of antiglaucomatous operation with auto drainage of the filtration zone in terminal glaucoma was proved in 75% of cases.

Biography

She is the head of the department of Ophthalmology at the Tashkent Medical Academy.

dr.ben58@mail.ru

Notes: