

# 13<sup>th</sup> Euro Obesity and Endocrinology Congress

and

# 16<sup>th</sup> World Congress on Nutrition and Food Chemistry

September 18-20, 2017 | Zurich, Switzerland



## Shougang Wei

*Capital Medical University, China*

### Obesity leads to iron retention in the duodenum of mice likely due to increased production of adipose-derived hepcidin

Obese people and animals have higher rates of iron deficiency (ID) than their normal weight peers. It was still uncertain whether obesity-related ID is a true or functional deficiency of iron. This study was to determine the effects and the possible underlying mechanisms of obesity on duodenal iron absorption and liver iron accumulation. C57BL/6J mice were randomly divided into high-fat diet-induced obese (DIO) group and normal control (NC) group to be fed respectively for 16 weeks. Oral iron absorption was tested by measuring serum iron, liver iron and the retained duodenal iron 90 min after intra gastric administration of  $^{57}\text{FeSO}_4$  solution. The protein expression levels of iron transporters in duodenum and liver were evaluated by western blotting. Hepcidin mRNA levels in the liver and adipose tissues were quantified by real-time RT-PCR. The results showed that DIO mice had significantly higher iron retention in the duodenum, lower iron concentration in plasma and liver than NC mice. The protein expression levels of ferroportin-1 (Fpn1) in duodenum and transferrin receptor-2 (TfR2) in the liver were markedly decreased in DIO mice. Hepcidin mRNA levels in visceral adipose tissue but not in the liver were higher in DIO mice than NC mice. In conclusion, obesity-related ID may be attributed to impaired intestinal iron absorption of which iron being retained in the duodenal enterocytes, not to that iron being accumulated in the liver. Increased expression of visceral adipose hepcidin probably is the immediate cause for the malabsorption of iron in obesity by inducing reduction of the duodenal Fpn1.

#### Biography

Shougang Wei serves as Professor, PhD supervisor and Deputy Director in Department of Children's and Women's Health, School of Public Health, Capital Medical University, Peking, China. He has been engaged in the study of child and adolescent health, mainly focused on the field of childhood obesity about its health risks, pathogenic factors, and preventive and treatment measures.

[shangwei@ccmu.edu.cn](mailto:shangwei@ccmu.edu.cn)

#### Notes: