

20th World Congress on

NUTRITION & FOOD SCIENCES

May 14-16, 2018 Tokyo, Japan

Alcohol exposure suppresses neural crest cells generation and differentiation during early chick embryo**Ping Zhang**

Jinan University, China

It is now known that excess alcohol consumption during pregnancy can cause fetal alcohol syndrome (FAS) in which several characteristic craniofacial abnormalities are often visible. However, the molecular mechanisms of how excess ethanol exposure affecting cranial neural crest cells (CNCCs), the progenitor cells of the cranial skeleton, is still not clear. In the study, we investigated the effects of ethanol exposure on CNCCs migration both in early chick embryo and *in vitro* explant culture. First of all, we demonstrated that ethanol treatment caused alizarin red-stained craniofacial developmental defects including parietal defect. Second, immuno-fluorescent staining with neural crest special markers indicated that CNCCs generation was inhibited by ethanol exposure. And, double immuno-fluorescent staining (Ap-2 α /PHIS3, HNK1/BrdU and AP-2 α /c-caspase3) revealed that ethanol exposure inhibited CNCCs proliferation and increased apoptosis. In addition, it inhibited NCCs production by repressing the expression level of key transcription factors which regulate neural crest development by altering expression of epithelial-mesenchymal transition (EMT)-related adhesion molecules in the developing neural crests. In sum, we have provided experimental evidence that excess ethanol exposure during embryogenesis disrupts CNCCs survival, EMT and migration, which in turn causes defective cranial bone development.

1696937253@qq.com