

17th International Conference and Exhibition on
NANOMEDICINE AND NANOTECHNOLOGY IN HEALTHCARE

November 23-24, 2017 Melbourne, Australia

Cytotoxic effects of bioactive compounds isolated of *Iris persica* L. on human cancer cell linesHawraz Ibrahim M Amin¹, Faiq H S Hussain¹ and Giovanni Vidari²¹Salahaddin University-Erbil, Iraq²University of Pavia, Italy

Genus *Iris* (Iridaceae) comprises over 300 species; 12 of them are present in Iraq. *Iris persica* has been used in Kurdish traditional medicine for the treatment of wound inflammation and tumor. However, Chemical and biological aspects of *I. persica* have not yet been investigated. The present study reports the first investigation on the isolation and characterization of bioactive compounds from flowers, bulbs and rhizomes of *I. persica* that has been collected from Kurdistan Region-Iraq and cytotoxicity effect of the isolated compounds against six human cancer cell lines were evaluated. Dry flowers, bulbs and rhizomes of *I. persica* were exhaustively extracted by maceration at room temperature, solvents of increasing polarity: hexane, methanol, methanol/water 70:30. Chlorophylls were removed from the methanolic extracts of flowers by filtration on a C-18 reversed phase column. Subsequently, the methanolic extracts of the flowers, bulbs and rhizomes were separately fractionated by repetitive preparative MPLC, on C-18 reversed phase, affording four compounds as the major products: tectorigenin (1), embinin (2), isovitexin (3) and trans-resveratrol-3-O- β -D-glucopyronoside (4). The structures of the compounds were identified on the basis of spectroscopic analyses and comparison with literature data. The cytotoxic activity was measured against six human cancer cell lines, the effects of two isolated compounds, Tectorigenin (1) and embinin (2), on the proliferation of tumor cells were evaluated in comparison with the well-known antitumor drug cis-diamminedichloroplatinum (II) (cisplatin) by MTT assays. In particular, MCF7 and SkBr3 breast, endometrial Ishikawa, ovarian BG-1, mesothelioma IST-MES1 and lung A549 cancer cells were treated for 48 hours with increasing concentrations of tested compounds. Compound P2 showed a stronger inhibitory activity than cisplatin in five cell lines from total of the six cell lines; MCF7, SkBr3, Ishikawa, BG-1, IST-MES1 and A549, embinin IC₅₀±SD of 6 (±3), 4(±1), 10(±3), 8(±2), 7(±2) and 9(±1) μ M, while cisplatin (standard) IC₅₀±SD of 17(±4), 10(±2), 10(±3), 12(±3), 12(±2) and 13(±2) μ M, respectively. In conclusion, this is first study on phytochemical study of *I. persica* as well as cytotoxic activity of embinin isolated flowers of *I. persica*, this study confirms that Embinin could be considered as a natural anticancer. At the same time, the present study confirms the traditional use of *Iris persica* L. in the treatment of tumor.

hawraz_px@yahoo.com