

# World Congress on Clinical Pediatrics and Pediatric Oncology & Care

November 12-13, 2018 | Lisbon, Portugal

Charisse B Pulmano<sup>1, 2</sup>

<sup>1</sup>University of the City of Manila,  
Philippines

<sup>2</sup>Hospital ng Maynila Medical Center,  
Philippines

## Association of breastfeeding practices and selected clinical outcomes of patients aged three months to two years old diagnosed with PCAP C in a tertiary government hospital in Manila from January 2013-January 2016

Pneumonia continues to be a significant cause of infant and childhood morbidity and mortality in the Philippines. In the Philippines, pneumonia still continues to be one of the leading causes of mortality and morbidity among infants. With this as a social burden, various strategies have been implemented to decrease childhood morbidity and mortality which is congruent to Millennium Development Goals. In line with this, the World Health Organization strongly advocates the importance of exclusive breastfeeding for the first six months of life. Supplemental breast feeding is not recommended until at least age two and exclusive breastfeeding may be continued thereafter. In local setting, the Department of Health has launched a campaign entitled “Breast feeding TSEK: Tama, Sapat, at Ekslusibo”. This program has been implemented and warrants strict compliance to reduce childhood mortality from inevitable diseases. In line with this, the author wants to determine the association of breastfeeding practices and selected clinical outcomes of the PCAP C patients aged three months to two years in a tertiary government hospital from January 2013-January 2016. Main outcomes were identified as to length of hospital stay (three-five days, six-10 days, >10 days), disposition (Discharged improved, Mortality), presence of complications such as sepsis, pneumothorax, pleural effusion, empyema, or warranting intubation (with/without) among others. This study used a retrospective cohort design gathering data from records of patients admitted in a tertiary hospital in a given set period. Plan of analysis used percentage, risk ratio/chi square test of association. From this study, it was concluded that exclusive breastfeeding can significantly lower the risk of complications of pneumonia and mostly patient dispositions are markedly improved. In fact, it was found that the risk of moderate to long hospitalization, risk of having complications, and risk of dying are 2.91 (95% CI: 1.91, 4.43) and 3.05 (95% CI: 1.86, 5.04), 3.33 (95% CI: 2.36, 4.70), and 2.23 (1.19, 4.20) times higher, respectively, among mixed fed patients compared to breastfed patients.

### Biography

Charisse B Pulmano, he is affiliated to The Hospital Manila Medical Centre in Philippines. Currently he is practising at Mansila-Ospital Longxodon where he belongs to the department of Pediatric Nutrition and he has published several articles in many reputed journals in his research towards pediatrics nutrition.

charisse.pulmano@yahoo.com

### Notes: