26th European Pediatrics Congress

October 22-23, 2018 | Amsterdam, Netherlands

Frequency and antimicrobial susceptibility pattern of microorganisms isolated from hospitalized infantile burn cases in a tertiary care hospital



Jalal Uddin Akber Baqai Medical University, Pakistan **Objective:** The aim of this study was to determine the frequency of organisms and their antimicrobial sensitivity pattern in infantile burn.

Methods: This retrospective study was conducted at the plastic surgery, burn unit of Patel Hospital during period of 7 years from January 2007 to December 2013. Children aged one year or less at the time of admission were included in the study. SPSS 21 version was used for statistical analysis.

Results: Total 789 pediatric burn patients were admitted in Patel Hospital during January 2007 to December 2013, in which 106 were infants. Eighty three (78.3%) infants had scald burn, 21 (19.8%) had fire burn and 2 (1.9%) had chemical burn. Out of 106 infantile burn cases, 28 (26.4%) had growth of organisms in wound cultures. Initially at the time of admission only 2 (7%) of infants had growth of organisms in wound cultures but on subsequent cultures the growth of organisms increased. Single organism was isolated in wound cultures of 9 (32.1%) patients, while two organisms were found in 8 (28.6%) and three organisms were found in 11 (39.3%) infants. The commonest organisms present both in scald and fire burn were Staphylococcus aureus 17 (60.7%) followed by Acinetobacter species 14 (50%) and Pseudomonas aeruginosa 13 (46.4%) respectively. If we see the sensitivity pattern, Staphylococcus aureus was 100% sensitive to vancomycin and linezolid followed by fusidic acid 47%. If we see the culture and sensitivity pattern of Pseudomonas aeruginosa, Acinetobacter species and Klebsiella species all were 100% sensitive to polymyxin B. While Providencia species and Proteus species were 100% sensitive to cefoperazone + sulbactam and meropenem.

Conclusion: This study high lights that *Staphylococcus aureus*, *Acinetobacter* species and *Pseudomonas aeruginosa* are the common organisms in infantile burn. While vancomycin and polymyxin B are the effective empirical therapy in our setup. Antibiotic resistance due to inappropriate use of drugs is a common finding in our environment and medical staff must be educated regarding the rational use of antibiotics. Wound swabs should be performed in all cases

Biography

Jalal uddin Akber graduated from Liaquat Medical College in 1988 and Presently working as Chairman and Head of Pediatrics at Children's Hospital Baqai Medical University, Karachi, since last 7 years. He is also working as consultant pediatrician at Patel hospital Karachi for last 18 years and a member of plastic surgery and burn unit. Dr. Jalal uddin Akber is a master trainer for different project of UNICEF, WHO and Pakistan Pediatric Association. He is a member of Pakistan Pediatrics Association, American Academic of Pediatrics as well as Polio Eradication Certification Committee, Government of Pakistan. He has been a guest speaker in more than 100 national and international conferences, seminars and workshops.

akber.jalal@yahoo.com

Notes: