

# 26<sup>th</sup> European Pediatrics Congress

October 22-23, 2018 | Amsterdam, Netherlands

## Cause specific management of shock in neonate

**Mohammad Monir Hossain**  
Dhaka Shishu Hospital, India

Shock is characterized by inadequate oxygen delivery to tissues to meet demand because of circulatory failure. The immediate aim of management of neonatal shock is to optimize perfusion and delivery of oxygen and nutrients to the tissues. Understanding the pathophysiology of neonatal shock helps to recognize and classify shock in the early compensated phase and initiate appropriate treatment. Hypovolemic shock in neonate is usually due to antepartum hemorrhage, post-natal blood loss iatrogenic, or secondary to disseminated intravascular coagulation or vitamin K deficiency, or excessive insensible water loss in extreme pre-terms. Cardiogenic shock in the neonate may be caused by myocardial ischemia due to severe intra-partum asphyxia, arrhythmias, primary structural heart disease and mechanical reduction of cardiac function or venous return secondary to tension pneumothorax or diaphragmatic hernia and disturbance of transitional circulation, due to persistent pulmonary hypertension in newborn, or patent ductus arteriosus in premature infants. Distributive shock caused by neonatal sepsis, vasodilation, myocardial depression, or endothelial injury and obstructive shock is caused from tension pneumothorax or cardiac tamponade. The immediate aim of management of neonatal shock is to optimize perfusion and delivery of oxygen and nutrients to the tissues. The American College of Critical Care Medicine estimates that 60 minutes is the average time needed to provide adequate circulatory support and block the development of shock. The first step in managing shock in the newborn during the first 5 minutes is to recognize cyanosis, respiratory distress and decreased perfusion. This should be followed immediately by airway access and ventilation to optimize oxygenation. Rapid peripheral, central venous or intraosseous access is of primary importance in the initial management of the newborn in shock. Any baby with shock and hepatomegaly, cyanosis or a pressure gap between upper and lower limbs should be treated with prostaglandin within 10 minutes of birth, until congenital heart disease is excluded. Inotropes like dopamine, dobutamine, epinephrine and norepinephrine are indicated via IV or IO route before central access is achieved when myocardial contractility remains poor despite adequate volume replacement. Delay increases mortality 20-fold.

## Notes: