

International Conference on

GYNECOLOGY & OBSTETRICS PATHOLOGY &

2nd World Congress on

EMBRYOLOGY & *IN VITRO* FERTILIZATION

March 30-31, 2018 | Orlando, USA

Special case of extra-uterine pregnancy

Achour Radhouane and Khaled Neji
Tunis El Manar University, Tunisia

Spontaneous bilateral ectopic pregnancy is rare; therefore preoperative diagnosis is uncommon indicating limitations of Ultrasonography. We report an unusual case of ectopic pregnancy in which patient had spontaneous bilateral tubal ectopic pregnancy. A 33-year-old gravida 8, para 3 woman was referred to the gynecology service with an approximate gestational age of 10 weeks. Emergency exploratory laparotomy revealed haemoperitoneum of approximately 2000 mL there was a ruptured fimbrial ectopic pregnancy with active bleeding on the right side. The left tube showed an intact ectopic pregnancy 10 cm in the ampullary region that was bleeding and forming an organized haematoma at the fimbrial end. This case underscores the importance of identifying and closely examining both tubes at the time of surgery, even in the presence of significant adhesive disease.

Biography

Achour Radhouane is Associate Professor at Faculty of Medicine, El Manar University of Tunis, Tunisia. He has published many basic and clinical articles in relation to gynecology and obstetrics. He serves as Associate Professor at Emergency Department of Gynecology and Obstetrics in Maternity and Neonatology Center Tunis, Tunisia. He also serves as Member of the Editorial Team of *Asian Pacific Journal of Reproduction*, *The Global Journal of Rare Diseases* and *Journal of Neonatal Biology*.

radhouane.a@live.com

Notes: