5th Asia Pacific

GYNECOLOGY AND OBSTETRICS CONGRESS

August 22-23, 2018 Tokyo, Japan

Intimate area reshaping with a new specific HA filler

Dawid Serafin Serafin Clinic, Poland

The intervention with dermal filler is indicated in the reshaping of congenital or after weight loss thin labia majora and it is also indicated to improve vaginal defects or complications after episiotomy or spontaneous perineal damage. The growing interest to improve the aesthetic and functional intimate area of women, has lead the research to develop specific resorbable injectables to correct and reshape this specific area. Considering that the hypotrophy or atrophy of labia majora need a volumizing gel (i.e. elasto-plastic rheological features), but also a gel able to preserve a very natural sensation and projection of the skin, it is of interest a new specific formulation of HA filler. The formulation (intense ROSE) presents the advantage to use an ultra-pure HA as row material (derived from a probiotic bacterium: *Bacillus subtilis*) and especially to use as stabilizer a polymeric cross-linker (PEG). This polymer allows to create gels with important filling effect (i.e. same volume with less amount of gel compared to other HA filler) and to obtain both very natural results and sensation thanks to the thermo-dynamic behavior of the gel and physiological distribution into the skin layers. The author will illustrate his personal approach and results in reshaping this area with this specific new formulation.

Biography

Dawid Serafin is a Specialist of Gynecology and Obstetrics. He is specialized in the treatment of aesthetic gynecology and the treatment of aesthetic problems with hyaluronic acid fillers. He is a Trainer in the field of Aesthetic Gynecology, as well as being a participant, lecturer and author of articles on aesthetic gynecology.

dawid@serafinclinic.pl

Notes: