

September 04, 2013 Holiday Inn Orlando International Airport, Orlando, FL, USA

Influence of tillage and organic manure sources on termite infestation, sprouting of cassava cuttings and plant growth in a termite infested soil in Owerri, south-eastern Nigeria

Timothy T. Epidi² and Kennedy.O Ogbedeh¹

Department of Crop Science and Technology, Federal University of Technology Owerri, Nigeria Department of Crop Production Technology, Niger Delta University, Nigeria

Termites have been identified as one of the major pests of field-grown cassava in Nigeria. The study investigated the influence of tillage and organic manure sources on termite infestation, sprouting of cassava cuttings and plant development in a termite infested soil in Owerri, South-Eastern Nigeria. Field trials were conducted in 2007, 2008 and 2009 at the Teaching and Research Farm of the Federal University of Technology Owerri. The experiments were laid out in a 3x6 split-plot factorial arrangement fitted into a randomized complete block design (RCBD) with three replications. The cassava cultivar used was TMS 4 (2) 1425. Treatments consisted of three tillage methods (Zero, flat and mound), two rates of municipal waste (1.5 and 3.0 tonnes/ha), two rates of *Azadirachta indica* (neem) leaves (20.0 and 30.0 tonnes/ha), control (0.0 tonnes/ha) and carbofuran (chemical check). Data were collected at planting (2WAP), growth and maturing phases of the plant. These were analyzed using a Mixed-model procedure of Statistic Analysis System (SAS). Means were separated using least significance difference (LSD) at 5% level of probability. Results showed *A. indica* leaves at 30 t/ha produced highest sprouting percentage (43.5-80.6%) of planted cassava cuttings, followed by zero tillage (43.46-75.93%). Also termite incidence and severity (infestation) were significantly depressed with the application of carbofuran; *A. indica* leaves at 20 and 30 t/ha, and zero tillage especially at 16 to 22 WAP. However, termite infestation remarkably increased at 2 to 8 WAP, but decreased at 10-16 WAP and finally increased again at 18-30 WAP.

tepidi@yahoo.com