

## International Conference on **Thyroid Disorders and Treatment** February 29-March 01, 2016 Philadelphia, Pennsylvania, USA

Chosen immunological aspects in autoimmune Thyroid diseases in developmental age

Artur Bossowski Medical University in Bialystok, Poland

**Introduction:** Up till now, altered balance of Th1 and Th2 immune cells has been postulated to play an important role in the pathogenesis of Autoimmune Thyroid Tiseases (AITD). However, recent studies on Thyroid diseases suggest a new role for Th17 (T helper 17) cells that have been classified as a new lineage, distinct from Th1, Th2 and T-reg cells. Despite wide interest, the role of Th17 cells in the pathogenesis of inflammatory and autoimmune diseases is still being debated. Th17 cells are involved in immune responses against extracellular pathogens and have the ability to secrete cytokines: IL-17, IL-17F, IL-21 and IL-23. Th17 cells can be characterized by several surface markers, i.e. CCR6 (CD196), IL-23R, IL-12Rbeta2 and CD161.

Aim: To estimate the proportions of circulating CD4+CD161+CD196+ and CD4+IL-17+ Th17 cells and serum concentrations of IL-17 & IL-23 in patients with Graves' disease (GD, n=42, mean age  $\pm$  SEM 14.2  $\pm$  4 years), Hashimoto's Thyroiditis (HT, n=67, mean age  $\pm$  SEM 14.8 $\pm$ 2 yrs) and in healthy controls (C, n=45, mean age  $\pm$  SEM 15.1  $\pm$  3 yrs).

**Material & Methods:** Polychromatic flow cytometry and several fluorochrome-conjugated monoclonal antibodies were applied to delineate Th17 cells with CD4+CD161+CD196+ or CD4+IL-17+ phenotype using apparatus FACS Calibur (BD Biosciences). The expression of IL-17 and IL-23 were analyzed by Bio-Tek ELx800 ELISA reader.

**Results:** In untreated HT children, we observed an increased percentage of CD4+CD161+CD196+ (p<0.04) and CD4+IL-17+ (p<0.01) Th17 lymphocytes in comparison to the healthy controls. In GD children we did not reveal such abnormalities in the population of these cells. In cases with HT, a positive correlation between the percentage of CD4+IL-17+ and CD4+CD161+CD196+ T cells and serum level of anti-TPO antibodies (r=0.48; p<0.025; r=0.65; p<0.01; respectively) was detected. In untreated patients with AITD we observed an increased levels of IL-23 in comparison to control group (GD: p=0.004, HT: p=0.046). Methimazole treatment in GD led to decrease in these cytokine levels in a period of 6-12 months. However, during 6-24 months of L-thyroxine therapy in HT there wasn't any reduction of IL-23 concentration compared with HC. IL-17 was elevated only in HT patients in comparison to the controls (p=0.021), which normalized during therapy.

**Conclusions:** We conclude that the increased percentage of Th17 cells and elevated level of IL-17 and IL-23 cytokines in children with HT can suggest their role in initiation and development of immune and inflammatory processes in this endocrinopathy.

## Biography

Artur Bossowski completed Doctorate in June 2001 and have already presented preliminary findings at various local and international meetings. He also started to performed investigation of lymphocyte subpopulations and co-stimulatory molecules and integrins in peripheral blood and in Thyroid tissues of patient with Graves' disease and non-toxic nodular goiter.

abossowski@hotmail.com

Notes: