

JOINT EVENT

World Congress on **Psychiatry & Psychological Syndromes**29<sup>th</sup> International Conference on &**Adolescent Medicine and Child Psychology**

December 06-07, 2018 | Rome, Italy

**Undiagnosed diabetes mellitus and associated factors among psychiatric patients receiving antipsychotic drugs at the University of Gondar Hospital, Northwest Ethiopia****Daniel Asmelash, Wondale Getnet, Belete Biadgo, Sintayehu Ambachew, Tadele Melak, Lemmesa Melese, Shiwaneh Damite, Habtamu Wondifraw Baynes and Molla Abebe**  
University of Gondar, Ethiopia

**Background:** Undiagnosed diabetes mellitus cases are at higher risk for diabetic related complications. In low income African countries, patients with undiagnosed diabetes mellitus account for 75% of diabetes cases. Psychiatric disorders have a greater impact on the global burden of diseases and disability associated with chronic diseases like diabetes mellitus and cardiovascular diseases.

**Objective:** To assess the prevalence of undiagnosed diabetes mellitus and its associated factors among psychiatric patients receiving antipsychotic drugs at the University of Gondar Hospital, Northwest Ethiopia.

**Methods:** Institution based cross-sectional study was conducted at the University of Gondar Hospital from February to April 2016. A total of 205 psychiatric patients aged above 15 years who were taking antipsychotics were included by the simple random sampling method. Fasting blood glucose, triglycerides and cholesterol level were determined from venous blood samples to evaluate diabetes mellitus based on WHO criteria.

**Results:** Among 205 psychiatric patients taking antipsychotics, 15(7.3%) had undiagnosed diabetes mellitus. Duration of antipsychotic treatment and sex had a statistically significant association with the prevalence of undiagnosed diabetes mellitus. As the duration of antipsychotic drug treatment increased by one year, the risk of having diabetes mellitus increase by 1.47 times (AOR:1.47 CI: 1.021-2.125).

**Conclusion:** The prevalence of undiagnosed diabetes mellitus among psychiatry patients taking antipsychotics was higher than the estimated diabetes national prevalence of Ethiopia. Screening of diabetes mellitus in particular patients having a longer duration of antipsychotic treatment is mandatory to bring more undiagnosed cases for medical attention.

roberto.cancio@lmu.edu