


5<sup>th</sup> World Congress and Expo on

# Applied Microbiology

November 12-13, 2018 | Edinburgh, Scotland



## Charambira K

*International Union against Tuberculosis and Lung Disease, Zimbabwe*

### Increased rifampicin mono-resistance prevalence in Zimbabwe – Is the higher prevalence of codon 523 to 529 mutation in the *rpoB* gene an attributing factor?

**Background:** Zimbabwe conducted a second anti-tuberculosis drug resistance survey (TB-DRS) in 2015/16 using the Xpert MTB/RIF assay. This assay uses molecular beacons in five overlapping regions of the *rpoB* DNA region. The probes detect mutations in the codons 507 to 511 (Probe A), 511 to 518 (Probe B), 518 to 523 (Probe C), 523 to 529 (Probe D), and 529 to 533 (Probe E). We report the frequencies of mutations in *rpoB* gene of the *Mycobacterium tuberculosis* (MTB) among the TB-DRS samples with rifampicin resistance tuberculosis (RR-TB) detected.

**Method:** A retrospective review of data collected through the TB-DRS and using the GxAlert platform to check the actual probe details for those tests from samples that had RR-TB strains. Sputum smear positive samples had an Xpert MTB/RIF assay done followed by phenotypic culture and drug susceptibility testing (DST) in those that had RR-TB detected.

**Results:** A total of 60 specimens had RR-TB detected on Xpert MTB/RIF assay. Of these, 50 (83.3%) had *Mycobacterium tuberculosis* growth on culture with 48 (96.0%) confirmed RR-TB on phenotypic DST. Among those confirmed RR-TB on phenotypic DST, 23 (47.9%) had rifampicin mono-resistance (RMR) detected and 25 had additional isoniazid resistance (MDR-TB). Probe E mutations occurred in 46% (23/50), probe D 34% (17/50), probe B 10% (5/50), probe A 2% (1/50) and probe C 2% (1/50) of the specimens. Among the RMR, probe D mutation occurred in 54.5% (12/22), probe E 36.4% (8/22), probe A 4.5% (1/22) and probe B 4.5% (1/22).

**Conclusion:** There is increase in the RMR prevalence from zero percent to 48% between the 1994/5 and 2015/6 TB-DRS. Rifampicin (RMP) seems to be associated with mutations in codons 523 to 529 of the *rpoB* gene of MTB DNA. GxAlert makes it possible to conduct such surveillance remotely and there is need for further studies to cement this.

### Biography

I am a medical doctor. I am the Programmatic Management of Drug Resistant TB Technical officer with The Union. I'm currently enrolled with the London School of Tropical Hygiene and Medicine.

[kcharambira@theunion.org](mailto:kcharambira@theunion.org)

### Notes: