Chromatography and Analytical Techniques

October 25, 2022 | Webinar

Valentin A. Krylov et al., J Chromatogr Sep Tech 2022, Volume 13

Features of gas chromatographic-mass spectrometric determination of o-phthalic acid esters in low alcohol wines coupled with emulsion liquidphase microextraction preconcentration

Valentin A. Krylov* and Pavel V. Mosyagin

N. I. Lobachevskii Nizhny Novgorod State University, Russia

Esters of o-phthalic acid are very dangerous for human health. Their occurrence in wines is connected with the Einflow from the plasticized polymer seals, plastic piping, tanks and stoppers. In this study the high sensitive gas chromatographic-mass spectrometric determination of phthalates in low alcoholic beverages (champagne, red and white wine) coupled ultrasound-assisted emulsification-microextraction was developed. As extractants environmentally friendly hydrocarbons n-heptane and n-hexane are proposed. The sources of possible systematic errors were investigated: leaking of o-phthalates from chromatographic septum; contamination of phthalates in solvents; influence of macro components of wines (sugar, alcohol, anthocyanins); the hydrolysis of o-phthalates and others. For the first time it is shown that the impact of these factors can lead to an overestimation or underestimation of the actual concentration of impurities by 1-2 orders of magnitude. The methods of accounting or elimination of systematic errors are proposed. Purification of solvents by <u>Rayleigh distillation method</u> allows to obtain samples with impurity content lower than (1-4).10-3 mgL-1. Containers for sampling and storage of samples to be analyzed should be made of borosilicate glass or quartz. The content of o-phthalates in wines was 0.03-1 mgL-1. The largest concentrations are characteristic for diethyl-, di-n-butyl- and di(2-ethylhexyl) o-phthalates.

The limits of detection of esters of o-phthalic acid in low alcohol beverages achieved are at the level of 10-6–10-5 mgL-1 and are highly competitive with the best world results. The relative expanded uncertainty of the determination of toxicants is at the level of 13- 30%.

Biography

Krylov V. A. has completed his PhD from Gorkiy State University, Russia. Currently, he is a professor at the Department of Analytical Chemistry of Nizhny Novgorod State University. The main direction of scientific research of professor Krylov is the development of the theory and applications of chromatography for the analysis of high purity substances, environmental objects and development of methods of the micro extraction. He is the author of two monographs and over 200 scientific papers, including reviews on the analytical chemistry of air, high purity volatile substances and liquid-liquid micro extraction. He presented papers in more than 75 Russian and international conferences.

Received: May 12, 2022; Accepted: May 13, 2022; Published: October 14, 2022