



Sleep, Anesthesiology, and the Neurobiology of Arousal State Control

Fethiye Aylin Sungur

Vita-Salute San Raffaele University, Milan, Italy

Abstract: (600 words)

Sleep, like breathing, is a biological cycle that the brain creates on its own. Neuronal networks that developed to govern normal sleep show a preference for modulating features that identify sedation and anaesthesia states. Sleep is divided into various stages, each of which has its own set of physiological and behavioural characteristics. Sleep and anaesthetic susceptibility are heritable phenotypes that are genetically modulated. This overview examines 40 years of research into the cellular and molecular systems that influence arousal state. The basic belief that sleep deprivation is a weakness has been refuted and replaced by clinical and preclinical research. Sleep restriction and deprivation reduce attentiveness, affect neuroendocrine control, and impair immune function. There is substantial evidence that lower levels of attentiveness have a negative influence on performance. The tidal change in public and legal viewpoints that today regard a sleep-deprived individual as handicapped is based on advances in neuroscience. ANESTHESIOLOGY has a big stake in figuring out how sleep affects clinical performance and career longevity. 1,2,3,4,5,6,7,8,9 The American Society of Anesthesiologists' emblem features attentiveness, because sleep deprivation decreases vigilance and neurocognitive performance. There is substantial evidence that lower levels of attentiveness have a negative influence on performance. Demanding call schedules, rising patient numbers, and a caregiver scarcity all add to the importance of weariness in medical practise. Recurrent sleep deprivation has been linked to severe neuroendocrine disruption in humans, according to a growing body of evidence. The Institute of Medicine's report, which estimates that medical errors cause up to 98,000 fatalities each year, has fueled research on the regulation of vigilance and tiredness management. The therapeutic importance of sleep-related changes in autonomic regulation has aided in the development of sleep medicine. Anaesthesiologists are incorporating advances in sleep medicine into their practise. The growing overlap between anesthesiology and sleep research necessitates a review that focuses on elements of sleep neurobiology that are particularly relevant to anesthesiology. This article reviews sleep phenomenology; arousal state regulation by multiple brain regions; the cholinergic model of rapid eye movement (REM) sleep; the regulation of traits that define arousal states; and neurochemical modulation of arousal states by acetylcholine, adenosine, -aminobutyric acid (GABA), monoamines, and hypocretin/orexin. The homeostatic management of sleep is demonstrated by this rebound rise. Normal sleep has a dynamic architecture, and its temporal organisation must be retained if sleep is to offer a pleasant and refreshing subjective experience. Any number of physiologic signs could have been chosen to name the two major phases of sleep. For example, REM sleep also is characterized by the presence of an activated cortical electroencephalogram.

Importance of Research: (200 words)

Anesthesiologists have claimed for more than 20 years that there is no one mechanism that causes states of anaesthesia. There is now widespread consensus that a single anaesthetic action mechanism cannot account for the physiologic and behavioural characteristics that distinguish anaesthetic states. Anesthesia and sleep are two distinct states with strikingly comparable

physiologic and behavioural characteristics. Anaesthetic and sleep share so many characteristics that patients are frequently assured that anaesthesia will put them to sleep. Sleep is a soothing metaphor for an altered arousal state brought on by hazardous chemicals, many of which have startlingly identical ED50 and LD50 values. Clinical and preclinical studies concur that anatomically scattered and chemically diverse neurons are responsible for spontaneously emerging states of arousal. The idea that neural networks evolved to govern naturally occurring sleep selectively modulate features that define sedation and anaesthesia has gotten a lot of support. The idea that specific neural functions are localised to specific brain regions is based on more than 150 years of clinical neurology. The intricacy of various brain regions contributing to the control of arousal states complicates efforts to comprehend the mechanisms by which sleep and anaesthesia reduce alertness. In, a schematic representation of the numerous brain areas involved in arousal state control is shown. Recent reviews provide in-depth information and diverse viewpoints on a variety of brain areas that regulate sleep and wakefulness. The deleterious effects of sleep deprivation on performance and state-dependent changes in neuronal excitability observed in preclinical investigations are consistent with brain region-specific variations in metabolism and blood flow revealed by advancements in sleep functional neuroimaging.

Biography: (200 words)

Fethiye Aylin Sungur is a General Surgeon with more than 30 years of experience as well as broad medical experience. She has excellent bedside manner and patient communication skills developed through more than three decades of combined schooling and teaching experience. She is the Chief of Minimally Invasive Surgery at San Raffaele Hospital, lead and assist in a variety of surgical procedures to address injuries, inflammatory and oncological diseases, communicate with patients and other medical professionals to create a treatment plan that includes preoperative preparations, surgical protocols and postoperative care and also prepare reports and other forms of documentation to keep patient charts updated around the clock during pre- and post-surgical hospital stays. She is also a Professor of Surgery and author of 88 paper published on PubMed and serves as an Editorial Board Member in various journals. During her bachelor degree. She was awarded multiple Gold medals during his student life. She is an active Medical Educator and continues to participate and present at various national and international meetings with more than 145 conference abstract presentations to her credit. He has deep interest for resuscitation, acute critical illness, trauma anesthesia, pediatric congenital cardiovascular anesthesia and perioperative patient safety.

Institute Photograph:



References:

1. Lydic R: The central regulation of sleep and autonomic physiology, *The Clinical Physiology of Sleep*. Edited by Lydic R, Biebuyck JF. Bethesda, American Physiological Society, 1988, pp 1–20 Lydic R, Biebuyck JF Bethesda, American Physiological Society
2. Dement WC, Mitler M: It's time to wake up to the importance of sleep disorders. *JAMA* 1993; 269:1548–50
3. [Kryger MH, Roth T, Dement WC: Principles and Practice of Sleep Medicine, 4th edition. Philadelphia, Elsevier Saunders, 2005, pp 1–1517 Philadelphia, Elsevier Saunders](#)
4. Loadsmann JA, Wilcox I: Is obstructive sleep apnoea a rapid eye movement-predominant phenomenon? *Br J Anaesth* 2000; 85:354–8
5. Loadsmann JA, Hillman DR: Anaesthesia and sleep apnoea. *Br J Anaesth* 2001; 86:254–66
6. [Chung F, Imarengiaye C: Management of sleep apnea in adults. Can J Anesth 2002; 49:R1–6](#)
7. Lichtor JL, Alessi R, Lane BS: Sleep tendency as a measure of recovery after drugs used for ambulatory surgery. *Anesthesiology* 2002; 96:878–83
8. Wu A, Drummond GB: Sleep arousal after lower abdominal surgery and relation to recovery from respiratory obstruction. *Anesthesiology* 2003; 99:1295–302
9. Fiset P: Research on anesthesia, consciousness or both? Understanding our anesthetic drugs and defining the neural substrate. *Can J Anesth* 2003; 50:R1–5
10. Brown KA, Lafemere BA, Moss IR: Recurrent hypoxemia in young children with obstructive sleep apnea is associated with reduced opioid requirement for analgesia. *Anesthesiology* 2004; 100:806–10
11. Backman SB, Fiset P, Plourde G: Cholinergic mechanisms mediating anesthetic induced altered states of consciousness. *Prog Brain Res* 2004; 145:197–206
12. den Herdner C, Schmeck J, Appleboom DJK, de Vries N: Risk of general anaesthesia in people with obstructive sleep apnoea. *BMJ* 2004; 329:955–9
13. Larijani GE, Goldberg ME, Hojat M, Khaleghi B, Dunn JB, Marr AT: Modafinil improves recovery after general anesthesia. *Anesth Analg* 2004; 98:976–81
14. Aserinsky E, Kleitman N: Regularly occurring periods of eye motility, and concomitant phenomena, during sleep. *Science* 1953; 118:273–4
15. Borbely AA: A two process model of sleep regulation. *Human Neurobiol* 1982; 1:195–204