

A Method for Purifying Native Transthyretin from Human Plasma

Aline G Santana^{1,2}, Paulo C Carvalho¹, Nilson IT Zanchin¹ and Tatiana ACB Souza^{1*}

¹Laboratory of Proteomics and Proteins Engineering, Institute Carlos Chagas, Curitiba-PR, Brazil

²Universidade Federal do Paraná (UFPR), Curitiba-PR, Brazil

Abstract

Transthyretin is a homotetrameric thyroid-hormone-transporting protein that binds to the retinol binding protein thus being involved in metabolism, growth, fertility, homeostasis of the cardiovascular and central nervous system, cell differentiation, reproduction, development and maintenance the cognitive processes during aging. Currently, there are several methodologies for natively purifying TTR from plasma, serum, tears, and amyloid fibrils; however, these procedures are laborious. Herein, a low-cost and simple protocol to purify TTR from human plasma is described. It involves the separation of plasma proteins by size exclusion and DEAE chromatography. The homogeneity was assessed by SDS-PAGE and by tandem mass spectrometry using an Orbitrap-XL (Thermo, San Jose-CA).

Keywords: Transthyretin; Plasma isolation

Introduction

Transthyretin was discovered in 1942 and originally named as pre albumin [1]; it is a highly conserved protein found in plasma and cerebrospinal fluid [1,2]. It is mainly expressed by hepatocytes [3] and epithelial cells of the choroid plexus [4]; yet it can also be expressed by the cerebral meninges [5], epithelial cells of the retina [6], pancreatic islets of Langerhans [7], visceral yolk sac [8], placenta [9], intestine [10] and at low-scale by stomach, heart muscle, and spleen [11].

The two major physiological functions of TTR are the transport of T4 hormone produced by thyroid and the transport of retinol (vitamin A) through interaction with retinol-binding protein RBP [12], thus being involved in metabolism, growth, fertility, and homeostasis of the cardiovascular and central nervous system [13], cell differentiation, reproduction, development and maintaining the cognitive processes during aging [14].

Previous reports describe evidences of TTR involvement in response to stress [15], in immunological pathways [16], metabolism of lipoproteins [17], and in neuroprotection against Alzheimer's disease by modulating the formation of β -amyloid plaques [18-20]. Paradoxically to this latter function, it is known that when some individuals inherit a mutated TTR gene, the mutated protein loses its functional structure and aggregates. This structural modification leads to formation of amyloid fibers and results in degenerative diseases that affect the nervous system, heart muscle, and other organs. Currently, the molecular mechanisms involved in the conversion of the TTR tetramer into aggregates of amyloid precursor fibers remains elusive. Although it is known that upon dissociation the TTR tetramer produces monomeric species with structural characteristics different from native monomer [21,22], the events associated with this non-native monomer in the initial process of TTR aggregation remain elusive.

The first reports for purifying TTR are from decades ago. Fex and Lindgren [23] purified a bovine counterpart to TTR from bovine serum by thiol-disulfide exchange chromatography on thiol-Sepharose 4B and affinity chromatography on human retinol-binding protein linked to Sepharose 4B. Berni et al. [24] purified TTR by using ammonium sulfate fractionation, followed by a hydrophobic interaction chromatography on phenyl-Sepharose and gel filtration on Sephadex G-50. Bashor et al. [25] developed a methodology for

TTR purification that involves precipitation of contaminating proteins with dilute aqueous phenol, ion-exchange chromatography on DEAE-Sephacel, and gel permeation chromatography on Sephadex G-100. Furuya et al. [26] were the first to recombinantly produce TTR fused with the *E. coli* outer membrane protein A (ompA) signal peptide. TTR has also been produced in the eukaryotic *Pichia Pastoris* system [27]. Lin et al. [28] created a method consisting of serum precipitation, anion exchange, thyroxine affinity chromatography and gel filtration. The latest methodologies for TTR purification include: 1) salting out, anion and cation exchange chromatographies, preparative electrophoresis, and size chromatography [29]; 2) ammonium sulfate fractionation followed by urea/Sephadex G-100 chromatography and a combination of two dye-affinity chromatographic steps on reactive yellow and cibacron blue coupled to agarose columns [30] and 3) extraction of TTR fibrils followed by sequential gel filtration after solubilization in a solution of guanidine hydrochloride and covalent chromatography [31]. This last method aims separating full-length TTR from C-terminal fragments found in TTR amyloid fibrils.

Aiming to establish a simple, fast and efficient method for purification of TTR, this work describes a purification strategy for TTR from blood plasma by employing two chromatographic methods in tandem. In our view, our method is one of the simplest yet for purifying TTR from a natural source; thus, ultimately aiding in studies that focus on the stabilization of the native tetramer for inhibition of their disassembly, prevention of amyloidogenic intermediates formation, inhibition of amyloidogenic intermediates aggregation or promotion of rupture of amyloid fibers [32-34]. A better understanding the TTR properties contribute to the development of more effective therapeutic strategies targeting pathologies related to this protein.

***Corresponding author:** Tatiana de Arruda Campos Brasil de Souza, Instituto Carlos Chagas, ICC- FIOCRUZ-PR, Rua Prof. Algacyr Munhoz Mader, 3775, bloco C, 81350-010, Curitiba-PR, Brazil, Tel: +55 41-2104-3225; E-mail: tatianabrasil@fiocruz.br

Received September 10, 2015; **Accepted** September 16, 2015; **Published** September 21, 2015

Citation: Santana AG, Carvalho PC, Zanchin NIT, Souza TACB (2015) A Method for Purifying Native Transthyretin from Human Plasma. *Biochem Pharmacol (Los Angel)* 4: 187. doi:10.4173/2167-0501.1000187

Copyright: © 2015 Santana AG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Material and Methods

Obtaining human plasma

Blood was collected from volunteer donors in collection tubes containing sodium citrate 3.2%. Blood samples were centrifuged at 15°C, 1500 rpm for 15 minutes. The supernatant (plasma) was collected and the pellet discarded.

Size exclusion chromatography

1 mL of human plasma obtained as described above was used as input for chromatography on Superdex 200 10/300 GL column. Sodium citrate 3.2% pH 8.0 was used as mobile phase for elution. The process was conducted on a FPLC system (Fast Performance Liquid Chromatography-GE-Healthcare). Eluted proteins were boiled in denaturant per 15 minutes and then loaded onto (15% acrylamide) as describe by [35].

Ion exchange chromatography

The fractions from gel filtration containing TTR were pooled and fractionated by ion exchange chromatography on DEAE Sepharose Fast Flow column with Sodium citrate 3.2% pH 8.0 (buffer A) and Sodium citrate 3.2% pH 8.0, 1M NaCl (Buffer B). Elution was performed with a linear gradient of 0-100% B in 10 column volumes with manual hold at each peak. The protein purification was analyzed by SDS PAGE (15% acrylamide) and by tandem mass spectrometry.

LC-MS/MS acquisition

The peptides were subjected to LC-MS/MS analysis using a Thermo Scientific Easy-nLC 1000 ultra-high performance liquid chromatography (UPLC) system coupled online to a LTQ-Orbitrap XL ETD mass spectrometer (Mass Spectrometry Facility-RPT02H PDTIS/ Carlos Chagas Institute-Fiocruz Paraná), as follows. The peptide mixtures were loaded onto a column (75 mm i.d., 15 cm long) packed in house with a 3.2 µm ReproSil-Pur C18-AQ resin (Dr. Maisch) with a flow of 500 nL/min and subsequently eluted with a flow of 250 nL/min from 5% to 40% ACN in 0.5% formic acid and 0.5% DMSO, in a 120 min gradient. The mass spectrometer was set in data-dependent mode to automatically switch between MS and MS/MS (MS2) acquisition. Survey full scan MS spectra (from m/z 300-2000) were acquired in the Orbitrap analyzer with resolution R=60,000 at m/z 400 (after accumulation to a target value of 1,000,000 in the linear trap). The ten most intense ions were sequentially isolated and fragmented in the linear ion trap. Previous target ions selected for MS/MS were dynamically excluded for 90 seconds. Total cycle time was approximately three seconds. The general mass spectrometric conditions were: spray voltage, 2.4 kV; no sheath and auxiliary gas flow; ion transfer tube temperature 100°C; collision gas pressure, 1.3 mTorr; normalized energy collision energy using wide-band activation mode; 35% for MS2. Ion selection thresholds were: 250 counts for MS2. An activation q=0.25 and activation time of 30 ms were applied in MS2 acquisitions.

Raw MS data analysis

The reviewed proteome set of Homo sapiens, composed of 20,187 sequences, was downloaded from the UniProt consortium on July 4th, 2014. PatternLab was used for generating a target-decoy database by grouping subset sequences, adding the sequences of 127 common mass spectrometry contaminants, and, for each sequence, including a reversed version of it. The final database used for PSM contained 105,551 sequences.

Peptide sequence matching

The Comet 2014 rev. 1 search engine [36], which is embedded into PatternLab for proteomics [37], was used to compare experimental tandem mass spectra against those theoretically generated from our sequence database and select the most likely peptide sequence candidate for each spectrum. Briefly, the search was limited to fully peptide candidates; we imposed carbamidomethylation of cysteine and oxidation of Methionine as fixed and variable modification, respectively. The search engine accepted peptide candidates within a 40-ppm tolerance from the measured precursor m/z and used the XCorr as the primary search engine score.

PSM validity was assessed using the search engine processor (SEPro) [38], which is embedded in PatternLab version 3.0.0.34. Briefly, identifications were grouped by charge state (+2 and > +3) resulting in two distinct subgroups. For each result, the Comet XCorr, DeltaCN, DeltaPPM, and Peaks Matched values were used to generate a Bayesian discriminator. The identifications were sorted in non-decreasing order according to the discriminator score. A cutoff score was established to accept a false-discovery rate (FDR) of 1% at the peptide level based on the number of labeled decoys. This procedure was independently performed on each data subset, resulting in an FDR that was independent of charge state. Additionally, a minimum sequence length of six amino-acid residues was required. Results were post-processed to only accept peptide spectrum matches with less than 6 ppm from the global identification average and proteins with two or more identified peptides.

Protein relative quantitation by extracted ion chromatograms

Relative quantitation by mass spectrometry describes strategies for comparing quantitative information of a same analyte between multiple samples. On the other hand, obtaining absolute quantitation values by mass spectrometry is challenging; some inherent difficulties are: a) different molecules ionize with different efficiencies in the mass spectrometer b) each protein will result in a different number of peptides after tryptic digestion. In a previous report, Zybaylov et al described a strategy for normalizing spectral counting quantitation data that provides quantitation values closer to absolute values [39]. We recall that spectral counting is a label-free strategy that correlates the total number of MS/MS spectra assigned to a protein with its relative abundance. Briefly, Zybaylov's normalization, the so-called Normalized Spectral Abundance Factor (NSAF), considers the fact that longer proteins tend to have more peptide identifications than shorter ones. Formally, the NSAF for a protein k is given by "the number of spectral counts (SpC) identifying a protein, k, divided by the protein's length (L), and divided by the sum of SpC / L for all N proteins in the experiment [39].

In this work, we employed a modified relative quantitation, having roots in NSAF, to assess the effectiveness of our purification approach. Briefly, instead of relying on spectral counts, or strategy, here termed Normalized Ion Abundance Factor (NIAF), replaces spectral count values by extracted ion chromatograms (XIC) values. Briefly, we recall that XICs are obtained by plotting the intensity of a particular peptide ion (m/z) over time and then integrating the area under the curve; for more on quantitative proteomic strategies we refer the reader to [http://dx.doi.org/10.1155/2013/674282]. XICs were obtained by using PatternLab's SEProQ module [37]. Our motivation in doing so is that protein ratios obtained by XICs yield more accurate estimates ratios than those obtained by spectral counting and therefore yielding estimates closer to the absolute values.

Results

Size exclusion chromatography (SEC), a chromatography that is widely used for purification and determination of the hydrodynamic radius of molecules [40], was the choice as first step for plasma fractionation. Human plasma, when chromatographed through Superdex 200 10/300 column is resolved into five distinct protein peaks (Figure 1).

The SDS-PAGE electrophoresis profile suggested that each peak from superdex 200 10/300 contained different blood proteins. In the fractions corresponding to the shoulder of the third peak, it was possible to verify the presence of a protein with expected size for TTR (Figure 1). Fractions containing the TTR were pooled and submitted to ion exchange chromatography on a DEAE column. IEC resulted in four distinct peaks (Figure 2A). The proteins were well separated and, in the fourth peak, it is possible to identify TTR by mass spectroscopy with two minor contaminants detected. The active TTR in the Uniprot has ~13.9 kDa and is composed of 127 amino acids. The sequence coverage was 88% and peptides comprising 112 aminoacids out of 127 were detected. We obtained NSIAF from all proteins, as detailed in the methodology; our results showed that 92.5% of the total NSIAF from the sample was decurrently from TTR (Figure 2B). In that same fraction, 4.9% corresponded to the keratin, a common contamination that happens during sample preparation, and 2.4% refers to other proteins.

All our raw mass spectrometric data is made available at <http://proteomics.fiocruz.br/tatiana/2015/>. Moreover, TTR was not detected

in any other fraction indicating that all TTR present in plasma is concentrated in this fraction. The yield of purification was ~0.09 mg of protein per ml of plasma. Our experimental procedure also could be helpful to separate albumin, Serotransferrin and Antithrombin-III from other blood proteins (Figure 2).

Discussion

Several reports of plasma protein purification by size exclusion chromatography are available; but none were performed with the type of column and buffer system used in this work. By comparing the chromatographic profile of the gel filtration of human plasma presented in Figure 1 with other works, it is possible to observe that the chromatographic profile can vary widely for the same sample at the expense the column used [41,42].

Sephacrose 4 FF is widely used for plasma fractionation [43], but due to its low resolution, it is most suitable in cases where the target protein and most contaminants have very distinct molecular masses. Other columns such as Superose 6, Sephadex G-200, Sephadex G-100 also appear in the literature for plasma proteins fractionation [43-46], but they all have lower resolution than Superdex 200 10/300. Sephadex G-200, the column closest to the required resolution, is capable of providing three distinct peaks when used in the first plasma fractionation step [44]. However, its use at the expense of Superdex 200 10/300, would probably require additional purification steps to obtain pure TTR due to higher amounts of remnant contaminants.

Another important aspect to be noted is that the same column and





Figure 1: Chromatogram of plasma fractionation by gel filtration on a Superdex 200 10/300 column and SDS PAGE of the eluted fractions during gel filtration. The fractions containing TTR are evidenced by the red box. First lane- Molecular weight marker (Bench marker protein ladder). The arrow (←) indicates the gel bands expected to be TTR.



sample can result in slightly different chromatographic profiles when using buffers with different pH and ionic strength. In this work, the buffer composed of 3.2% Sodium Citrate pH 8.0 was used in all steps of protein purification. The majority of the previously solutions used for plasma purification were buffered with Tris-HCl [23,41,42]. The choice of 3.2% Sodium Citrate was fundamental to better separation of plasma proteins since this buffer prevents the activation of the coagulation cascade, by sequestering calcium, and it is a good buffering at a pH close to physiological, favoring proteins to maintain the conformation found in plasma. Our experimental procedure also could be helpfull to separate albumin, Serotransferrin and Antithrombin-III from other blood proteins (Figure 2). Albumin was found in the second peak of IEC (Figure 2). This protein represents 50% -70% of all plasma proteins [45], and have a remarkable ability to bind to exogenous and endogenous components [46], making its removal important for application related to the remaining plasma proteins.

This method can also effectively be used to isolate two other important proteins: serotransferrin and antithrombin-III (Figure 2). Serotransferrin is associated with Fe³⁺ uptake, distribution and control of its quantity [47]. Antithrombin-III is an endogenous anticoagulant capable of inhibit key-factors of the coagulation cascade such as thrombin and FXa, thus, having a crucial role in homeostasis [42]. About 25% of the total NSIAF from peak 1 and peak 3 (Figures 2A and 2C) was decurrently from serotransferrin and antithrombin-III, respectively. These proteins can be further purified through additional chromatographic steps. All these results show a simple and low-cost method for chromatographic purification of human plasma TTR. The procedure is also helpfull to remove albumin and to purify two other blood proteins. TTR are often submitted to drug development screenings due to its role in important pathologies such as Senile Systemic Amyloidosis (SSA) and Familial Amyloid Polyneuropathy (FAP) [48] with some promising amyloidogenesis inhibitors [49-54] such as genistein [54,55], other flavonoids of similar structure [55], T4

analogues [49,52,53] being indentified. However all screenings for such compounds were performed with recombinant molecules.

References

1. Kabat EA, Moore DH, Landow H (1942) An Electrophoretic Study Of The Protein Components In Cerebrospinal Fluid And Their Relationship To The Serum Proteins. *J Clin Invest* 21: 571-577.
2. Seibert FB, Nelson JW (1942) Electrophoretic study of the blood response in tuberculosis. *J Biol Chem* 143: 29-38
3. Murakami T, Yasuda Y, Mita S, Maeda S, Shimada K, et al. (1987) Prealbumin gene expression during mouse development studied by in situ hybridization. *Cell Differ* 22: 1-9.
4. Aleshire SL, Bradley CA, Richardson LD, Parl FF (1983) Localization of human prealbumin in choroid plexus epithelium. *J Histochem Cytochem* 31: 608-612.
5. Herbert J, Wilcox JN, Pham KT, Fremreau RT Jr, Zeviani M, et al. (1986) Transthyretin: a choroid plexus-specific transport protein in human brain. The 1986 S. Weir Mitchell award. *Neurology* 36: 900-911.
6. Martone RL, Herbert J, Dwork A, Schon EA (1988) Transthyretin is synthesized in the mammalian eye. *Biochem Biophys Res Commun* 151: 905-912.
7. Kato M, Kato K, Blaner WS, Chertow BS, Goodman DS (1985) Plasma and cellular retinoid-binding proteins and transthyretin (prealbumin) are all localized in the islets of Langerhans in the rat. *Proc Natl Acad Sci U S A* 82: 2488-2492.
8. Soprano DR, Soprano KJ, Goodman DS (1986) Retinol-binding protein and transthyretin mRNA levels in visceral yolk sac and liver during fetal development in the rat. *Proc Natl Acad Sci U S A* 83: 7330-7334.
9. McKinnon B, Li H, Richard K, Mortimer R (2005) Synthesis of thyroid hormone binding proteins transthyretin and albumin by human trophoblast. *J Clin Endocrinol Metab* 90: 6714-6720.
10. Loughna S, Bennett P, Moore G (1995) Molecular analysis of the expression of transthyretin in intestine and liver from trisomy 18 fetuses. *Hum Genet* 95: 89-95.
11. Soprano DR, Herbert J, Soprano KJ, Schon EA, Goodman DS (1985) Demonstration of transthyretin mRNA in the brain and other extrahepatic tissues in the rat. *J Biol Chem* 260: 11793-11798.
12. Kanai M, Raz A, Goodman DS (1968) Retinol-binding protein: the transport protein for vitamin A in human plasma. *J Clin Invest* 47: 2025-2044.

13. Hulbert AJ (2000) Thyroid hormones and their effects: a new perspective. *Biol Rev Camb Philos Soc* 75: 519-631.
14. Brouillette J, Quirion R (2008) Transthyretin: a key gene involved in the maintenance of memory capacities during aging. *Neurobiol Aging* 29: 1721-1732.
15. Bernstein LH, Leukhardt-Fairfield CJ, Pleban W, Rudolph R (1989) Usefulness of data on albumin and prealbumin concentrations in determining effectiveness of nutritional support. *Clin Chem* 35: 271-274.
16. Burton PM, Hung P, Lin T, Lovelace C, White A (1985) The effects of homogeneous human prealbumin on in vitro and in vivo immune responses in the mouse. *Int J Immunopharmacol* 7: 473-481.
17. Sousa MM, Berglund L, Saraiva MJ (2000) Transthyretin in high density lipoproteins: association with apolipoprotein A-I. *J Lipid Res* 41: 58-65.
18. Riisøen H (1988) Reduced prealbumin (transthyretin) in CSF of severely demented patients with Alzheimer's disease. *Acta Neurol Scand* 78: 455-459.
19. Schwarzman AL, Gregori L, Vitek MP, Lyubski S, Strittmatter WJ, et al. (1994) Transthyretin sequesters amyloid beta protein and prevents amyloid formation. *Proc Natl Acad Sci U S A* 91: 8368-8372.
20. Serot JM, Christmann D, Dubost T, Couturier M (1997) Cerebrospinal fluid transthyretin: aging and late onset Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 63: 506-508.
21. Quintas A, Vaz DC, Cardoso I, Saraiva MJ, Brito RM (2001) Tetramer dissociation and monomer partial unfolding precedes protofibril formation in amyloidogenic transthyretin variants. *J Biol Chem* 276: 27207-27213.
22. Lindgren M, Sörgjerd K, Hammarström P (2005) Detection and characterization of aggregates, prefibrillar amyloidogenic oligomers, and protofibrils using fluorescence spectroscopy. *Biophys J* 88: 4200-4212.
23. Fex G, Lindgren R (1977) Purification of a bovine counterpart to human prealbumin and its interaction with a bovine retinol-binding protein. *Biochim Biophys Acta* 493: 410-417.
24. Berni R, Ottonello S, Monaco HL (1985) Purification of human plasma retinol-binding protein by hydrophobic interaction chromatography. *Anal Biochem* 150: 273-277.
25. Bashor MM, Hewett J, Lackey A, Driskell WJ, Neese JW (1987) Purification of prealbumin from human serum. *Prep Biochem* 17: 209-227.
26. Furuya H, Nakazato M, Saraiva MJ, Costa SP, Sasaki H, et al. (1989) Tetramer formation of a variant type human transthyretin (prealbumin) produced by *Escherichia coli* expression system. *Biochem Biophys Res Commun* 163: 851-859.
27. Prapunpoj P, Leelawatwatana L, Schreiber G, Richardson SJ (2006) Change in structure of the N-terminal region of transthyretin produces change in affinity of transthyretin to T4 and T3. *FEBS J* 273: 4013-4023.
28. Lin Q, Su T, Liu G, Gu J (1996) Purification of transthyretin by high performance affinity chromatography from human plasma. *Prep Biochem Biotechnol* 26: 245-257.
29. Blirup-Jensen S (2013) Protein standardization I: Protein purification procedure for the purification of human prealbumin, orosomucoid and transferrin as primary protein preparations. *Clin Chem Lab Med* 39: 1076-1089.
30. Raghu P, Ravinder P, Sivakumar B (2003) A new method for purification of human plasma retinol-binding protein and transthyretin. *Biotechnol Appl Biochem* 38: 19-24.
31. Westermark P, Westermark GT (2005) Purification of transthyretin and transthyretin fragments from amyloid-rich human tissues. *Methods Mol Biol* 299: 255-260.
32. Damas AM, Saraiva MJ (2000) Review: TTR amyloidosis-structural features leading to protein aggregation and their implications on therapeutic strategies. *J Struct Biol* 130: 290-299.
33. Saraiva MJ (2002) Hereditary transthyretin amyloidosis: molecular basis and therapeutic strategies. *Expert Rev Mol Med* 4: 1-11.
34. Mason JM, Kokkoni N, Stott K, Doig AJ (2003) Design strategies for anti-amyloid agents. *Curr Opin Struct Biol* 13: 526-532.
35. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
36. Eng JK, Jahan TA, Hoopmann MR (2013) Comet: an open-source MS/MS sequence database search tool. *Proteomics* 13: 22-24.
37. Carvalho PC, Fischer JS, Xu T, Yates JR 3rd, Barbosa VC (2012) PatternLab: from mass spectra to label-free differential shotgun proteomics. *Curr Protoc Bioinformatics Chapter 13: Unit13*.
38. Carvalho PC, Fischer JS, Xu T, Cociorva D, Balbuena TS, et al. (2012) Search engine processor: Filtering and organizing peptide spectrum matches. *Proteomics* 12: 944-949.
39. Zybailov B, Mosley AL, Sardi ME, Coleman MK, Florens L, et al. (2006) Statistical analysis of membrane proteome expression changes in *Saccharomyces cerevisiae*. *J Proteome Res* 5: 2339-2347.
40. Mory S, Barth HG (1999) Size exclusion chromatography. Springer, Berlin.
41. JosiÅž D, Horn H, Schulz P, Schwinn H, Britsch L (1998) Size-exclusion chromatography of plasma proteins with high molecular masses. *J Chromatogr A* 796: 289-298.
42. Kaersgaard P, Barington KA (1998) Isolation of the factor VIII-von Willebrand factor complex directly from plasma by gel filtration. *J Chromatogr B Biomed Sci Appl* 715: 357-367.
43. Mourey L, Samama JP, Delarue M, Choay J, Lormeau JC, et al. (1990) Antithrombin III: structural and functional aspects. *Biochimie* 72: 599-608.
44. Perrier H, Delcroix JP, Perrier C, Gras J (1973) An attempt to classify the plasma proteins of the rainbow trout (*salmo gairdnerii richardson*) using disc electrophoresis, gel filtration and salt solubility fractionation. *Comp. Biochem. Physiol* 46: 475-482.
45. Silverlight JJ, Prysor-Jones RA, Jenkins JS (1985) Growth hormone in normal female rat plasma appears on gel filtration as a large molecular weight form. *Life Sci* 36: 1927-1932.
46. Hu T, Liu Y (2015) Probing the interaction of cefodizime with human serum albumin using multi-spectroscopic and molecular docking techniques. *J Pharm Biomed Anal* 107: 325-332.
47. Fanali G, di Masi A, Trezza V, Marino M, Fasano M, et al. (2012) Human serum albumin: from bench to bedside. *Mol Aspects Med* 33: 209-290.
48. Chasteen DN (1977) Human serotransferrin: Structure and Function. *Coord Chem Rev* 22: 1-36.
49. Mourey L, Samama JP, Delarue M, Choay J, Lormeau JC, et al. (1990) Antithrombin III: structural and functional aspects. *Biochimie* 72: 599-608.
50. Palmieri Lde C, Lima LM, Freire JB, Bleicher L, Polikarpov I, et al. (2010) Novel Zn²⁺-binding sites in human transthyretin: implications for amyloidogenesis and retinol-binding protein recognition. *J Biol Chem* 285: 31731-31741.
51. Miroy GJ, Lai Z, Lashuel HA, Peterson SA, Strang C, et al. (1996) Inhibiting transthyretin amyloid fibril formation via protein stabilization. *Proc Natl Acad Sci U S A* 93: 15051-15056.
52. Peterson SA, Klabunde T, Lashuel HA, Purkey H, Sacchetti JC, et al. (1998) Inhibiting transthyretin conformational changes that lead to amyloid fibril formation. *Proc Natl Acad Sci U S A* 95: 12956-12960.
53. Baures PW, Oza VB, Peterson SA, Kelly JW (1999) Synthesis and evaluation of inhibitors of transthyretin amyloid formation based on the non-steroidal anti-inflammatory drug, flufenamic acid. *Bioorg Med Chem* 7: 1339-1347.
54. Klabunde T, Petrassi HM, Oza VB, Raman P, Kelly JW, et al. (2000) Rational design of potent human transthyretin amyloid disease inhibitors. *Nat Struct Biol* 7: 312-321.
55. Petrassi HM, Johnson SM, Purkey HE, Chiang KP, Walkup T, et al. (2005) Potent and selective structure-based dibenzofuran inhibitors of transthyretin amyloidogenesis: kinetic stabilization of the native state. *J Amer Chem Soc* 127: 6662-6671.
56. Green NS, Foss TR, Kelly JW (2005) Genistein, a natural product from soy, is a potent inhibitor of transthyretin amyloidosis. *Proc Natl Acad Sci U S A* 102: 14545-14550.
57. Trivella DBB, Bleicher L, Palmieri LC, Wiggers HJ, Montanari CA, et al. (2010) Conformation differences between the wild type and V30M mutant transthyretin modulate its binding to genistein: implications to tetramer stability and ligand-binding. *J Struct Biol* 170: 522-531.