

Altered signal transduction and cellular permeability in retinal bipolar cells caused by mutated TRPM1 and MYO7A could reveal new insights of retinal degeneration pathways

Luigi Donato^{1,2}, Concetta Scimone^{1,2}, Simona Alibrandi^{1,3}, Carmela Rinaldi¹, Rosalia D'Angelo^{1,2}, Antonina Sidoti^{1,2}

¹Department of Biomedical and Dental Sciences and Morphofunctional Images, Division of Medical Biotechnologies and Preventive Medicine, University of Messina, Messina, Italy;

²Department of Biomolecular strategies, genetics and avant-garde therapies, I.E.M.E.S.T., Palermo, Italy;

³Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Messina, Italy.

Abstract:

The earliest step of the visual processing is the generation of parallel information channels responding to increases versus decreases in light intensity. Such ON and OFF responses begin at the first retinal synapse where two classes of postsynaptic bipolar cells react with opposite polarities to glutamate released by photoreceptors. While dendrites of OFF-bipolar cells contain ionotropic glutamate receptors of the AMPA/kainite class, the ON-bipolar cell dendrites express a unique metabotropic glutamate receptor 6 (mGluR6). TRPM1 is a component of the transduction cation channel negatively regulated by the mGluR6 cascade in ON bipolar cells, and forms a macromolecular complex with other proteins including the just cited mGluR6, GPR179, nyctalopin, and the regulator of G protein signaling proteins. Mutations of human TRPM1 are associated with hereditary and acquired diseases in which the retinal ON pathway is selectively affected, such as congenital stationary night blindness. It represents a clinically and genetically heterogeneous group of retinal disorders, whose affected patients lack rod function and suffer from night blindness starting in early childhood. We present data coming from whole exome sequencing of a family in which 2 sons were diagnosed for an orphan form of retinal dystrophy, even if characterized by two different phenotypes. Both patients presented a causative mutation of Usher Syndrome in MYO7A gene, but only one showed a causative mutation of CSNB in TRPM1 gene

(c.470C>T, Ser157Phe). We evaluated possible consequences of identified variants on each corresponding protein, analyzing their possible interaction and biological processes that their alterations could impair.

Keywords: Biology, Cell documents, Gene therapy

Results:

Histological examination of the removed appendix has shown "Adenocarcinoma of the appendix in the base of epithelial adenoma, with negative surrounding lymph nodes". Following the histopathology report the patient has underwent an elective right hemicolectomy at a later stage.

Conclusions:

The neoplasms of the appendix are rare (1% of the cases) and are usually expressed as acute appendicitis. Adenocarcinomas correspond to 10% of the cases of tumors of the appendix whereas there are three histological types: mucosal (more often), intestinal (colonic) and Signet- ring cell carcinoma. The treatment always involves surgical intervention, right hemicolectomy, which gives a better five year survival rate (73%) compared to appendicectomy (44%). In some cases a simple appendicectomy was the treatment of choice if the margins of the tumor stop at the mucosal or sub-mucosal layer. Chemotherapy has not been proved to contribute to treatment.