

An Update on the Sense of Taste in Chickens: A Better Developed System than Previously Appreciated

Hong-Xiang Liu*, Prasangi Rajapaksha, Zhonghou Wang, Naomi E Kramer and Brett J Marshall

Regenerative Bioscience Center, Department of Animal and Dairy Science, College of Agricultural and Environmental Sciences, University of Georgia, USA

Abstract

Taste is important in guiding nutritive choices and motivating food intake. The sensory organs for taste are the taste buds, that transduce gustatory stimuli into neural signals. It has been reported that chickens have a low taste bud number and thus low taste acuity. However, more recent studies indicate that chickens have a well-developed taste system and the reported number and distribution of taste buds may have been significantly underestimated. Chickens, as a well-established animal model for research, are also the major species of animals in the poultry industry. Thus, a clear understanding of taste organ formation and the effects of taste sensation on nutrition and feeding practices is important for improving livestock production strategies. In this review, we provide an update on recent findings in chicken taste buds and taste sensation indicating that the chicken taste organ is better developed than previously thought and can serve as an ideal system for multidisciplinary studies including organogenesis, regenerative medicine, feeding and nutritional choices.

Keywords: Chicken; Taste; Poultry; Feed; Nutrition

Introduction

Taste sensation is conserved in vertebrate animals – most species have a well-developed taste system comprised of taste sensory organs, the innervating nerves, and the central nervous system. In this review we will focus on the sense of taste in chickens, including their sensory organs and behavioral responses.

Recent studies have advanced our understanding of basic information concerning chicken taste bud number, distribution, structure, and development. Using molecular markers to label chicken taste buds in the oral epithelial sheet, many more taste buds have been observed. This indicates that chickens have a better developed taste system and thus a larger impact of taste on their feeding behavior than previously appreciated.

Sensory Organs for Taste in Chickens

The sensory organs for taste are taste buds, which detect different types of tastants and transduce gustatory stimuli into neural signals conveyed to the brain for taste perception. Among different species, taste bud distribution varies. For example, mammalian taste buds are primarily located in the tongue, though they are also observed in the soft palate, epiglottis, pharynx, larynx, uvula [1-4]. In contrast, the avian taste organ system is a prominent example of a non-lingual taste system. Chicken taste buds differ from those of mammals in many respects.

Taste bud number, distribution, and structure

Initially, it was reported that chickens do not have taste buds [5] and later about 70 taste buds were found in the oral cavity [6]. This number is low compared to mammals, e.g., rats (~1000 taste buds), humans (~10,000 taste buds) [1] and cattle, which have about 15,000-20,000 taste buds [7]. Further studies demonstrated that chickens have a higher number of taste buds varying in number from 240-360 on average according to the breed, e.g., broilers have more taste buds compared to the layer-type [8-10]. In addition, our recent studies have demonstrated that the number of taste buds varies among lines in the same type (broiler-type) of chickens at P3, i.e., female line males have more taste buds in the base of the oral cavity than females and male line males [11].

Chicken taste buds are present in both the anterior and posterior regions of the oral cavity, primarily distributed in three regions of oral epithelium – the palate (~69%) (anterior maxillary gland opening region, middle palatine papilla region, and posterior region), the base of the oral cavity (anterior mandibular gland region) (~29%), and posterior ventrolateral regions of the keratinized anterior tongue and posterior region of the tongue (region posterior to the lingual spine) (~2%) [8,9]. In the oral epithelium, taste buds are mainly located near salivary gland openings (>20 µm in diameter) [8,11]. The two clusters of taste buds in the anterior maxillary gland opening region of the palate are large and dense which may be important for immediate detection as soon as the feed enters the oral cavity. The lower number of lingual taste buds in chickens suggests that the tongue is not the primary organ for taste in chickens; rather it primarily facilitates food processing.

Chicken taste buds, comprised of a cluster of specified fusiform cells, are ovoid (“egg” shaped), which is in contrast to mammals whose taste buds are onion-like/bud-shaped. Using 2-photon microscopy in the oral epithelial sheet immunostained with molecular markers, large tube-like taste buds were also seen in the posterior region of the palate [11]. Unlike mammals, chickens do not have specialized structures like lingual taste papillae (i.e., fungiform, foliate, circumvallate) to host taste buds. The taste buds are embedded in the epithelium, and grouped in clusters that surround the salivary gland openings in a rosette pattern [8,11]. Regarding the distribution of taste buds, we found that taste buds in the base of oral cavity extend to the lateral edge [11], indicating a broader distribution than previously observed.

***Corresponding author:** Hong-Xiang Liu, Assistant Professor, Regenerative Bioscience Center, Department of Animal and Dairy Science, College of Agricultural and Environmental Sciences, University of Georgia, 425 River Road, 30602, USA, Tel: 1-706-5427048; Fax: 706-583-0274; E-mail: lhx@uga.edu

Received February 26, 2018; Accepted March 16, 2018; Published March 31, 2018

Citation: Liu HX, Rajapaksha P, Wang Z, Kramer NE, Marshall BJ (2018) An Update on the Sense of Taste in Chickens: A Better Developed System than Previously Appreciated. J Nutr Food Sci 8: 686. doi: [10.4172/2155-9600.1000686](https://doi.org/10.4172/2155-9600.1000686)

Copyright: © 2018 Liu HX, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The previous data on the number and distribution of taste buds have been obtained from the observation of taste pores (2-10 μm in diameter) with scanning electron microscopy [8]. More recently, we used molecular markers, α -Gustducin and Vimentin, to label chicken taste buds in the peeled epithelial sheet and found many more taste buds than previously reported [11]. The numbers vary by gender even in the same breed (COBB 500) – female-line male chickens have more taste buds than females and male-line males. The broiler-type female-line male chickens have up to 500 taste buds in the palate and ~260 in the base of oral cavity at P3, which is much higher than the previously reported number with scanning electron microscopy (218 in palate and 91 in the base of oral cavity) [8]. It has been reported that the ratio of salivary gland openings to taste buds in chickens is around 1:2.5, a ratio which does not change with age [9]. However, using molecular markers to label taste buds, larger clusters are observed in the base of the oral cavity, e.g., 4 buds per cluster on average (up to 14) in males and 3 buds in females on average (up to 9) [11]. The high number of chicken taste buds implicates a better-developed taste organ system and potentially larger than previously appreciated impact of taste on feeding behaviors in birds.

Previous studies have shown that the total number of taste buds is similar in young and adult chickens [12]. However, a more detailed analysis showed that the numbers of taste buds and clusters change with age from P0-P8. For instance, taste bud number in broiler-type, female line males increases after hatch, peaks at P3, and then decreases at later stages. The continuing development of taste buds in chicks after P0 provides a time window when taste bud formation, and therefore taste sensing, can be modified in early hatched chickens [11].

Taste bud cell types

Similar to mammals, the chicken taste bud cell population is heterogeneous histologically [8,13]. In mammals, there are four types of taste bud cells, type I cells (dark), type II cells (light) that are considered to be “receptor cells”, type III (intermediate) and type IV (basal) cells. In chickens, different taste bud cell types have been identified based on ultra-structural studies using transmission electron microscopy.

At least four taste bud cell types have been classified based on appearance, including basal cells, dark cells, light cells, and flattened/intermediate cells [14]. The dark cells, the most abundant cell type in chicken taste buds, have cytoplasmic extensions (similar to microvilli in mammalian taste bud cells) and their main function is regarded as support of the taste bud [15]. In addition, dark cells have dense cytoplasm with scattered chromatin and fewer vesicles. The light cells, similar to type II taste bud cells in mammals, have less dense cytoplasm but more vesicles compared to dark cells. The intermediate cells have certain degrees of characteristics of both light and dark cells [14]. Basal cells, as the name indicates, are located in the basal region of the taste buds, have an irregular shaped nucleus, dense cytoplasm and are darker than the other cell types [14].

Specific markers for the three types of differentiated taste bud cells in rodents have been well-characterized, e.g., NTPDase II for type I [16], α -Gustducin for type II [17,18], SNAP25 for type III [19]. However, there are not established molecular markers to identify the specific taste cell types in chickens, although molecular markers (Vimentin and α -Gustducin) are available to label taste buds [11,20-22].

Development and maintenance of taste buds

Chicken taste bud development: Chicken beaks and tongue, the regions where the oral tissues hosting taste buds are located, develop

by embryonic day (E) 8, and taste buds emerge at E17 (Hamburger Hamilton stage 43) as spherical cluster of cells in the base of the epithelium. From E17-18, the cell clusters further develop and increase in number, though without a pore penetrating the surface of the epithelium. At E19, taste bud cells elongate and form an ovoid structure with a narrow and shallow taste pore penetrating the surface of the epithelium [12]. Compared to rodents, chicken taste buds develop and mature early, i.e., before hatch [23]. In rodents, taste bud development and maturation occur after birth, and taste bud maturation is completed 2-3 weeks after birth [24].

At E19, the number of taste buds peaks and has been reported to remain consistent after hatch in most chickens [12]. It has been assumed that taste bud number remained relatively static in young chicks versus adults, with no age-related effects on taste bud development [25]. However, as discussed earlier, in broiler-type Cobb 500 chickens, the female-line males continue to develop taste buds after hatch, peaking at P3, and then decreasing to a stable level [11].

Chicken taste bud maintenance: Taste buds reside in the epithelium and contain cells which possess epithelial features – they have a short life span and undergo continuous turnover. It has been suggested that the cell turnover rate depends on the species, age, and location of the taste buds [26,27]. In chickens, the average life span of taste bud cells is shorter compared to other vertebrates that have been reported so far. The life span of chicken taste bud cells in the base of the oral cavity is 3-4 days on average [26,27], in contrast to 7-14 days in mammals [26,28,29]. The high turnover rate of taste bud cells requires a rigorous progenitor/stem cell niche in order to maintain proper taste function. Type IV/basal cells are regarded as one of these stem cell niches. In addition, the “edge/perigemmal” cells immediately surrounding taste buds are highly proliferative which is implicit for their function in taste bud cell renewal.

Our recent studies have demonstrated a population of proliferating cells within chicken taste buds that are primarily not labeled by other known chicken taste cell markers (i.e., Vimentin and α -Gustducin), implying that these are a distinct and undifferentiated population of taste cells. This is unique because in rodents, taste bud cells are largely post-mitotic – their proliferating cells existing chiefly in the surrounding epithelium [30] and potentially the connective tissue [31]. The unique localization of proliferating cells in chicken taste buds, lining its basal layer, suggests the possibility that the taste buds are predominantly independent structures with “built-in” progenitors to meet the needs of their large size and rapid turnover. However, proliferating cells are also located in other tissue compartments such as the epithelium and connective tissue, so lineage tracing must be carried out to confirm this conjecture.

Regulation of taste bud development and maintenance: Taste organs, like other epithelial appendages, require epithelial-mesenchymal interactions and involvement of multiple signaling pathways for proper formation. Sonic hedgehog (Shh) and bone morphogenetic proteins (BMP) signaling cascades have been identified in the development of the avian tongue [32], where a small population of taste buds are located. The signaling mechanisms underlying chicken taste bud development are largely unknown.

Knowledge regarding the regulation of taste organ development in rodents may give us perspectives and be beneficial for in-depth studies in chickens. In rodents, numerous molecular pathways have been identified that regulate the development of embryonic tongue, developing taste papillae, and taste buds. For example, e.g., Hedgehog

[33-38], Wnt/ β -catenin [39-43] TGF- β /BMP [44-47], Notch [48-50] fibroblast growth factor (FGF) [45,51,52], and Erbb [53-55].

Recently, we used RNA-Seq analysis to map the transcriptomic architecture of developing gustatory tissues in chickens [56]. Although chickens have taste buds throughout the oral cavity, including the epithelia of the palate, base of the oral cavity, and posterior tongue, analyses between the epithelia and the underlying mesenchyme of each tissue demonstrated that the epithelium of the base of the oral cavity contained the most differentially expressed genes (DEGs) related to taste, including but not limited to GNAT3 and TAS1R3. A deeper analysis of this tissue using regions of the epithelium containing (gustatory) or not containing (non-gustatory) taste buds, as well as the mesenchyme situated beneath the gustatory epithelium, revealed components of several pathways involved in organogenesis that were differentially expressed. Indeed, TGF β /BMP, FGF, Notch, SHH, and ERbb signaling cascades were differentially expressed between gustatory epithelium and gustatory mesenchyme, implicating the regulatory roles of these tissues in taste bud development and maintenance.

In summary, chicken taste organs provide an ideal system for multidisciplinary studies, including organogenesis and regenerative medicine, considering their many features as outlined below: (1) a unique distribution pattern in the gustatory tissue of the oral cavity; (2) a much shorter lifespan (~4 days) compared to mammals (~10–12 days in rodents) indicating a more active progenitor cell niche and allowing for a more efficient way to study taste bud cell renewal compared to the rodent model; (3) similarity to humans in the connective tissue cell marker Vimentin being expressed in a large population of taste bud cells, suggesting a comparable mechanism underlying the contribution of connective tissue to taste buds in both organisms; (4) development of new taste buds after hatching which provides a time window to study the regulation of taste bud development; (5) other beneficial aspects of using chickens as a research model, e.g., convenience of in vivo embryo manipulation, high availability and rapid development.

Behavioral Responses of Chickens to Taste Stimuli

Chickens respond to taste stimuli immediately after hatch, and newly hatched chicks respond to different taste stimuli and show aversion/acceptance behavior for different tastants [23,57]. Behavioral studies have identified the typical response to tastants including shaking the head, wiping the beak, and tongue/beak movements. In addition, the ability of tastants to signal to the brain has been analyzed with electroencephalogram (EEG) [58].

Taste sensitivity of chickens

The sensitivity of taste in chickens positively correlates with the total number of taste buds, i.e., the more the taste buds, the more sensitive the bitter taste [10,59]. Broiler-type males are more sensitive to taste stimuli than layer-type males because they have more taste buds [8,9]. There has been a broad consensus that birds have a lower taste acuity compared to mammals due to their low taste bud numbers. However, the recent finding using molecular markers to label taste buds shows that birds have a well-developed taste system and a high number of taste buds relative to the volume of mouth cavity [11].

Indeed, chickens (from P0 to adult) respond to chemical stimulants (e.g., hydrochloric acid, acetic acid) even at the low concentrations (Gentle, 1972). Although taste sensitivity cannot be quantified by behavioral responses, there exists a correlation between oral response and taste sensitivity. Going forward, it will be important to establish an ideal method for evaluating the actual response to chemical stimuli.

Real-time Ca²⁺ imaging in isolated chicken taste buds has been used to investigate the responses of taste cells to bitter, salt and umami tastants, and has proved to be a functional approach for analyzing the taste senses of the chicken [60]. Moreover, a simple method developed recently for labeling chicken taste buds with molecular markers in the intact epithelial sheet of palate and base of oral cavity may provide an efficient way to determine an accurate number and overall distribution pattern of all chicken taste buds, and may facilitate studies correlating taste bud quantity with feeding behaviors in chickens [11].

Taste quality of chickens

It has been widely accepted that there are five basic taste qualities (sweet, bitter, umami, sour, salty). The different taste receptor molecules and ion channels that are localized in the cell membrane of different types of taste bud cells are the mediating molecules for transducing different taste stimuli. Therefore, the taste receptor and ion channel gene expressions in taste bud cells are responsible for taste qualities. For example, in mammals taste quality is determined by taste receptor and channel gene expressions in different taste cell types, e.g., sweet by T1R2+T1R3 in type II, umami by T1R1+T1R3 in type II, bitter by T2Rs in type II, salt by ENaC in type I, sour by PCKD channels in type III cells [61].

Compared to mammals, chickens have fewer genes for taste receptors, e.g., lacking the taste receptor T1R2 for sweet [62,63] and the bitter taste receptor repertoire is small, consisting of only 3 members (T2R1, T2R2 and T2R7), in contrast to humans (25), cows (11), and mice (35) [64,65]. Thus, in contrast to mammals, which have five taste qualities, it is presumed that chickens are only able to detect four tastes (sour, umami, salt and bitter).

Early behavioral studies have been performed in chickens using some commonly used tastants, including sucrose, saccharine, quinine acid, sodium chloride, acetic acid, and hydrochloric acid [57]. Newly hatched chicks were able to distinguish bitter and sour tastes, exhibiting aversive responses [23,57]. Bitter stimuli (e.g., quinine chloride) activates all three bitter receptors [66] for aversive responses [57,67]. Chickens' aversive responses to quinine chloride in a dose-dependent manner are similar to mammals. In contrast, chickens do not display significant responses towards 'sweet' stimuli presumably due to the absence of T1R2 receptor [23,57]. Finally, chickens' umami taste is detected by the GPCR T1R family receptors (T1R1 and T1R3) [60,68,69].

In addition to the tastants described above, there are studies which indicate that chickens may have other taste qualities. For example, chicks are responsive to water while there was no significant response when using egg fluid, suggesting that water alone is a strong stimulus for birds [70]. Moreover, a recent study has demonstrated that G-protein-coupled receptor-120 mediates the response of taste cells to oleic and linoleic acid, and is regarded as one of the functional fat taste receptors in chickens [71].

Oral responses in chickens are different for unique stimuli and some stimuli are not perceived until a certain high concentration (fructose, sucrose etc.) [67]. Thus far, there has not been a standard method available to determine the taste responses. Different outcomes have been reported from individual studies which could be attributed to inconsistencies between testing methods [57,67]. Therefore, establishing a standard for studying the oral responses when subjected to unique stimuli will be important for furthering our knowledge in this area.

Although chickens have unique taste bud cell types, it is unclear whether individual cell types express specific taste receptors that contribute to specific tastes. A recent report showed that Gustducin⁺ spindle-shaped cells isolated from taste buds respond to umami and bitter taste stimuli [60], similar to Gustducin⁺ cells in mammals [18]. Gustducin is a G-protein present in type II taste bud cells in mammals, which has been reported to be expressed in sweet and bitter sensing taste cells in mice, hamsters and rats [17,72,73]. However, α -Gustducin has also been identified in a large subset of chicken taste cells [20]. The expression of α -Gustducin in chicken taste bud cells suggests a signaling mechanism that is similar to mammals. Further studies on the differentiation of individual taste cell types and the expression of different taste receptor genes specific for unique taste qualities will be significant for deepening our understanding of how taste quality is determined in chickens.

Association of taste sensitivity and quality in chickens

In chickens, taste sensitivity is different for specific taste stimuli. For instance, chickens are more tolerant to 'sour' taste compared to mammals, but they are highly sensitive to 'bitter' taste, despite having a lower sub-type number of bitter taste receptors [74], and only two of the three sub-types responsible for bitter taste [75]. Chickens also respond strongly to umami taste stimuli that are composed of inosine-5' monophosphate and monopotassium L-glutamate, suggesting that the 'umami' taste is highly conserved from birds to mammals [69]. However, they are only responsive to sweet and salty taste stimuli at very high concentrations (i.e., sucrose, 0-5N) [23,57].

Taste sensitivity for specific taste qualities may be altered under certain conditions. Zinc-deficiency in chickens affects water intake and enhances responses to bitter and salt taste stimuli, in stark contrast to reports in humans and rats that zinc deficiency leads to taste loss [76,77]. Although no morphological changes in taste buds were associated with these deficiencies, feed supplemented with these minerals caused a significant increase in responses to taste stimuli [78]. Vitamin A deficiency in chickens is also reported to cause a decrease in response to taste stimuli [79], presumably because Vitamin A is important for the maintenance of integrity of the epithelium. Similarly, Vitamin A deficiency leads to a significant decrease in response to NaCl and quinine chloride stimuli in rats [80].

In summary, chickens are sensitive to taste stimuli, which is consistent to the well-developed taste system indicated by the recent studies and indicates a higher than previously thought impact on feeding behaviors. The response of birds (including chickens) to each group of taste stimuli for primary taste qualities are reviewed by Roura et al. [81]. Further behavioral studies are needed for the practical applications of taste perceptions on dietary intake in poultry industry.

Impact of Understanding Chicken's Taste Sensation on Poultry Industry

Chickens are one of the major livestock animals used for egg production and meat. Taste buds are the sensory organs that guide nutritive choices and motivate feed intake, and thus have a direct impact on the productivity of these animals. Thorough knowledge regarding taste bud development, regulation and taste response to different stimuli helps to improve the feed efficiency, thereby increasing the productivity. Further mechanistic studies on how taste bud development and taste receptor gene expression are regulated will provide information beneficial for improving feed intake and animal performance.

Acknowledgements

This study was supported by the National Institutes of Health (grant number R01 DC012308 to HXL) and University of Georgia Start-up fund to HXL.

References

- Doty RL (2015) Handbook of olfaction and gustation (3rd Edn). Wiley Blackwell, Hoboken, New Jersey, p: 1240.
- Höfer D, Püschel B, Drenckhahn D (1996) Taste receptor-like cells in the rat gut identified by expression of alpha-gustducin. Proc Natl Acad Sci USA 93: 6631-6634.
- Lalonde ER, Eglitis JA (1961) Number and distribution of taste buds on the epiglottis, pharynx, larynx, soft palate and uvula in a human newborn. Anat Rec 140: 91-95.
- Travers SP, Nicklas K (1990) Taste bud distribution in the rat pharynx and larynx. Anat Rec 227: 373-379.
- Lindenmaier P, Kare MR (1959) The taste end-organs of the chicken. Poult Sci J 38: 545-550.
- Berkhoudt H (1985) Special sense organs: structure and function of avian taste receptors. Form and function in birds 3: 463-496.
- Davies RO, Kare MR, Cagan RH (1979) Distribution of taste buds on fungiform and circumvallate papillae of bovine tongue. Anat Rec 195: 443-446.
- Ganchrow D, Ganchrow JR (1985) Number and distribution of taste buds in the oral cavity of hatchling chicks. Physiol Behav 34: 889-894.
- Kudo K, Nishimura S, Tabata S (2008) Distribution of taste buds in layer-type chickens: scanning electron microscopic observations. Anim Sci J 79: 680-685.
- Kudo K, Shiraishi J, Nishimura S, Bungo T, Tabata S (2010) The number of taste buds is related to bitter taste sensitivity in layer and broiler chickens. Anim Sci J 81: 240-244.
- Rajapaksha P, Wang Z, Venkatesan N, Tehrani KF, Payne J, et al. (2016) Labeling and analysis of chicken taste buds using molecular markers in oral epithelial sheets. Sci Rep 6: 37247.
- Ganchrow JR, Ganchrow D (1987) Taste bud development in chickens (*Gallus gallus domesticus*). Anat Rec 218: 88-93.
- Kurosawa T, Kusuhara S, Ishida K, Niimura S (1983) Morphological studies of taste buds in chickens. Jap J Zootech Sci.
- Ganchrow D, Ganchrow JR, Goldstein RS (1991) Ultrastructure of palatal taste buds in the perihatching chick. Am J Anat 192: 69-78.
- Ganchrow D, Ganchrow JR, Royer SM, Dovidpor S, Kinnamon JC (1998) Identified taste bud cell proliferation in the perihatching chick. Chem Senses 23: 333-341.
- Bartel DL, Sullivan SL, Lavoie ÉG, Sévigny J, Finger TE (2006) Nucleoside triphosphate diphosphohydrolase-2 is the ecto-ATPase of type I cells in taste buds. J Comp Neurol 497: 1-12.
- Boughter Jr JD, Pumplin DW, Yu C, Christy RC, Smith DV (1997) Differential expression of α -gustducin in taste bud populations of the rat and hamster. J Neurosci 17: 2852-2858.
- Yang R, Tabata S, Crowley HH, Margolskee RF, Kinnamon JC (2000) Ultrastructural localization of gustducin immunoreactivity in microvilli of type II taste cells in the rat. J Comp Neurol 425: 139-151.
- Yang R, Crowley HH, Rock ME, Kinnamon JC (2000) Taste cells with synapses in rat circumvallate papillae display SNAP25 like immunoreactivity. J Comp Neurol 424: 205-215.
- Kudo K, Wakamatsu K, Nishimura S, Tabata S (2010) Gustducin is expressed in the taste buds of the chicken. Anim Sci J 81: 666-672.
- Venkatesan N, Rajapaksha P, Payne J, Goodfellow F, Wang Z, et al. (2016) Distribution of α -gustducin and vimentin in premature and mature taste buds in chickens. Biochem Biophys Res Commun 479: 305-311.
- Witt M, Reutter K, Ganchrow D, Ganchrow JR (2000) Fingerprinting taste buds: intermediate filaments and their implication for taste bud formation. Philos Trans R Soc Lond B Biol Sci 355: 1233-1237.
- Ganchrow JR, Steiner JE, Bartana A (1990) Behavioral reactions to gustatory stimuli in young chicks (*Gallus gallus domesticus*). Dev Psychobiol 23: 103-117.

24. Hosley MA, Oakley B (1987) Postnatal development of the vallate papilla and taste buds in rats. *Anat Rec* 218: 216-222.
25. Shin Y, Cong W, Cai H, Kim W, Maudsley S, et al. (2011) Age-related changes in mouse taste bud morphology, hormone expression, and taste responsiveness. *J Gerontol A Biol Sci Med Sci* 67A: 336-344.
26. Ganchrow D, Ganchrow JR, Romano R, Kinnamon JC (1994) Ontogenesis and taste bud cell turnover in the chicken. I. Gemmal cell renewal in the hatchling. *J Comp Neurol* 345: 105-114.
27. Ganchrow JR, Ganchrow D, Royer SM, Kinnamon JC (1993) Aspects of vertebrate gustatory phylogeny: morphology and turnover of chick taste bud cells. *Microsc Res Tech* 26: 106-119.
28. Beidler LM, Smallman RL (1965) Renewal of cells within taste buds. *J Cell Biol* 27: 263-272.
29. Perea-Martinez I, Nagai T, Chaudhari N (2013) Functional cell types in taste buds have distinct longevities. *PLoS ONE* 8: e53399.
30. Sullivan JM, Borecki AA, Oleskevich S (2010) Stem and progenitor cell compartments within adult mouse taste buds. *Eur J Neurosci* 31: 1549-1560.
31. Boggs K, Venkatesan N, Mederacke I, Komatsu Y, Stice S, et al. (2016) Contribution of underlying connective tissue cells to taste buds in mouse tongue and soft palate. *PLoS ONE* 11: e0146475.
32. Huang R, Lang ER, Otto WR, Christ B, Patel K (2001) Molecular and cellular analysis of embryonic avian tongue development. *Anat Embryol* 204:179-187.
33. Ermilov AN, Kumari A, Li L, Joiner AM, Grachtchouk MA, et al. (2016) Maintenance of taste organs is strictly dependent on epithelial hedgehog/GLI signaling. *PLoS Genet* 12: e1006442.
34. Hall JM, Hooper JE, Finger TE (1999) Expression of sonic hedgehog, patched, and gli1 in developing taste papillae of the mouse. *J Comp Neurol* 406: 143-155.
35. Kim JY, Lee MJ, Cho KW, Lee JM, Kim YJ, et al. (2009) Shh and ROCK1 modulate the dynamic epithelial morphogenesis in circumvallate papilla development. *Dev Biol* 325: 273-280.
36. Liu HX, Maccallum DK, Edwards C, Gaffield W, Mistretta CM (2004) Sonic hedgehog exerts distinct, stage-specific effects on tongue and taste papilla development. *Dev Biol* 276: 280-300.
37. Liu HX, Ermilov A, Grachtchouk M, Li L, Gumucio DL, et al. (2013) Multiple Shh signaling centers participate in fungiform papilla and taste bud formation and maintenance. *Dev Biol* 382: 82-97.
38. Mistretta CM, Liu HX, Gaffield W, MacCallum DK (2003) Cyclopamine and jervine in embryonic rat tongue cultures demonstrate a role for Shh signaling in taste papilla development and patterning: fungiform papillae double in number and form in novel locations in dorsal lingual epithelium. *Dev Biol* 254: 1-18.
39. Gaillard D, Barlow LA (2011) Taste bud cells of adult mice are responsive to Wnt/ β -catenin signaling: implications for the renewal of mature taste cells. *Genesis* 49: 295-306.
40. Gaillard D, Bowles SG, Salcedo E, Xu M, Millar SE, et al. (2017) β -catenin is required for taste bud cell renewal and behavioral taste perception in adult mice. *PLoS Genet* 13: e1006990.
41. Iwatsuki K, Liu H, Grönder A, Singer MA, Lane TF, et al. (2007) Wnt signaling interacts with Shh to regulate taste papilla development. *Proc Natl Acad Sci USA* 104: 2253-2258.
42. Liu F, Millar SE (2010) Wnt/ β -catenin signaling in oral tissue development and disease. *J Dent Res* 89: 318-330.
43. Liu F, Thirumangalathu S, Gallant NM, Yang SH, Stoick-Cooper CL, et al. (2007) Wnt- β -catenin signaling initiates taste papilla development. *Nat Genet* 39: 106-112.
44. Beites CL, Hollenbeck PLW, Kim J, Lovell-Badge R, Lander AD, et al. (2009) Follistatin modulates a BMP autoregulatory loop to control the size and patterning of sensory domains in the developing tongue. *Development* 136: 2187-2197.
45. Han D, Zhao H, Parada C, Hacia JG, Bringas P, et al. (2012) A TGF β -Smad4-Fgf6 signaling cascade controls myogenic differentiation and myoblast fusion during tongue development. *Development* 139: 1640-1650.
46. Kawasaki K, Pomtaveetus T, Oommen S, Ghafoor S, Kawasaki M, et al. (2012) Bmp signalling in filiform tongue papillae development. *Arch Oral Biol* 57:805-813.
47. Zhou Y, Liu H, Mistretta CM (2006) Bone morphogenetic proteins and noggin: inhibiting and inducing fungiform taste papilla development. *Dev Biol* 297: 198-213.
48. Ren W, Aihara E, Lei W, Gheewala N, Uchiyama H, et al. (2017) Transcriptome analyses of taste organoids reveal multiple pathways involved in taste cell generation. *Sci Rep* 7: 4004.
49. Seta Y, Seta C, Barlow LA (2003) Notch-associated gene expression in embryonic and adult taste papillae and taste buds suggests a role in taste cell lineage decisions. *J Comp Neurol* 464: 49-61.
50. Seta Y, Toyono T, Kataoka S, Toyoshima K (2005) Regulation of taste bud cell differentiation by notch signaling pathway. *Chem Senses* 30: i48-i49.
51. Ookura T, Kawamoto K, Tsuzaki H, Mikami Y, Ito Y, et al. (2002) Fibroblast and epidermal growth factors modulate proliferation and neural cell adhesion molecule expression in epithelial cells derived from the adult mouse tongue. *In Vitro Cell Dev Biol Anim* 38: 365-372.
52. Petersen CI, Jheon AH, Mostowfi P, Charles C, Ching S, et al. (2011) FGF signaling regulates the number of posterior taste papillae by controlling progenitor field size. *PLoS Genet* 7: e1002098.
53. Liu H, Henson BS, Zhou Y, D'Silva NJ, Mistretta CM (2008) Fungiform papilla pattern: EGF regulates inter-papilla lingual epithelium and decreases papilla number by means of PI3K/Akt, MEK/ERK, and p38 MAPK signaling. *Dev Dyn* 237: 2378-2393.
54. McLaughlin SK (2000) Erb and c-Kit receptors have distinctive patterns of expression in adult and developing taste papillae and taste buds. *J Neurosci* 20: 5679-5688.
55. Morris-Wiman J, Segó R, Brinkley L, Dolce C (2000) The effects of sialoadenectomy and exogenous EGF on taste bud morphology and maintenance. *Chem Senses* 25: 9-19.
56. Cui X, Marshall B, Shi N, Chen SY, Rekaya R, et al. (2017) RNA-Seq analysis on chicken taste sensory organs: an ideal system to study organogenesis. *Sci Rep* 7: 9131.
57. Gentle MJ (1972) Taste preference in the chicken (*Gallus domesticus* L). *Br Poult Sci* 13: 141-155.
58. Halpern BP (1962) Gustatory nerve responses in the chicken. *Am J Physiol* 203: 541-544.
59. Miller Jr IJ, Reedy Jr FE (1990) Variations in human taste bud density and taste intensity perception. *Physiol Behav* 47: 1213-1219.
60. Kudo K, Kawabata F, Nomura T, Aridome A, Nishimura S, et al. (2014) Isolation of chicken taste buds for real-time Ca²⁺ imaging. *Anim Sci J* 85: 904-909.
61. Chandrashekar J, Hoon MA, Ryba NJ, Zuker CS (2006) The receptors and cells for mammalian taste. *Nature* 444: 288-294.
62. Cheled-Shoval SL, Druyan S, Uni Z (2015) Bitter, sweet and umami taste receptors and downstream signaling effectors: expression in embryonic and growing chicken gastrointestinal tract. *Poult Sci J* 94: 1928-1941.
63. Lagerström MC, Hellström AR, Gloriam DE, Larsson TP, Schiöth HB, et al. (2006) The G protein-coupled receptor subset of the chicken genome. *PLoS Comput Biol* 2: e54.
64. Go Y (2006) Lineage-specific expansions and contractions of the bitter taste receptor gene repertoire in vertebrates. *Mol Biol Evol* 23: 964-972.
65. Meyerhof W (2005) Elucidation of mammalian bitter taste. *Rev Physiol Biochem pharmacol* 154: 37-72.
66. Cheled-Shoval SL, Behrens M, Meyerhof W, Niv MY, Uni Z (2014) Perinatal administration of a bitter tastant influences gene expression in chicken palate and duodenum. *J Agric Food Chem* 62: 12512-12520.
67. Gentle MJ, Harkin C (1979) The effect of sweet stimuli on oral behaviour in the chicken. *Chem Senses* 4: 183-190.
68. Cheled-Shoval SL, Reicher N, Niv MY, Uni Z (2017) Detecting thresholds for bitter, umami, and sweet tastants in broiler chicken using a 2-choice test method. *Poult Sci J* 96: 2206-2218.
69. Yoshida Y, Kawabata Y, Kawabata F, Nishimura S, Tabata S (2015) Expressions of multiple umami taste receptors in oral and gastrointestinal tissues, and umami taste synergism in chickens. *Biochem Biophys Res Commun* 466: 346-349.
70. Vince MA (1977) Taste sensitivity in the embryo of the domestic fowl. *Anim Behav* 25: 797-805.

71. Sawamura R, Kawabata Y, Kawabata F, Nishimura S, Tabata S (2015) The role of G-protein-coupled receptor 120 in fatty acids sensing in chicken oral tissues. *Biochem Biophys Res Commun* 458: 387-391.
72. Cho YK, Farbman AI, Smith DV (1998) The timing of α -gustducin expression during cell renewal in rat vallate taste buds. *Chem Senses* 23: 735-742.
73. McLaughlin SK, McKinnon PJ, Margolskee R (1992) Gustducin is a taste-cell-specific G protein closely related to the transducins. *Nature* 357: 563-569.
74. Hirose N, Kawabata Y, Kawabata F, Nishimura S, Tabata S (2015) Bitter taste receptor T2R1 activities were compatible with behavioral sensitivity to bitterness in chickens. *Biochem Biophys Res Commun* 460: 464-468.
75. Dey B, Kawabata F, Kawabata Y, Yoshida Y, Nishimura S, et al. (2017) Identification of functional bitter taste receptors and their antagonist in chickens. *Biochem Biophys Res Commun* 482: 693-699.
76. Catalanotto FA, Nanda R (1977) The effects of feeding a zinc-deficient diet on taste acuity and tongue epithelium in rats. *J Oral Pathol Med* 6: 211-220.
77. Henkin RI (1984) Zinc in taste function: a critical review. *Biol Trace Elem Res* 6: 263-280.
78. Gentle M, Dewar W, Wight P (1981) The effects of zinc deficiency on oral behaviour and taste bud morphology in chicks. *Br Poult Sci* 22: 265-273.
79. Gentle M, Dewar W (1981) The effects of vitamin A deficiency on oral gustatory behaviour in chicks. *Br Poult Sci* 22: 275-279.
80. Bernard RA, Halpern BP, Kare MR (1961) Effect of vitamin A deficiency on taste. *Exp Biol Med* 108: 784-786.
81. Roura E, Baldwin MW, Klasing KC (2013) The avian taste system: potential implications in poultry nutrition. *J Ani Feed Sci Tech* 108: 1-9.