

Assessment of Trace Elements in Bissap (*Hibiscus sabdariffa*) Sold in the City of Abidjan in Côte d'Ivoire

M'boh Gervais Melaine^{1*}, N'guessan Ismaël Gbabi Zié², Ackah Jacques Auguste Alfred Bognan², Konan Kouassi Martin³ and Djaman Allico Joseph^{1,3}

¹Institut Pasteur de Côte d'Ivoire, Ivory Coast

²University of Lorougnon Guédé, Ivory Coast

³University Félix Houphouët Boigny, Ivory Coast

Abstract

Pollutants such as lead, cadmium and mercury are harmful to living things at low concentrations. In Côte d'Ivoire, the drink of *Hibiscus sabdariffa* Linn generally called "Bissap juice" is widely consumed by the population. However, recent studies have revealed a significant presence of heavy metals in the cultivated soils as well as in the plants taken from these soils in the city of Abidjan. Our study aimed to assess the health risks associated with the consumption of Bissap marketed in three (3) communes of Abidjan. Thus, evaluation of trace metallic elements (Pb, Cd, Cu, Zn) in *Hibiscus sabdariffa* L. calyces given as average value: Cd ($23,142 \pm 13,332$ mg/kg), Pb ($77,949 \pm 15,620$ mg/kg) Zn (78.96 ± 15.584 mg/kg) and Cu (42.894 ± 28.394 mg/kg). Bissap juice showed average concentrations Zn (0.139 ± 0.076 mg/L), Pb (0.544 ± 0.143 mg/L) and Cd (0.221 ± 0.064 mg/L). These data compared to Codex Alimentarius and European Commission standards revealed a significant risk of lead and cadmium poisoning in calyces and Bissap juice and a lower value for metal copper compared to zinc. Our study requires further investigation in order to evaluate all the factors involved in the chain of contamination and to assess possibly the toxicological and pathophysiological risks incurred by the population due to abuse.

Keywords: *Hibiscus sabdariffa*; Trace elements; Bissap; Chronic poisoning

Introduction

Awareness of the dangers of metal trace elements (ETM) is unequivocal for scientists and leaders around the world. The issues raised by the ETM or heavy metals are mainly environmental but above all health [1]. Trace metals (Copper, Zinc, Iron, Arsenic, etc.) are all toxic or toxic at a certain threshold [2]. Some of these pollutants such as lead, cadmium and mercury are harmful to living things at low concentrations [3]. One of the main sources of human intoxication to heavy metals is diet.

The use of chemical fertilizers, food additives, environmental pollutants and others related, promote the presence of heavy metals (HMT) in plants, water, fish and other products used for human consumption [4]. HMTs are a real public health problem with metabolic disorders and chronic diseases (cancer, high blood pressure, neurological diseases, infertility, etc.) and the high costs of managing these diseases [5].

Zinc (Zn) is essential for normal immune function [6] and has been shown to reduce the incidence of diarrhea and pneumonia [7] but also involved with Cu as cofactors of superoxide dismutase enzyme to fight against oxidative stress. The need for Cu also derives from its involvement in a myriad of biological processes, including antioxidant defense, neuropeptide synthesis and immune function [8,9]. Cu deficiency may result in impaired development of the cardiovascular system, bone malformation and ongoing neurologic and immunologic abnormalities into infancy and beyond [10,11].

In terms of health, the level of morbidity and mortality remains high in Côte d'Ivoire, mainly affecting women and children. General morbidity remains characterized by communicable diseases (Tuberculosis) and by non-communicable diseases (NCDs) in a health context of concern [12]. According to the WHO, NCDs show a marked increase and are responsible for 31% of deaths in Côte d'Ivoire in

recent years [13]. Thus, no communicable diseases are a national health concern and remain dominated by metabolic diseases, cardiovascular diseases and cancers [14].

Hibiscus sabdariffa Linnaeus var. *Sabdariffa*, known in Côte d'Ivoire under its vernacular name of Bissap, Dah or Dà foura [15], is sold in almost all public spaces, making Ivorian eating habits (feast, accompaniment of snacks, refreshing-toning, etc.).

It's very affordable cost makes it one of the street foods, the most consumed by the population especially by the younger who are fond of sweets. The vegetable growing of Bissap (*Hibiscus sabdariffa* L.) occurs in the urban and peri-urban areas of the cities of Abidjan and Yamoussoukro [16]. However, in recent years, several studies have highlighted a worrying presence of heavy metals in these vegetable soils in Abidjan and in the crops grown there [17-19]. Moreover, the presence of these contaminants was also noted in the plants sold on the Abidjan markets [20]. The present study generally aims to determine the levels of four (4) trace metals, namely cadmium (Cd), copper (Cu), lead (Pb) and zinc (Zn), in *Hibiscus sabdariffa* L. (Dried calyces and Bissap juice) sold in the city of Abidjan. It is part of a contribution to the assessment of the level of exposure of populations to MTEs and therefore to the reduction of the risks of proliferation of metabolic and chronic diseases.

***Corresponding author:** M'boh Gervais Melaine, Institut Pasteur de Côte d'Ivoire, Ivory Coast, Tel: 07357030; E-mail: gervaismbh@pasteur.ci

Received August 22, 2017; **Accepted** September 07, 2017; **Published** September 13, 2017

Citation: M'boh GM, N'Guessan IGZ, Ackah JAAB, Konan KM, Djaman AJ (2017) Assessment of Trace Elements in Bissap (*Hibiscus sabdariffa*) Sold in the City of Abidjan in Côte d'Ivoire. J Plant Biochem Physiol 5: 197. doi: [10.4172/2329-9029.1000197](https://doi.org/10.4172/2329-9029.1000197)

Copyright: © 2017 M'boh GM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Materials and Methods

Study area

Abobo, Yopougon and Adjamé were the three communes of the District of Abidjan (Côte d'Ivoire) which served as a framework for our research. These communes are distinguished by their demographics and their fluctuations (human, commercial, etc.) Abobo covers an area of about 9,000 hectares, with a population estimated at 1,030,658 inhabitants. According to the latest census, it is the second most populous commune in Abidjan (and Côte d'Ivoire) after Yopougon [21]; first Ivorian commune by its demography with a population estimated at 1,071,543 habitants. Unlike the two previous municipalities, Adjamé has an estimated population of 372,978 habitants and constitutes only the 5th most populated municipality [21]. However, this commune is distinguished by the external affluence it records every day. Indeed, Adjamé represents a very important trade and human transit hub for the District of Abidjan.

Sampling

The biological material was composed of dry calyces of *Hibiscus sabdariffa* Linn. and Bissap beverages or "Bissap juice".

Our study sites were selected on the basis of the demographic factor [21] and previous studies carried out [20]. In the three communes, the calyces were bought in the main supply markets, while beverages came from markets as well as other busy public spaces (bus stations, crossroads, sidewalks, etc.). Only traders with a high daily customer base were included in this study. The goal is to have an excellent representativeness of the consumer population of Bissap.

Samples kept in their original containers were transported in coolers chilled by carboglaces; Calyces were packed in hermetically sealed plastic bags. Each sample was labeled to facilitate future identification.

Samples preparation

Concerning the calyces, it consisted in passing the dry flowers to the machine for a crude fragmentation. This step is essential to optimize the extraction of Bissap juice; traders ensured this phase of fragmentation for their customers.

Concerning the drink, the dry calyces of *Hibiscus sabdariffa* were then soaked in water at room temperature for an average of 1 to 2 days or boiled at least 100°C in water between thirty (30) minutes and one (1) hour. Then, after filtration, sugar was added for the formulation of the final beverage. However, other ingredients (fruits, mint leaves, chemical flavors, etc.) could be added.

Determination of trace elements

The determination of the metallic trace elements (ETM) was carried out by atomic absorption spectrophotometry with an air-acetylene flame (Varian AA20 Pattern®, France). The different concentrations were read in triplet by ranges of standard elements of copper, zinc, cadmium and lead at the following wavelengths: 229.6 nm for Cd, 217.8 nm for Pb, 325.5 nm for Cu and 214.6 nm for Zn. Concerning calyces, a mass of 0.3 g of dried calyces milled and placed in a porcelain crucible was placed in an oven at 600°C for 5 hours. After cooling, 5 mL of nitric acid (1N) were added to the ash obtained and then brought to total evaporation on a sand bath previously heated. To the residue are added 5 ml of hydrochloric acid (0.1N). The solution was then re-baked at 400 °C. for 30 min. The final residue was recovered in 10 ml hydrochloric acid (0.1N) and then transferred to a 50-ml flask. The procedure was repeated three (3) times and then the contents of the completed flask were diluted with hydrochloric acid to 50 ml. The trace elements contained in the solution were then assessed after blank tests [22,23]. Concerning the Drink (Bissap juice), a volume of 6 mL of each sample was taken and then placed in dry tubes before assaying the ETMs contained in the solution.

Statistical analysis

The GraphPad Prism 7 software was used as a means of statistical data processing. The ANOVA test and multiple comparison (Tukey) of the variation of means were applied to the data. The differences were considered significant for $p < 0.05$ to $\alpha = 0.05$.

Results

The majority of calyce sold in the market was found in Adjamé market and the Bissap juice performed in majority following a hot extraction (Figure 1).

Distribution of trace metal in Bissap calyces

The results showed the presence of cadmium, lead, zinc and copper at varying concentrations in the calyces of *Hibiscus sabdariffa*. Bissap calyces sold in the city of Abidjan (Figure 2) had relatively high Cd contents (23.14 ± 13.33 mg/kg), Pb (77.95 ± 15.62 mg/kg), Zn (78.96 ± 15.58 mg/kg) and Cu (42.90 ± 28.39 mg/kg) with substantially equal lead and zinc.

The trace metal content (Table 1) was lower in Adjame than in the other two communes. Yopougon calyces were richer in Zn and Cu, while Abobo's calyces were richer in Cd and Pb. However, average concentrations between study areas showed significant differences ($P < 0.001$), except Cd in the calyces at Adjamé and Yopougon with ($P = 0.144$).







Figure 1: Distribution of calyce in the market (A); Preparation about the juice extraction (B).



Distribution of trace metal in Bissap juice

The presence of trace copper was found in the beverages of Hibiscus Sabdariffa. Zinc (0.14 ± 0.07 mg/L) was present in Bissap juices at very low levels compared to toxic metals (Cadmium and Lead). However, Pb (0.54 ± 0.14 mg/L) was twice as high as Cd (0.22 ± 0.06 mg/L) in Bissap juice sold in Abidjan (Figures 3 and 4).

Mean Pb and Zn levels showed significant differences between the three study areas. On the other hand, those of the Cd were statistically identical except between Abobo and Adjame ($P=0.09$). Bissap dosed

Area market	Trace elements (mg/Kg)			
	Cadmium	Lead	Zinc	Copper
Abobo	$40.64 \pm 2.34^*$	$93.26 \pm 3.04^*$	75.74 ± 4.03	32.24 ± 2.82
Adjame	12.33 ± 1.42	58.34 ± 2.45	63.13 ± 3.95	16.83 ± 1.09
Yopougon	16.45 ± 1.24	82.25 ± 2.12	$98.01 \pm 1.93^*$	$79.60 \pm 2.42^*$

* $p < 0.001$

Table 1: Repartition of trace metallic elements in calyces according to area market.

Area market	Trace elements (mg/L)			
	Cadmium	Lead	Zinc	Copper
Abobo	0.16 ± 0.03	$0.72 \pm 0.03^*$	0.14 ± 0.04	ND
Adjame	0.21 ± 0.03	0.50 ± 0.04	$0.22 \pm 0.03^*$	ND
Yopougon	$0.29 \pm 0.05^*$	0.41 ± 0.03	0.06 ± 0.02	ND

* $p < 0.001$

Table 2: Repartition of trace metallic elements in the Bissap juice according to area market.

juice showed higher levels of Cd to Yopougon, Pb to Abobo and Zinc to Adjame (Table 2).

Discussion

In this study, the concentrations of metals (Cd, Pb, Zn and Cu) recorded in the dry calyces of Bissap were compared with those observed by other researchers. Thus, in their work on the synthesis of Hibiscus sabdariffa, [24] reported concentrations of 6 mg/kg of Cu; 1.8 mg/kg in Pb and up to 65 mg/kg in Zn in the dry calyces of Bissap from Mali and Nigeria. These Cu and Pb data are much lower than those obtained in the calyces sold on the markets of Abidjan (Côte d'Ivoire) but identical for the Zn. However, the Zn levels (814 mg/Kg) presented by [25] for purple red calyces were much larger than ours. Hibiscus sabdariffa can be used for soil phytoremediation [26]. The presence of trace metals in calyces could be explained by bioaccumulation resulting from soil contamination in crop soils [27], but also by exposure to environmental dust, Engine exhaust etc. [20].

Adjame was the main calyx supply area and the least populated; this may justify the lower grades recorded in this study area. Thus, the levels of TME in the calyces of Adjame would be an indicator to better understand the level of initial contamination and by extrapolation the level of exposure to metals for each of the study areas. However, by referring to the maximum permitted Cd (0.2 mg/kg) and Pb (0.3 mg/kg) limits for edible plants by [28]; in Zn (20 mg/kg) and Cu (100 mg/kg) reported in the publication by [17], there would be a high risk of Cd, Pb and Zn contamination for consumers in various forms of dried calyces from Abobo, Adjame and Yopougon. On the other hand, the Cu content had been found to be below the recommended limit value for this metal. In the study by [29], the testicular cytotoxicity of rats with water extracts of Hibiscus sabdariffa could be justified by the presence of lead [30] in the Bissap calyces. The presence of Cd, Pb, Cu and Zn in the Bissap juice sold in Abidjan would be likely due to the presumed existence of these metals in the calyces of Hibiscus sabdariffa. However, the copper contents were below the detection limit and could not be evaluated.

We have statistically reconciled our data to those of [31] who assessed the Cu and Zn concentrations of two types of Hibiscus sabdariffa beverage marketed in Nigeria: "Sorrel cordial juice" (Cu: 0.056 mg; Zn: 1.46 mg) and the "Black currant cordial juice" (Cu: 0.008 mg; Zn: 0.70 mg). These data as well as those reported by [32] (Zn: 814 mg and Cu: 24.4 mg), were higher compared to our study (Zn: 0.03-0.25 mg; Cu in trace). However, Pb and Zn levels in drinks consumed

in Nigeria based on *Hibiscus sabdariffa*, showed some results identical to ours [33]. Moreover, when we referred to Codex Alimentarius Codex Stan 179-1991 for fruit and vegetable juices [34] with maximum limits for Pb at 0.1 mg/kg, for Zn at 5 mg/kg and Cu at 5 mg/kg, there was a potential for lead poisoning [0.37-0.75 mg/l] for Bissap juice consumers; this risk would be enhanced by the presence of Cadmium [0.12-0.34 mg/L]. Lead and cadmium are among the top ten (10) most toxic to humans [35].

The low value of Zn and Cu in bissap juice could be justified by the long duration related to the hot cooking of this drink used in majority by the population. The lack of Cu observed in the bissap juice would lead to a bad choice of this drink as a dietary Cu intake; being given that Cu with Zinc are both cofactor of SOD enzyme, important to fight against the oxydative stress in the organism. Other factors could also be considered in assessing the potential hazards to the population: Firstly, the low presence of Zinc, which plays an important role in protecting the organism against cadmium [36,37]; but also in many other biological phenomena such as cell growth and proliferation [38], the fight against free radicals [39], immune responses [40], reproduction [41]. Then, we could also highlight the method of hot extraction which would considerably reduce the anthocyanin content knowing their important role in the antioxidant activity of Bissap [42]. Finally, the ignorance of the actors of the sector of Bissap and the bad manufacturing practices would favor the microbiological risks [43,44].

Prolonged and regular consumption of Bissap which contains Pb and Cd could lead to chronic heavy metal poisoning with disastrous consequences for the health of populations [45]. Works demonstrated the cumulative effect Pb and Cd on rats [46]. This chronic poisoning would have led to an increase in oxidative stress in the heart and liver as well as an alteration of these organs. Researchers have also demonstrated in rats that chronic co-exposure to lead and cadmium, a groundwater pollutant in a city in Algeria, would increase the risk of oxidative stress among the population [47]. Thus, regular and prolonged consumption of Bissap juice could be the cause of chronic diseases [48] such as high blood pressure, renal insufficiency, cancer, neurological disorders, sexual sterility etc.

Bissap juice is a drink of interest for its high consumption but also for its richness in anthocyanin which gives it many therapeutic properties. However, its poverty in essential minerals such as copper and zinc and the inappropriate presence of lead and cadmium beyond accepted recommendations do not militate to its advantage. Thus, given the impact of these heavy metals on the body, regular consumption of this drink by the population does not exclude the risks of prevalence of chronic diseases such as high blood pressure, renal insufficiency, cancer, neurological disorder etc. Our study requires further investigation in order to evaluate all the factors involved in the chain of contamination and to assess possibly the toxicological and pathophysiological risks incurred by the population due to abuse.

Contributors

MGM, AJAAB and DAJ conceived and designed the experiments. MGM and NIGZ performed the experiments. MGM, AJAAB, KKM analyzed the data. MGM and DAJ wrote the paper.

Conflict of Interest

The authors declare that there is no conflict of interest.

Acknowledgements

We thank the National Polytechnic Institute Félix Houphouët-Boigny (INPHB) for their support, and the population for their cooperation and contributions in the collected of samples.

References

- Papp R (2001) The environmental properties of trace elements. National College of Chemical Industry Environment Experts, CNEEIC, Paris, p: 112.
- Atli G, Canli M (2011) Essential metal (Cu, Zn) exposures alter the activity of ATPases in gill, kidney and muscle of tilapia *Oreochromis niloticus*. *Ecotoxicology* 20: 1861-1869.
- Waseem A, Arshad J, Iqbal F, Sajjad A, Mehmood Z, et al. (2014) Pollution status of Pakistan: a retrospective review on heavy metal contamination of water, soil, and vegetables. *BioMed Research International*, pp: 410-417.
- Gauthier C (2013) The impact of urban pollution on urban food production. Final report of the research carried out on behalf of the Brussels Institute for Environmental Management, Green Loop, p: 35.
- Jaffiol C (2016) Chronic Diseases: The New Challenge for Health in the 21st Century. Academy of Sciences and Letters of Montpellier (France), pp: 241-252.
- Shankar AH, Prasad AS (1998) Zinc and immune function: the biological basis of altered resistance to infection. *Am J Clin Nutr* 68: 447S-463S.
- Black RE (1998) Therapeutic and preventive effects of zinc on serious childhood infectious diseases in developing countries. *Am J Clin Nutr* 68: 476S-479S.
- Bonham M, O'Connor JM, Hannigan BM, Strain JJ (2002) The immune system as a physiological indicator of marginal copper status? *Br J Nutr* 87: 393-403.
- Uriu-Adams JY, Keen CL (2005) Copper, oxidative stress, and human health. *Mol Aspects Med* 26: 268-298.
- Cerhan JR, Saag KG, Merlino LA, Mikuls TR, Criswell LA (2003) Antioxidant micronutrients and risk of rheumatoid arthritis in a cohort of older women. *Am J Epidemiol* 157: 345-354.
- Georgieff MK (2007) Nutrition and the developing brain: nutrient priorities and measurement. *Am J Clin Nutr* 85: 614S-620S.
- WHO (2016) Strategies for cooperation with Côte d'Ivoire. World Health Organization WHO/CCU/16.03/Cote d'Ivoire, p: 2.
- WHO (2014) Country Profiles for Non-Communicable Diseases (NCDs) - Côte d'Ivoire. World Health Organization, p: 1.
- Ministry of Health and Public Hygiene (2016) National policy to improve the quality of health care and services in Côte d'Ivoire. Abidjan, p: 68.
- Roberty G (1954) Small flora of West Africa. French Republic Ministry of Overseas France, p: 441.
- FAO (2009) National report on the state of plant genetic resources for food and agriculture. Republic of Côte d'Ivoire, National Center for Agronomic Research, Abidjan, Côte d'Ivoire, Food and Agriculture Organization of the United Nations, p: 152.
- Kouakou KJ, Sika AE, Gogbeu SJ, Yao KB, Bounakhla M, et al. (2015) Level of exposure to metallic trace elements (Cd, Cu, Pb, Zn, Ni) of amaranth (*Amaranthus paniculatus* L.) and lettuce (*Lactuca sativa* L.) grown on marshy sites in the city of Abidjan (Abidjan, Ivory Coast). *International Journal of Innovation and Applied Studies* 10: 21-29.
- Kouassi JK, Ahoua ES, Denezon OO, Bekro Y, Baize D, et al. (2012) Levels of trace metals in market gardening soils in the city of Abidjan (Côte d'Ivoire). *International Journal of Biological and Chemical Sciences* 6: 2252-2262.
- Touré N, Yao-Kouamé A, Alui KA, Guety PT (2010) Assessment of major elements and metallic traces of an agricultural production environment in the Niéki valley in southeastern Côte d'Ivoire. *Journal of Applied Biosciences* 34: 2134-2144.
- Kouamé KB, Koné WM (2017) Evaluation of the treatment and storage conditions of medicinal plants in the markets of the District of Abidjan (Côte d'Ivoire). *European Scientific Journal* 13: 396-415.
- INS (2014) General Population and Housing Census 2014, Implementation Report and presentation of the main results. National Institute of Statistics, Côte d'Ivoire, p: 49.
- IITA (2001) Analysis of soil and plant samples. International Institute of Tropical Agriculture Manuel no 1, p: 160.
- AOAC (1990) Analytical chemists, food composition, Official Analytical Chemists, additives natural contaminant. Official Analytical Chemists, Additives Natural Contaminant, Adrich RC, USA, p: 256.

24. Cissé M, Dornier M, Sakho M, Ndiaye A, Reynes M, et al. (2009) Bissap (*Hibiscus sabdariffa* L.): composition and main uses. Fruits 64: 179-193.
25. Thongam CA, Thokchom R, Subapriya MS, Thokchom J, Singh SS (2016) *Hibiscus sabdariffa*-A natural micro nutrient source. Int J Adv Res Biol Sci 3: 243-248.
26. Arbaoui S, Paul R, Bettaieb T (2015) Phytoextraction of zinc and cadmium by kenaf (*Hibiscus cannabinus* L.) grown on soil polluted with trace metals. Nature & Technology, pp: 108-114.
27. Guety TP, Kone B, Yao GF (2015) Concentrations of Cadmium, Copper, Lead and Zinc in Soils and Vegetable Organs from Periurban Agriculture Areas of Abidjan in Cote d'Ivoire. Journal of Agriculture and Ecology Research International 3: 12-23.
28. FAO/WHO (2015) Report of the Codex Alimentarius on the General Standard for Contaminants and Toxins in Foods and Animal Products Codex Stan 193-1995. Codex Alimentarius, Food and Agriculture Organization of the United Nations and World Health Organization, p: 34.
29. Orisakwe OE, Husaini DC, Afonne OJ (2004) Testicular effects of sub-chronic administration of *Hibiscus sabdariffa* calyx aqueous extract in rats. Reproductive Toxicology 18: 295-298.
30. Hamadouche AN (2010) Effects of Chronic Lead Exposure on the Reproductive System and the Hypothalamic-Pituitary Axis in the Wistar Male Rat: Histological and Biochemical Study. Thesis of Doctorate of State in Applied Biochemistry. Ahmed Ben Bella University of Oran1 Es Senia, Algeria, p: 241.
31. Onyeke NG (2013) Development of Products from Sorrel (*Hibiscus Sabdariffa*) and Evaluation of their Nutrients, Phytochemicals, Antinutrients, Toxicants, Physico-Chemical, Sensory and Storage Properties, Doctoral thesis of the University of Nsukka, Nigeria, p: 164.
32. Ajala LO, Oyeleke GO, Olagunju EO (2013) Nutritional assessment of juice extract from *Hibiscus sabdariffa*. Journal of Biology, Pharmacy and Allied Sciences 1: 10-19.
33. Bakare-Odunola MT, Mustapha KB (2014) Identification and quantification of heavy metals in local drinks in Northern Zone of Nigeria. Journal of Toxicology and Environmental Health Sciences 6: 126-131.
34. FAO/WHO (2000) Proposed Draft Revised Codex Standard for Vegetable Juices CODEX STAN 179-1991. Codex Alimentarius, Food and Agriculture Organization of the United Nations and World Health Organization, Rome (Italy), p: 5.
35. WHO (2017) International Program on Chemical Safety, Sections: Ten chemicals that pose a major public health problem.
36. Najafi S, Moshtaghie AA, Noori A (2016) Protective Effects of Long Term Administration of Zinc on Bone Metabolism Parameters in Male Wistar Rats Treated with Cadmium. Zahedan University of Medical Sciences, pp: 1-6.
37. Berroukche A, Slimani M, Kahloulou K, Kafi H, Cheikh A (2014) Evaluation of the activity of cadmium, in the presence of zinc, on structures of tissues regulating metabolism in the Wistar rat. International Journal of Biological and Chemical Sciences 8: 1796-1807.
38. Prasad AS (2013) Essential and Toxic Element: Trace Elements in Human Health and Disease. Elsevier, p: 544.
39. Chimienti F, Seve M, Richard S, Mathieu J, Favier A (2001) Role of cellular zinc in programmed cell death: temporal relationship between zinc depletion, activation of caspases, and cleavage of Sp family transcription factors. Biochemical Pharmacology 62: 51-62.
40. Schroeder JJ, Cousins RJ (1990) Interleukin 6 regulates metallothionein gene expression and zinc metabolism in hepatocyte monolayer cultures. Proceedings of the National Academy of Sciences 87: 3137-3141.
41. Chia SE, Ong CN, Chua LH, Ho LM, Tay SK (2000) Comparison of zinc concentrations in blood and seminal plasma and the various sperm parameters between fertile and infertile men. Journal of Andrology 21: 53-57.
42. Mourtzinou I, Makris DP, Yannakopoulou K, Kalogeropoulos N, Michali I, et al. (2008) Thermal Stability of Anthocyanin Extract of *Hibiscus sabdariffa* L. in the Presence of β -Cyclodextrin. Journal of Agricultural and Food Chemistry 56: 10303-10310.
43. Ndiaye NA, Dieng M, Kane A, Cisse M, Montet D, et al. (2015) Diagnosis and microbiological characterization of artisanal processes for the manufacture of beverages and concentrates of *Hibiscus sabdariffa* L in Senegal. Africa Science, International Journal of Science and Technology 11: 197-210.
44. Noura B, Tchame G, Tchiosah R, Ngang JJE, Etoa FX (2014) Influence du procédé de fabrication sur la qualité microbiologique du jus de folere (*Hibiscus sabdariffa*) vendu dans trois villes du Cameroun: Maroua, Mokolo et Mora. Innovative Space of Scientific Research Journals 9: 786-796.
45. Jaishankar M, Tseten T, Anbalagan N, Mathew BB, Beeregowda KN (2014) Toxicity, mechanism and health effects of some heavy metals. Interdisciplinary Toxicology 7: 60-72.
46. Markiewicz-Górka I, Januszewska L, Michalak A, Prokopowicz A, Januszewska E, et al. (2015) Effects of chronic exposure to lead, cadmium, and manganese mixtures on oxidative stress in rat liver and heart. Archives of Industrial Hygiene and Toxicology 66: 51-62.
47. Belli N, Mesbah L, Chebab S, Tekouk M, Leghouchi E (2010) Oxidative stress induced by coexposure to lead and cadmium: two groundwater contaminants from Oued Nil (Jijel-Algeria). Revue des Sciences de l'Eau 23: 289-301.
48. WHO (2003) Diet, nutrition and chronic disease prevention. Technical Report Series, Report of a WHO/FAO Expert Consultation (Geneva), p: 127.