



Bioenergetics and Type 2 Diabetes

Doliba NM*

Department of Biochemistry and Biophysics, Institute for Diabetes, Obesity and Metabolism, University of Pennsylvania School of Medicine, PA, USA

*Corresponding author: Doliba NM, Department of Biochemistry, Biophysics and Institute for Diabetes, Obesity and Metabolism, University of Pennsylvania School of Medicine, PA 19104, USA, Tel: 215-898-4366; Fax: 215-898-5408; E-mail: nicolai@mail.med.upenn.edu

Received date: Jan 05, 2016; Accepted date: Jan 16, 2017; Published date: Jan 23, 2017

Copyright: © 2017 Doliba NM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Hyperglycemia of type 2 diabetes mellitus (T2DM) develops when pancreatic β -cells damaged by chronic exposure to elevated blood glucose and lipids (glucolipototoxicity) fail to synthesize and secrete sufficient quantities of insulin for maintaining plasma glucose level. Despite intensive studies in this field, the molecular mechanism by which fatty acids (FA) cause β -cell impairment is not well understood. It still remains unknown what are the lipid- or glucose-derived molecules directly responsible for the impairment of β -cell function? Our studies showed that in addition to impaired insulin secretion and loss of biphasicity, T2D islets exhibited altered cell bioenergetics as evidenced by decreased oxygen consumption rate as compared to those of the control islets. We also discovered that fuel overload (high level of FA and glucose leads to incomplete FA oxidation and results in accumulation of "toxic" long-chain 3-OH-fatty acids that could induce oxidative stress and disrupt mitochondrial function. Time-dependent impairments of bioenergetics due to chronic exposure to elevated blood glucose and lipids would be the consequence leading to pancreatic β -cell failure.

Keywords: Bioenergetics; Pancreatic islets; Fatty acids; Glucolipototoxicity; Type 2 diabetes; Insulin secretion

Glucolipototoxicity and Diabetes

Overconsumption of calorie rich diets increases the incidence of type 2 diabetes mellitus (T2DM) in genetically predisposed individuals, which has resulted in the worldwide epidemic of T2DM now afflicting about 350 million people. Hyperglycemia of T2DM develops when pancreatic β -cells damaged by chronic exposure to elevated blood glucose and lipids fail to synthesize and secrete sufficient quantities of insulin for maintaining plasma glucose level at a critical level of 5 mM. "Glucolipototoxicity", the operationally defined condition resulting from caloric overload, is proposed to worsen or cause β -cell damage which eventually leads to T2D. The term "glucolipototoxicity" implies that repeated or continued exposure to high blood glucose and lipid levels are required for β -cell damage and functional dysfunction to occur. However, a compelling mechanistic molecular explanation of "glucolipototoxicity" affecting pancreatic β -cells is still lacking. In an attempt to model "glucolipototoxicity" *in vitro*, pancreatic islets are usually cultured for several days in high glucose and fatty acids (FA) concentrations. Studies have described multiple cellular processes involved in the pathogenesis of β -cell dysfunction, including changes in gene expression [1,2] intermediary metabolism [3], mitochondrial function [4], ion channel activity [5-7], insulin synthesis and exocytosis [8]. Different molecular mechanisms of FA-induced β -cell dysfunction have been proposed including accumulation of ceramide [9], apoptosis of β -cells due to oxidative [10-11] and endoplasmic reticulum (ER) stress [12,13] as well as others mechanisms [14,15]. Many of these mechanisms remain controversial. For example, the exposure of human islets for 24 hours to elevated FA and glucose conditions was found in one study to initiate apoptosis [16], but other studies have failed to find evidence of any significant apoptosis following long-term exposure. Olofsson et al. reported that

inhibition of glucose-stimulated insulin secretion (GSIS) by long-term exposure to the FAs oleate and palmitate was not related to any signs of increased β -cell death, reduced insulin synthesis, impaired glucose metabolism, KATP channel regulation, or Ca^{2+} signaling. These discrepancies could be due to differences in acute or chronic islet responses [17,18]. Albumin/FA ratios in *in vitro* studies are often suboptimal (i.e., $\leq 3:1$), glucose concentrations are often excessive to be meaningful (i.e., >16 mM), there are significant limitations inherent in animal models, and there is a lack of a clear definition of the "glucolipototoxicity" phenomenon [19]. While various molecular and cellular mechanisms of glucolipototoxicity and their roles in obesity and diabetes have been described in animal models [19-22], it is unclear whether these models recapitulate the pathogenesis of human T2D.

Pancreatic Islet Bioenergetics and Diabetes

A faulty bioenergetic process is a plausible explanation for defective insulin secretion in T2D. Normally the ATP, generated in glycolysis and oxidative phosphorylation by catabolism of glucose, amino acids, and FAs, serves as coupling factor in fuel stimulated insulin secretion [23]. The unique role of ATP as a critical messenger in the stimulus-secretion coupling was clearly shown in our previous studies with mouse, rat and human islets where oxygen consumption, glycolysis, glucose oxidation were related to insulin secretion [24]. Such measurements allowed us to calculate the ATP production rate in pancreatic β -cells as a function of the glucose concentration and insulin secretion. Our calculation was based on a reasonable assumption that islet glycogen stores are negligible [25] and that coupling of oxidative phosphorylation is intact. Despite major differences in insulin profiles (Figure 1A), the ATP-Production/Insulin-Secretion curves were similar for mouse, rat and human islets. The data for all species fitted a single sigmoidal curve (Figure 1B), indicating a clear relationship between ATP production rate and insulin secretion [24].




Figure 1: Fusion of mouse, rat and human islets ATP-production/insulin-secretion diagrams. Panel A: Secretion profiles of mouse, rat, and human islets as function of the glucose concentration. Panel B: Combined energy production/insulin-secretion curve with data on ATP production and insulin release from all 3 types of islets projected on 1 single continuous sigmoidal curve. ATP production was calculated based on oxygen consumption and glucose usage data.

The ATP-Production/Insulin-Secretion curve has a threshold for glucose stimulated insulin secretion at about 16 pmole/islet/min and a Hill coefficient of about 11. This data indicate a highly cooperative coupling mechanism between ATP production and insulin secretion with half-maximal effective rate (ER50) of 22 pmole/islet/min. We speculate that the "ATP-Production/Insulin-Secretion" curve has clinical significance comparable to that of the classical "Frank-Starling" curve of the heart. We also showed that ATP-Production/Insulin-Secretion curve is modified by GLP-1 and a glucokinase activator piragliatin [24]. We speculate that the "ATP-Production/Insulin-Secretion" curve is modified in T2D islets.


The literature proposes various ways by which mitochondrial energy metabolism can be altered due to fuel overload. FAs may act as uncouplers and inhibitors of mitochondrial respiration and oxidative phosphorylation [26], operating as protonophors and by inhibiting the electron transport [27-29]. In addition, FAs increase the expression of Uncoupling Protein (UCP2) in pancreatic islets [14,30-31]. FAs may inhibit complex I [32] and can also inhibit ATP and ADP exchanges [33]. FAs may promote opening of the permeability transition pore [34-37].

FAs also increase the expression of PGC-1 α which may alter bioenergetics in pancreatic β -cells [38]. PGC-1 α is elevated in islets from different animal models of diabetes and in human studies [38-41]. PGC-1 α promotes mitochondrial biogenesis in brown tissue [42], however adenovirus-mediated expression of PGC-1 α led to a marked inhibition of glucose-stimulated insulin secretion [38], by suppressing glucose oxidation or decreasing the cell's ability to drive ATP production. PGC-1 α increases the transcription of UCP2 [43] by PGC-1-mediated upregulation of β -cell sterol element binding protein (SREBP) gene expression. The higher expression of UCP2 may result in a decreased efficiency of ATP production [44] by translocation of protons across the mitochondrial membrane. The latest effect should lead to changes in oxygen consumption and oxidative ATP synthesis. However, such data are limited and related to insulinoma cells [45] and only measurements of total ATP reported in islets exposed to FA [8]. Because in pancreatic islets ATP is co-secreted with insulin, it is difficult to dissociate between the effects of FA on ATP syntheses and

changes of ATP content in insulin granules. In fact, the insulin content is decreased in islets chronically exposed to FFA [8].

In order to access β -cell bioenergetics and its relationship to insulin secretion, we performed two sets of experiments: (i) The bioenergetics, ionic and secretion profiles of pancreatic islets isolated from healthy and T2D organ donors were examined; (ii) Isolated normal human islets were exposed to a glucolipototoxicity condition (high glucose and FFAs) in organ culture and the bioenergetics and insulin secretion were studied in perfusion experiments. Diabetic islets exposed to a "staircase" increase in glucose concentration in the perfusion setup showed a significant decrease in insulin secretion profile (Figure 2A) and oxygen consumption (Figure 2C) as compared to those of the control islets. The baselines for both parameters are comparable. The difference in insulin secretion profile is most pronounced at the 6 and 12 mM glucose stimulation (Figure 2B) indicating decreased rates of insulin secretion and loss of biphasicity. The glucose-dependency curve of the OCR of the diabetic islets was right shifted and reduced by 50% (Figure 2C).

The uncoupler of respiration and oxidative phosphorylation FCCP (5 μ M) stopped insulin secretion instantly and transiently increased respiration in control and diabetic islets to the same extent indicating a strong coupling between islets respiration and oxidative phosphorylation in both types of islets (Figure 2C). Glucose dependency of oxygen consumption (VO₂) was sigmoidal for both type of islets (Figure 2D). However, islets from T2D exhibited a right shift and lower maximal stimulation of respiration (V_{max}) by glucose. The S0.5 in T2D islets rose from 4.39 \pm 0.01 in control to 5.43 \pm 0.13 mM (Figure 2D). Panel E of Figure 2 presents changes in intracellular [Ca²⁺] of human islets. Low glucose (3 mM) produced a transient and high glucose (9 mM) produced a biphasic and a sustained increase in [Ca²⁺] of control islets. Diabetic islets did not respond to 3 mM glucose and the responses to 9 mM glucose were delayed and lower than in controls.



This data indicates that impaired pancreatic islet β -cell bioenergetics resulting in reduced ATP production is a critical factor in the molecular pathogenesis of T2D. Importantly, the glucokinase activator piragliatin was able to correct the defect of respiration and glucose-stimulated insulin secretion [24]. In our second set of experiments, the glucolipotoxicity which is a hallmark of T2D, was mimicked *in vitro* by culturing the islets for 3 or 5 days with 0.5 mM palmitic acid or a mixture of palmitic and oleic acid at 1% albumin, and different concentrations of glucose: 10, 16 and 25, 46. As a result, chronic

exposure of mouse islets to glucolipotoxic condition led to bioenergetic failure, as evidenced by decreased OCR and reduced glucose-stimulated insulin secretion. In addition, the islet ATP levels and glucose induced ATP rise were reduced as well as mitochondrial DNA and expression of mitochondrial transcription factor A (Tfam). We also discovered accumulation of carnitine esters of hydroxylated long chain FA [47], that have been shown to uncouple the respiration and oxidative phosphorylation in heart and brain mitochondria [48,49]. We propose that mitochondrial accumulation of unsaturated hydroxylated long-chain FA uncouples and ultimately inhibits pancreatic islet β -cell respiration and that this effect of the toxic FA metabolite causes a slow decline of mitochondrial ATP production resulting in bioenergetic failure as the main cause of impaired insulin secretion and reduced β -cell mass, both hallmarks of T2D.

Effects of Fatty Acids in Humans

FAs play an essential role in the function of pancreatic β -cell in humans [50]. In the fasting state, free FAs support basal insulin secretion and promote efficient nutrient-stimulated insulin secretion when the fast is terminated. Elevated plasma FFA levels lead to chronic hyperinsulinemia in insulin-resistant obese subjects [51]. Removal of this FFA stimulus by overnight reduction of plasma FFAs with nicotinic acid impairs insulin secretion stimulated by glucose [51]. Despite the evidence from *in vivo* studies, the effects of prolonged elevation of FFA on insulin secretion in humans remain controversial. Boden and colleagues demonstrated that 48-h elevation of plasma FFA potentiated glucose-stimulated insulin secretion in healthy subjects at glucose levels clamped at 8.6 mM [52] but insulin secretion was defective in T2D patients [53]. In contrast, Carpentier et al. [54] have reported that acute stimulation of insulin secretion by FA in healthy humans is lost with chronic FA elevation. This loss of hormone secretion was specific to glucose because the response to arginine was normal [55]. It is of interest that obese but not diabetic subjects are more sensitive to the inhibitory effect of lipids on glucose-stimulated insulin secretion [56]. Kashyap et al. [57] have studied both insulin secretion and insulin action in normal subjects with and without a family history of T2D during a 4-day lipid infusion. The most striking finding is that a 4-day intralipid infusion stimulated insulin secretion in normal subjects but inhibits glucose-stimulated insulin secretion in individuals with family history of T2D [57]. These data suggest that β -cell lipotoxicity may play an important role in the progression from normal glucose tolerance to overt hyperglycemia in subjects with a high risk of developing T2D. In support of this data, the antilipolytic agent acipimox improved the first phase of insulin secretion in nondiabetic patients with a family history of type 2 diabetes [58].

Conclusion

To conclude, based on the existing literature it is clear that excessive glucose and fatty acids levels have time-dependent deteriorating effects on pancreatic β -cell pathophysiology in diabetes. These effects are different at the various stages of β -cell dysfunction during development of T2D. When insulin resistance develops the β -cells mount a compensatory response that increases insulin biosynthesis, insulin secretion and β -cell mass. This compensatory β -cell response is genetically determined [57,59-60] and plays an important role in the long-term ability to maintain glucose homeostasis during insulin resistance. In individuals that genetically predisposed to diabetes, β -cell compensation eventually may become insufficient to maintain a

secretory response that meets the demand imposed by insulin resistance.

The failure of β -cells to compensate for insulin resistance is a major component of impaired glucose homeostasis and overt diabetes. This defect is the consequence of a decline of insulin response to glucose due to functional β -cell deficiency. Bioenergetics of β -cell plays an essential role in stimulus-secretion coupling and contributes to molecular pathogenesis of T2D. It is also the consequence of an inability of the endocrine pancreas to adapt the β -cell mass which eventually leads to a decrease in functional β -cells. This idea has resulted in considerable attention being paid to the development of new therapeutic strategies aimed toward preserving or regenerating functional β -cell mass [61]. GLP-1 enhances β -cell survival by activating β -cell proliferation and differentiation, and inhibiting β -cell apoptosis and thus contributing to the long-term regulation of insulin secretion by maintaining a functional β -cell mass. It should be pointed out that any intervention to improve insulin secretion should start early in the disease when the endogenous insulin secretion and presumably the number of functional β -cells has not decreased excessively [62-64].

References

- Biden TJ, Robinson D, Cordery D, Hughes WE, Busch AK (2004) Chronic effects of fatty acids on pancreatic beta-cell function: new insights from functional genomics. *Diabetes* 1: S159-S165.
- Gremlich S, Bonny C, Waeber G, Thorens B (1997) Fatty acids decrease IDX-1 expression in rat pancreatic islets and reduce GLUT2, glucokinase, insulin, and somatostatin levels. *J Biol Chem* 272: 30261-30269.
- Iizuka K, Nakajima H, Namba M, Miyagawa J, Miyazaki J, et al. (2002) Metabolic consequence of long-term exposure of pancreatic beta cells to free fatty acid with special reference to glucose insensitivity. *Biochim Biophys Acta* 1586: 23-31.
- Koshkin V, Wang X, Scherer PE, Chan CB, Wheeler MB (2003) Mitochondrial functional state in clonal pancreatic beta-cells exposed to free fatty acids. *J Biol Chem* 278: 19709-19715.
- Branstrom R, Corkey BE, Berggren PO, Larsson O (1997) Evidence for a unique long chain acyl-CoA ester binding site on the ATP-regulated potassium channel in mouse pancreatic beta cells. *J Biol Chem* 272: 17390-17394.
- Branstrom R, Leibiger IB, Leibiger B, Corkey BE, Berggren PO, et al. (1998) Long chain coenzyme A esters activate the pore-forming subunit (Kir6. 2) of the ATP-regulated potassium channel. *J Biol Chem* 273: 31395-31400.
- Reimann F, Huopio H, Dabrowski M, Proks P, Gribble FM, et al. (2003) Characterisation of new KATP-channel mutations associated with congenital hyperinsulinism in the Finnish population. *Diabetologia* 46: 241-249.
- Somesh BP, Verma MK, Sadasivuni MK (2013) Chronic glucolipotoxic conditions in pancreatic islets impair insulin secretion due to dysregulated calcium dynamics, glucose responsiveness and mitochondrial activity. *BMC Cell Biol* 14: 31.
- Boslem E, Meikle PJ, Biden TJ (2012) Roles of ceramide and sphingolipids in pancreatic beta-cell function and dysfunction. *Islets* 4: 177-187.
- Gehrmann W, Elsner M, Lenzen S (2010) Role of metabolically generated reactive oxygen species for lipotoxicity in pancreatic beta-cells. *Diabetes Obes Metab* 2: 149-158.
- Morgan D, Oliveira-Emilio HR, Keane D (2007) Glucose, palmitate and pro-inflammatory cytokines modulate production and activity of a phagocyte-like NADPH oxidase in rat pancreatic islets and a clonal beta cell line. *Diabetologia* 50: 359-369.
- Kharroubi I, Ladriere L, Cardozo AK, Dogusan Z, Cnop M, et al. (2004) Free fatty acids and cytokines induce pancreatic beta-cell apoptosis by different mechanisms: role of nuclear factor-kappa B and endoplasmic reticulum stress. *Endocrinology* 145: 5087-5096.
- Cnop M, Igoillo-Estevé M, Cunha DA, Ladriere L, Eizirik DL (2008) An update on lipotoxic endoplasmic reticulum stress in pancreatic beta-cells. *Biochem Soc Trans* 36: 909-915.
- Joseph JW, Koshkin V, Saleh MC (2004) Free fatty acid-induced beta-cell defects are dependent on uncoupling protein 2 expression. *J Biol Chem* 279: 51049-51056.
- Cnop M, Welsh N, Jonas JC, Jorns A, Lenzen S, et al. (2005) Mechanisms of pancreatic beta-cell death in type 1 and type 2 diabetes: many differences, few similarities. *Diabetes* 2: S97-S107.
- El-Assaad W, Buteau J, Peyot ML (2003) Saturated fatty acids synergize with elevated glucose to cause pancreatic beta-cell death. *Endocrinology* 144: 4154-4163.
- Kelpe CL, Moore PC, Parazzoli SD, Wicksteed B, Rhodes CJ, et al. (2003) Palmitate inhibition of insulin gene expression is mediated at the transcriptional level via ceramide synthesis. *J Biol Chem* 278: 30015-30021.
- Olofsson CS, Collins S, Bengtsson M, Eliasson L, Salehi A, et al. (2007) Long-term exposure to glucose and lipids inhibits glucose-induced insulin secretion downstream of granule fusion with plasma membrane. *Diabetes* 56: 1888-1897.
- Poitout V, Amyot J, Semache M, Zarrouki B, Hagman D, et al. (2010) Glucolipotoxicity of the pancreatic beta cell. *Biochimica et Biophysica Acta* 1801: 289-298.
- Prentki M, Matschinsky FM, Madiraju SR (2013) Metabolic signaling in fuel-induced insulin secretion. *Cell Metab* 18: 162-185.
- Prentki M, Madiraju SR (2012) Glycerolipid/free fatty acid cycle and islet beta-cell function in health, obesity and diabetes. *Mol Cell Endocrinol* 353: 88-100.
- Poitout V, Robertson RP (2008) Glucolipotoxicity: fuel excess and beta-cell dysfunction. *Endocr Rev* 29: 351-366.
- Ashcroft FM, Harrison DE, Ashcroft SJ (1984) Glucose induces closure of single potassium channels in isolated rat pancreatic beta-cells. *Nature* 312: 446-448.
- Doliba NM, Qin W, Najafi H, Liu C, Buettger CW, et al. (2012) Glucokinase activation repairs defective bioenergetics of islets of Langerhans isolated from type 2 diabetics. *Am J Physiol Endocrinol Metab* 302: E87-E102.
- Matschinsky FM, Ellerman JE, Landgraf, krzanowski J, Kotler-Brajtburg J, et al. (1971) Quantitative histochemistry of glucose metabolism in the islets of Langerhans. *Curr Probl Clin Biochem* 3: 143-182.
- Wojtczak L, Schonfeld P (1993) Effect of fatty acids on energy coupling processes in mitochondria. *Biochimica et Biophysica Acta* 1183: 41-57.
- Schonfeld P, Reiser G (2006) Rotenone-like action of the branched-chain phytanic acid induces oxidative stress in mitochondria. *J Biol Chem* 281: 7136-7142.
- Schonfeld P, Wojtczak L (2007) Fatty acids decrease mitochondrial generation of reactive oxygen species at the reverse electron transport but increase it at the forward transport. *Biochimica et Biophysica Acta* 1767: 1032-1040.
- Schonfeld P, Wojtczak L (2008) Fatty acids as modulators of the cellular production of reactive oxygen species. *Free Radical Biol Med* 45: 231-241.
- Chan CB, Saleh MC, Koshkin V, Wheeler MB (2004) Uncoupling protein 2 and islet function. *Diabetes* 1: S136-S142.
- Zhang CY, Baffy G, Perret P (2001) Uncoupling protein-2 negatively regulates insulin secretion and is a major link between obesity, beta cell dysfunction, and type 2 diabetes. *Cell* 105: 745-755.
- Loskovich MV, Grivennikova VG, Cecchini G, Vinogradov AD (2005) Inhibitory effect of palmitate on the mitochondrial NADH:ubiquinone oxidoreductase (complex I) as related to the active-de-active enzyme transition. *Biochem J* 387: 677-683.

33. Klingenberg M (2008) The ADP and ATP transport in mitochondria and its carrier. *Biochimica et Biophysica Acta* 1778: 1978-2021.
34. Scorrano L, Penzo D, Petronilli V, Pagano F, Bernardi P (2001) Arachidonic acid causes cell death through the mitochondrial permeability transition. Implications for tumor necrosis factor- α apoptotic signaling. *J Biol Chem* 276: 12035-12040.
35. Bernardi P, Penzo D, Wojtczak L (2002) Mitochondrial energy dissipation by fatty acids. Mechanisms and implications for cell death. *Vitamins and Hormones* 65: 97-126.
36. Penzo D, Tagliapietra C, Colonna R, Petronilli V, Bernardi P (2002) Effects of fatty acids on mitochondria: implications for cell death. *Biochimica et Biophysica Acta* 1555: 160-165.
37. Penzo D, Petronilli V, Angelini A, Cusan C, Colonna R, et al. (2004) Arachidonic acid released by phospholipase A(2) activation triggers Ca(2+)-dependent apoptosis through the mitochondrial pathway. *J Biol Chem* 279: 25219-25225.
38. Yoon JC, Xu G, Deeney JT, Yang SN, Rhee J, et al. (2003) Suppression of beta cell energy metabolism and insulin release by PGC-1 α . *Dev Cell* 5: 73-83.
39. Ek J, Andersen G, Urhammer SA, Gaede PH, Drivsholm T, et al. (2001) Mutation analysis of peroxisome proliferator-activated receptor- γ coactivator-1 (PGC-1) and relationships of identified amino acid polymorphisms to Type II diabetes mellitus. *Diabetologia* 44: 2220-2226.
40. Oberkofler H, Linnemayr V, Weitgasser R, Klein K, Xie M, et al. (2004) Complex haplotypes of the PGC-1 α gene are associated with carbohydrate metabolism and type 2 diabetes. *Diabetes* 53: 1385-1393.
41. Oberkofler H, Hafner M, Felder T, Krempler F, Patsch W (2009) Transcriptional co-activator peroxisome proliferator-activated receptor (PPAR) γ co-activator-1 β is involved in the regulation of glucose-stimulated insulin secretion in INS-1E cells. *J Mol Med* 87: 299-306.
42. Puigserver P, Wu Z, Park CW, Graves R, Wright M (1998) A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 92: 829-839.
43. Oberkofler H, Klein K, Felder TK, Krempler F, Patsch W (2006) Role of peroxisome proliferator-activated receptor- γ coactivator-1 α in the transcriptional regulation of the human uncoupling protein 2 gene in INS-1E cells. *Endocrinology* 147: 966-976.
44. Klingenberg M, Huang SG (1999) Structure and function of the uncoupling protein from brown adipose tissue. *Biochim Biophys Acta* 1415: 271-296.
45. Barlow J, Affourtit C (2013) Novel insights into pancreatic beta-cell glucolipotoxicity from real-time functional analysis of mitochondrial energy metabolism in INS-1E insulinoma cells. *Biochem J* 456: 417-426.
46. Doliba NM, Chen P, Chen J, Naji A, Bennet M, et al. (2015) Long-chain beta-hydroxylated fatty acids as mediators of pancreatic beta-cell bioenergetic failure. American Diabetes Association, Boston, MA: Diabetes p: 2354.
47. Doliba NM, Changong LI, Chen P, Chen J, Naji A (2015) Long-Chain beta-Hydroxylated Fatty Acids as Mediators of Pancreatic beta-Cell Bioenergetics Failure Diabetes.
48. Tonin AM, Amaral AU, Busanello EN, Grings M, Castilho RF, et al. (2013) Long-chain 3-hydroxy fatty acids accumulating in long-chain 3-hydroxyacyl-CoA dehydrogenase and mitochondrial trifunctional protein deficiencies uncouple oxidative phosphorylation in heart mitochondria. *J Bioenerg Biomembr* 45: 47-57.
49. Tonin AM, Ferreira GC, Grings M, Viegas CM, Busanello EN, et al. (2010) Disturbance of mitochondrial energy homeostasis caused by the metabolites accumulating in LCHAD and MTP deficiencies in rat brain. *Life Sci* 86: 825-831.
50. McGarry JD (2002) Banting lecture 2001: dysregulation of fatty acid metabolism in the etiology of type 2 diabetes. *Diabetes* 51: 7-18.
51. Dobbins RL, Chester MW, Daniels MB, McGarry JD, Stein DT (1998) Circulating fatty acids are essential for efficient glucose-stimulated insulin secretion after prolonged fasting in humans. *Diabetes* 47: 1613-1618.
52. Boden G, Chen X, Rosner J, Barton M (1995) Effects of a 48-h fat infusion on insulin secretion and glucose utilization. *Diabetes* 44: 1239-1242.
53. Boden G, Chen X (1999) Effects of fatty acids and ketone bodies on basal insulin secretion in type 2 diabetes. *Diabetes* 48: 577-583.
54. Carpentier A, Mittelman SD, Lamarche B, Bergman RN, Giacca A (1999) Acute enhancement of insulin secretion by FFA in humans is lost with prolonged FFA elevation. *Am J Physiol* 276: E1055-E1066.
55. Carpentier A, Giacca A, Lewis GF (2001) Effect of increased plasma non-esterified fatty acids (NEFAs) on arginine-stimulated insulin secretion in obese humans. *Diabetologia* 44: 1989-1997.
56. Carpentier A, Mittelman SD, Bergman RN, Giacca A, Lewis GF (2000) Prolonged elevation of plasma free fatty acids impairs pancreatic beta-cell function in obese nondiabetic humans but not in individuals with type 2 diabetes. *Diabetes* 49: 399-408.
57. Kashyap S, Belfort R, Gastaldelli A, Pratipanawatr T, Berria R, et al. (2003) A sustained increase in plasma free fatty acids impairs insulin secretion in nondiabetic subjects genetically predisposed to develop type 2 diabetes. *Diabetes* 52: 2461-2474.
58. Paolisso G, Tagliamonte MR, Rizzo MR, Gualdiero P, Saccomanno F, et al. (1998) Lowering fatty acids potentiates acute insulin response in first degree relatives of people with type II diabetes. *Diabetologia* 41: 1127-1132.
59. Fadista J, Vikman P, Laakso EO, Mollet GI, Esguerra JL, et al. (2014) Global genomic and transcriptomic analysis of human pancreatic islets reveals novel genes influencing glucose metabolism. *Proc Natl Acad Sci* 111: 13924-13929.
60. Rosengren AH, Braun M, Mahdi T, Andersson SA, Travers ME, et al. (2012) Reduced insulin exocytosis in human pancreatic beta-cells with gene variants linked to type 2 diabetes. *Diabetes* 61: 1726-1733.
61. Karaca M, Magnan C, Kargar C (2009) Functional pancreatic beta-cell mass: involvement in type 2 diabetes and therapeutic intervention. *Diabetes Metabol* 35: 77-84.
62. Grill V, Bjorklund A (2002) Type 2 diabetes-effect of compensatory over secretion as a reason for beta-cell collapse. *Int J Exp Diabetes Res* 3: 153-158.
63. Karvestedt L, Andersson G, Efendic S, Grill V (2002) A rapid increase in beta-cell function by multiple insulin injections in type 2 diabetic patients is not further enhanced by prolonging treatment. *J Inter Med* 251: 307-316.
64. Doliba NM, Qin W, Vatamaniuk MZ, Buettger CW, Collins HW, et al. (2006) Cholinergic regulation of fuel-induced hormone secretion and respiration of SUR1 mouse islets. *Am J Physiol Endocrinol Metab* 291: E525-E535.