OPEN ACCESS Freely available online

Journal of Physical Chemistry & Biophysics

Perspective

Biophysical Perspectives on Cellular Communication through Bioelectricity

Hyung Na*

Department of Celluar Biology, University of the South Pacific, Suva, Fiji

INTRODUCTION

Cellular communication is fundamental for maintaining physiological functions in multicellular organisms. One of the most essential forms of communication occurs through bioelectricity, which involves the transmission of electrical signals within and between cells. This type of communication plays an integral role in processes such as nerve signalling, muscle contraction, and tissue regeneration. Bioelectric signals are not merely electrical currents; they are also influenced by molecular and cellular properties that govern their behaviour and transmission.

DESCRIPTION

Bioelectricity in cellular communication

Bioelectricity refers to the electrical signals produced by the movement of ions across cell membranes. The electrical potential difference across the membrane of a cell, known as the membrane potential, is a key factor in bioelectric communication. This potential difference is created by the selective permeability of the cell membrane to ions, such as sodium, potassium, calcium, and chloride, which are actively transported in and out of the cell by ion channels, pumps, and exchangers.

The resting membrane potential of a cell typically ranges from -60 to -70 millivolts, depending on the cell type. When a cell is stimulated, ion channels open, allowing the rapid influx or efflux of ions, which generates a change in the membrane potential. This electrical activity can propagate along the membrane and influence neighbouring cells, either through direct cell-to-cell interactions or through more complex signalling pathways.

In excitable cells such as neurons and muscle cells, bioelectricity plays an important role in transmitting signals across long distances. The action potential, a rapid and transient change in membrane potential, is the fundamental electrical signal used by neurons to communicate. This electrical signal travels along the axon and is transmitted across synapses to other neurons or muscle cells, enabling communication within the nervous system and coordination of movement.

Ion channels and their role in bioelectric signalling

Ion channels are integral to bioelectric communication. These protein structures embedded in the cell membrane allow ions to flow in and out of the cell in response to various stimuli. Ion channels are highly selective, meaning that each channel typically permits the passage of only one type of ion. For instance, sodium channels allow sodium ions to flow in, while potassium channels allow potassium ions to flow out. This selective permeability is essential for maintaining the ion gradients that are necessary for cellular function.

The opening and closing of ion channels are regulated by various factors, including voltage, chemical signals, and mechanical stress. Voltage-gated ion channels, for example, open or close in response to changes in the membrane potential, while ligand-gated ion channels open in response to the binding of specific molecules, such as neurotransmitters. Mechanosensitive ion channels are activated by mechanical forces, such as pressure or stretch, which are important for processes like touch sensation and blood flow regulation.

These channels do not function in isolation but interact in highly coordinated networks. The coordinated activity of ion channels across many cells is essential for proper tissue function, including the synchronized contraction of heart muscle and the transmission of nerve impulses. Disruptions in ion channel function, such as mutations that alter channel activity, can lead to diseases like epilepsy, cardiac arrhythmias, and cystic fibrosis.

Bioelectricity in development and regeneration

Bioelectricity is not only important for the day-to-day function of cells but also plays a significant role in development and regeneration. During embryonic development, bioelectric signals are involved in shaping the body plan, guiding cell differentiation, and directing tissue morphogenesis. Recent studies have shown that altering bioelectric patterns can influence the fate of stem cells, potentially leading to new methods for regenerative medicine.

Correspondence to: Hyung Na, Department of Cellular Biology, University of the South Pacific, Suva, Fiji; E-mail: na_hyung@gmail.com

Received: 06-Jan-2025, Manuscript No. JPCB-25-36732; **Editor assigned:** 09-Jan-2025, PreQC No. JPCB-25-36732 (PQ); **Reviewed:** 23-Jan-2025, QC No. JPCB-25-36732; **Revised:** 18-Feb-2026, Manuscript No. JPCB-25-36732 (R); **Published:** 25-Feb-2026, DOI: 10.35841/2161-0398.26.16.444

Citation: Na H (2026) Biophysical Perspectives on Cellular Communication through Bioelectricity. J Phys Chem Biophys. 16:444.

Copyright: © 2026 Na H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

In regenerative processes, bioelectricity helps to coordinate the repair and replacement of damaged tissues. In animals like axolotls, which can regenerate entire limbs, bioelectric signals are thought to play a critical role in guiding the regeneration process. By modulating the electrical activity in the area surrounding the injury, it is possible to encourage cells to revert to a more plastic state, allowing them to proliferate and differentiate into the necessary tissue types to replace the lost structures.

Cellular communication beyond the individual cell

Bioelectricity also facilitates communication between cells. In tissues such as the heart, cells are electrically coupled, allowing the electrical signal generated by one cell to spread to adjacent cells. This forms a network of interconnected cells that can coordinate their activity, such as in the synchronous contraction of cardiac muscle during each heartbeat. This type of communication is achieved through gap junctions, which are specialized connections between cells that allow ions and small molecules to pass directly from one cell to another.

In addition to direct cell-to-cell communication, bioelectric signals can also influence the extracellular environment. For example, electrical fields can modulate the behaviour of cells in a tissue, guiding their migration and orientation. This type of communication is particularly important in processes like tissue formation, wound healing, and cancer metastasis.

Electrical fields can also affect the activity of ion channels and receptors, influencing the response of cells to other signalling pathways, such as those mediated by growth factors and hormones. Thus, bioelectric signals serve as an important layer of regulation that can fine-tune cellular responses to external cues, contributing to the overall function of tissues and organs.

Advances in bioelectric research

- Advances in bioelectric research have led to the development of new therapeutic strategies, including the use of electrical stimulation for tissue repair, nerve regeneration, and pain management.
- Bioelectric therapies, such as transcranial direct current stimulation (tDCS) and electroceuticals, are showing promise as non-invasive methods to treat a range of conditions, from depression to chronic pain.

CONCLUSION

Bioelectricity is a fundamental aspect of cellular communication that underlies many physiological processes, including development, regeneration, and disease. By understanding how bioelectric signals influence cellular behaviour, we gain valuable insights into the functioning of tissues and organs, as well as the mechanisms of disease. As research in bioelectricity continues to evolve, it holds the potential to provide new approaches for the treatment of a wide range of medical conditions, from neurological disorders to tissue regeneration.