

Caffeine supplementation changes inflammatory biomarkers after exercise

Sílvio Terra Stefanello¹, Félix Alexandre Antunes Soares^{1,2}, Rômulo Pillon Barcelos^{1,2*}

¹Departamento de Bioquímica e Biologia Molecular, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria, Santa Maria, CEP 97105-900, Brazil

²Laboratório de Bioquímica do Exercício (BioEx), Centro de Educação Física e Desportos, Universidade Federal de Santa Maria (UFSM), Santa Maria, RS, Brazil

Abstract

Caffeine is widely consumed by the world population due to its major constituent of beverages like coffee and tea. Additionally, caffeine supplementation is also taken by many athletes due to its ergogenic properties. Thus, several studies demonstrated that caffeine supplementation has significant antioxidant and anti-inflammatory activity. Our study evaluated the caffeine intake associated with swimming exercise in rats. We found that caffeine and exercise training did not change the mitochondrial parameters as compared with the control group. However, our results demonstrated that caffeine intake is highly efficacious in modulating inflammation biomarkers.

Keywords: Caffeine; Anti-inflammatory; Antioxidant; Mitochondria

Introduction

Caffeine is a methylxanthine (1,3,7-trimethylxanthine) that is highly consumed by people because it is a key component of beverages such as coffee and tea [1,2].

In addition, caffeine presents *in vitro* antioxidant activity, presenting effectiveness in scavenging mainly hydroxyl radicals [3,4]. *In vivo*, we have proved that caffeine intake has been associated with beneficial antioxidant effects in liver, such as decreased lipid peroxidation and elevated superoxide dismutase and glutathione peroxidase enzymes activity [5]. Recent studies have also been shown that caffeine intake prevents oxidative stress in neurodegenerative diseases and has beneficial effects against inflammatory injury [6-9]. In addition, clinical study demonstrated that coffee consumption decreased significantly the serum concentration of proinflammatory immune mediators [10].

Our study demonstrated that caffeine promotes a significant decrease in myeloperoxidase (MPO) activity, indicating that caffeine supplementation acts as anti-inflammatory, probably by reducing neutrophil infiltration and its inflammatory mechanisms activation [11]. Besides, the remarkable decrease in acetylcholinesterase (AChE) activity corroborated the anti-inflammatory effect, since this inhibition leads to an increase in acetylcholine levels, which is linked to anti-inflammatory functions and suppressed pro-inflammatory cytokines production [12-15]. Interestingly, most of these anti-inflammatory effects of caffeine are not associated with exercise training, but to an effect of caffeine *per se*.

Furthermore, caffeine supplementation is massively used by athletes during training or competitions [16]. Thus, several studies have been evaluating the mechanism of action of caffeine association with exercise training [17,18].

During the last years a consensus has emerged among sports scientists: ROS and inflammation are determinant to muscle cell regeneration and differentiation/proliferation, and the use of antioxidants and nonsteroidal anti-inflammatory drugs (NSAIDs) may blunt important exercise adaptations on young healthy subjects [19,20]. Therefore, drugs that mainly help on softening oxidative stress and inflammation after an exhaustive exercise should be preferred when comparing to the ones that blunt or completely block these necessary adaptive pathways of skeletal muscle regeneration. Among the natural compounds consumed on a daily basis, creatine (Cr, N-[aminoiminomethyl]-N-methyl glycine) and caffeine consumed by athletes [21,22]. Cr is among the most consumed ergogenic aids

by athletes mainly due to its positive effects on fatigue resistance, and improved exercise recovery after an exhaustive exercise [23]. On the same way, caffeine is considered an ergogenic compound due to its antioxidant and anti-inflammatory role. These effects were proved by our previous study as it is described here [5,11].

Original Study

Our study examined the effect of chronic caffeine intake and exercise training in male Wistar rats. The animals were divided into four groups (control, control/caffeine, exercise, and exercise/caffeine). Animals from Control and Exercise groups received saline, while groups Control/Caffeine and Exercise/Caffeine received caffeine supplementation (6 mg/kg/day) during the training protocol [24]. The exercise chosen was the swimming training, which consists of four weeks of swimming, 50 min per day and five sessions per week [25]. Our data identified that caffeine supplementation, associated or not with exercise training, did not promote any hepatic morphological or functional mitochondrial changes, maintaining a similar mitochondrial membrane potential, swelling, ROS production and mitochondrial complex I activity between groups. Nonetheless, both plasma myeloperoxidase and AChE activities were altered by the chronic caffeine intake (Figure 1). We attribute this decrease in MPO and AChE activity to caffeine anti-inflammatory effect, which is a view shared by other authors [7,26].

Final Considerations


In conclusion, we can observe that even though caffeine antioxidant properties, its intake did not modulate significantly the mitochondrial ROS production. In addition, we also verified that exercise training also did not change the liver mitochondrial parameters, indicating that the rats were well conditioned to the swimming training protocol. It should be noticed that caffeine intake did not interfere in this training adaptation.

***Corresponding author:** Rômulo Pillon Barcelos, Departamento de Bioquímica e Biologia Molecular – CCNE, Universidade Federal de Santa Maria - 97105-900 - Santa Maria, RS, Brazil, Tel: +55-55-3220-9522 (or) +55-55-3220-9558; E-mail: romulo1604@hotmail.com

Received March 02, 2015; Accepted April 21, 2016; Published April 28, 2016

Citation: Stefanello ST, Soares FAA, Barcelos RP (2016) Caffeine supplementation changes inflammatory biomarkers after exercise. J Yoga Phys Ther 6: 240. doi:10.4172/2157-7595.1000240

Copyright: © 2016 Stefanello ST, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Additionally, it is well known that exercise training produce *per se* an inflammatory process before the adaptation cycle. Thus, our study provided evidences indicating that caffeine supplementation can diminish this inflammatory process decreasing MPO and AChE activity.

Therefore, exercise training and caffeine supplementation may be a viable and effective strategy to diminish the side effects of inflammatory process. If this is a good or a bad effect in medium and long-term for training adaptation and health we cannot be sure right now. The research to find metabolic-related molecules able to modulate the oxidative-inflammatory metabolism but on the same way allow muscle cell differentiation/proliferation is of clinical interest, considering that different muscle disorder patients will benefit from this information.

References

- Ashihara H, Sano H, Crozier A (2008) Caffeine and related purine alkaloids: Biosynthesis, catabolism, function and genetic engineering. *Phytochemistry* 69: 841-856.
- Mejia EGD, Ramirez-Mares MV (2014) Impact of caffeine and coffee on our health. *Trends Endocrinol Metab* 25: 489-492.
- Brezova V, Slebodova A, Stasko A (2009) Coffee as a source of antioxidants: An EPR study. *Food Chemistry* 114: 859-868.
- León-Carmona JR, Galano A (2011) Is Caffeine a Good Scavenger of Oxygenated Free Radicals? *The Journal of Physical Chemistry B* 115: 4538-4546.
- Barcelos RP (2014) Caffeine supplementation modulates oxidative stress markers in the liver of trained rats. *Life Sciences* 96: 40-45.
- Lv X (2010) Caffeine protects against alcoholic liver injury by attenuating inflammatory response and oxidative stress. *Inflammation Research* 59: 635-645.
- Gaascht F, Dicato M, Diederich M (2015) Coffee provides a natural multitarget pharmacopeia against the hallmarks of cancer. *Genes & Nutrition* 10: 51.
- Ullah F (2015) Caffeine prevents d-galactose-induced cognitive deficits, oxidative stress, neuroinflammation and neurodegeneration in the adult rat brain. *Neurochemistry International* 90: 114-124.
- Souza MA (2013) Antioxidant activity elicited by low dose of caffeine attenuates pentylentetrazol-induced seizures and oxidative damage in rats. *Neurochemistry International* 62: 821-830.
- Kempf K (2010) Effects of coffee consumption on subclinical inflammation and other risk factors for type 2 diabetes: a clinical trial. *The American Journal of Clinical Nutrition* 91: 950-957.
- Barcelos R (2014) Caffeine Intake May Modulate Inflammation Markers in Trained Rats. *Nutrients* 6: 1678.
- Borovikova LV (2000) Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* 405: 458-62.
- Pavlov VA, Tracey KJ (2004) Neural regulators of innate immune responses and inflammation. *Cell Mol Life Sci* 61: 2322-2331.
- Pavlov VA, Tracey KJ (2006) Controlling inflammation: the cholinergic anti-inflammatory pathway. *Biochem Soc Trans* 34: 1037-1040.
- Wang H (2004) Cholinergic agonists inhibit HMGB1 release and improve survival in experimental sepsis. *Nat Med* 10: 1216-1221.
- Tunncliffe JM (2008) Consumption of dietary caffeine and coffee in physically active populations: physiological interactions. *Applied Physiology, Nutrition, and Metabolism* 33: 1301-1310.
- Tarnopolsky MA (2008) Effect of caffeine on the neuromuscular system — potential as an ergogenic aid. *Applied Physiology, Nutrition, and Metabolism* 33: 1284-1289.
- Lou Y (2013) Oral caffeine during voluntary exercise markedly inhibits skin carcinogenesis and decreases inflammatory cytokines in UVB-treated mice. *Nutr cancer* 65: 1002-1013.
- Lee S (2011) Mitochondrial H(2)O(2) generated from electron transport chain complex I stimulates muscle differentiation. *Cell Research* 21: 817-834.
- Markworth JF (2014) Ibuprofen treatment blunts early translational signaling responses in human skeletal muscle following resistance exercise. *Journal of Applied Physiology* 117: 20-28.
- Wyss M, Kaddurah-Daouk R (2000) Creatine and Creatinine Metabolism. *Physiological Reviews* 80: 1107-1213.
- Tarnopolsky MA (2010) Caffeine and creatine use in sport. *Ann Nutr Metab* 2: 1-8.
- Rawson ES (2011) Low-dose creatine supplementation enhances fatigue resistance in the absence of weight gain. *Nutrition* 27: 451-455.
- Fredholm BB (1999) Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Pharmacol Rev* 51: 83-133.
- Gobatto CA (2001) Maximal lactate steady state in rats submitted to swimming exercise. *Comp Biochem Physiol A Mol Integr Physiol* 130: 21-27.
- Cechella JL (2014) Moderate swimming exercise and caffeine supplementation reduce the levels of inflammatory cytokines without causing oxidative stress in tissues of middle-aged rats. *Amino Acids* 46: 1187-1195.