Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Opinion Article

# Cancer Stem Cell Plasticity Regulating Disease Progression and Therapeutic Resistance

**Sofia Reinhardt***

*Department of Experimental Oncology, Central European Institute of Biomedical Sciences, Vienna, Austria*

## DESCRIPTION

Cancer is a biologically diverse disease composed of multiple cellular populations displaying considerable variation in morphology, metabolism, proliferative activity, molecular characteristics, and responses to environmental signals. This diversity has encouraged continuous investigation into the mechanisms responsible for disease initiation, progression, recurrence, and therapeutic response. Among the most actively studied cellular populations are cancer stem cells, which possess the capacity for prolonged self-renewal, multilineage differentiation, and adaptation to changing biological environments. Rather than representing a fixed group of cells, these populations demonstrate remarkable plasticity, continuously altering their biological characteristics in response to metabolic conditions, extracellular signaling, tissue architecture, and therapeutic exposure. This dynamic behavior has established cancer stem cell plasticity as an important subject within cancer research and immuno-oncology because it influences tumor heterogeneity, metastatic dissemination, disease recurrence, and long-term clinical outcomes.

The concept of cancer stem cells originated from observations that only a relatively small fraction of malignant cells could efficiently regenerate new tumors following transplantation into experimental models. These findings suggested that malignant tissues contain specialized populations possessing greater proliferative and developmental capacity than surrounding cancer cells. Subsequent investigations demonstrated that these cells express distinct molecular markers, display unique metabolic programs, and occupy specialized microenvironments supporting long-term survival. Modern research, however, indicates that cellular identity remains highly dynamic, with differentiated malignant cells capable of reacquiring stem cell–like characteristics under appropriate biological conditions. This reversible transition illustrates the importance of cellular plasticity during disease progression.

Normal tissue stem cells maintain organ homeostasis by generating differentiated cells required for tissue renewal throughout life. Their biological activity is tightly regulated through interactions with surrounding stromal populations, extracellular matrix proteins, vascular structures, and biochemical signaling pathways. Comparable regulatory mechanisms operate within malignant tissues, although they become progressively altered during cancer development. Cancer stem cells respond continuously to environmental signals while modifying neighboring cellular populations through reciprocal communication. These interactions contribute to extensive biological diversity observed within individual malignancies.

The microenvironment surrounding cancer stem cells contains fibroblasts, endothelial cells, macrophages, dendritic cells, lymphocytes, adipocytes, extracellular matrix proteins, and soluble signaling molecules that collectively influence cellular behavior. Rather than functioning independently, these components establish highly coordinated communication networks regulating proliferation, metabolism, migration, differentiation, and survival. Variability in local environmental conditions contributes to regional differences in stem cell activity throughout malignant tissues, resulting in multiple cellular states existing simultaneously within the same lesion.

Fibroblasts represent one of the principal stromal populations influencing cancer stem cell plasticity. Under healthy physiological conditions, fibroblasts maintain connective tissue organization through production of collagen, laminins, fibronectin, elastin, and additional extracellular matrix components. During malignant progression, continuous exposure to signaling molecules released by cancer cells modifies fibroblast physiology, leading to altered secretion of cytokines, growth factors, extracellular vesicles, and matrix-remodeling proteins. These biological changes influence cellular differentiation and contribute to maintenance of stem cell–associated phenotypes within specialized tissue regions.

Endothelial cells forming blood vessels create specialized vascular niches supporting cancer stem cell maintenance. These microenvironments provide oxygen, nutrients, metabolic substrates, and signaling molecules while facilitating communication with additional stromal populations. Irregular

**Correspondence to:** Sofia Reinhardt, Department of Experimental Oncology, Central European Institute of Biomedical Sciences, Vienna, Austria, E-mail : sofia.reinhardt@ceibs-austria.edu

**Received:** 01-Jun-2026, Manuscript No. JCRIO-26-43993; **Editor assigned:** 03-Jun-2026, PreQC No. JCRIO-26-43993(PQ); **Reviewed:** 17-Jun-2026, QC No. JCRIO-26-43993; **Revised:** 24-Jun-2026, Manuscript No. JCRIO-26-43993 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2684-1266.26.12.293

**Citation:** Reinhardt S (2026). Cancer Stem Cell Plasticity Regulating Disease Progression and Therapeutic Resistance. J Cancer Res Immunooncol. 11:293.

**Copyright:** © 2026 Reinhardt S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

vascular organization characteristic of malignant tissues produces regional differences in oxygen concentration and nutrient availability, generating distinct biological environments. Cancer stem cells adapt to these fluctuating conditions by modifying metabolic pathways and intracellular signaling, demonstrating remarkable physiological flexibility.

Dendritic cells also influence the biological environment occupied by cancer stem cells through production of signaling molecules and interactions with lymphocyte populations. Variations in dendritic cell activity affect communication throughout malignant tissues and contribute to differences in local biological conditions. Although numerous molecular mechanisms remain under investigation, current evidence indicates that antigen-presenting cells participate in regulation of complex cellular networks extending beyond conventional inflammatory pathways.

Lymphocyte populations display considerable diversity within malignant tissues, including cytotoxic T lymphocytes, helper T cells, regulatory T cells, natural killer cells, B lymphocytes, and plasma cells. These populations interact continuously with stromal cells and malignant cells through direct contact, soluble mediators, and extracellular vesicles. Differences in lymphocyte composition among tissue regions contribute to variability in cellular communication influencing cancer stem cell physiology and adaptation.

Mitochondria perform essential functions supporting these metabolic adaptations. Beyond generating cellular energy, mitochondria regulate calcium homeostasis, lipid metabolism, reactive oxygen species production, and biosynthetic pathways required for cellular maintenance. Continuous remodeling through mitochondrial fusion, fission, intracellular trafficking, and quality-control mechanisms enables cancer stem cells to respond efficiently to environmental stress. Variability in mitochondrial function among individual stem cell populations further contributes to biological heterogeneity.

Autophagy provides an additional adaptive mechanism supporting long-term cellular survival. Through lysosomal degradation of damaged proteins, organelles, and intracellular components, autophagy maintains metabolic homeostasis during nutrient limitation and oxidative stress. Cancer stem cells frequently modify autophagic activity according to local physiological conditions, allowing efficient recycling of intracellular materials required for biosynthesis and energy production. This adaptive process illustrates the close relationship between intracellular quality control and cellular plasticity.

## CONCLUSION

Future progress in cancer research and immuno-oncology will continue to improve understanding of cancer stem cell plasticity through integration of single-cell technologies, spatial molecular profiling, metabolomics, extracellular vesicle analysis, artificial intelligence, and liquid biopsy. Dynamic interactions among malignant cells, fibroblasts, endothelial cells, macrophages, dendritic cells, lymphocytes, and extracellular matrix components create highly adaptable biological environments supporting continuous cellular change. Comprehensive investigation of these communication networks is expected to strengthen biomarker development, improve individualized patient assessment, and contribute to more effective therapeutic strategies across a broad spectrum of human malignancies.