

## Carbonic Anhydrase between Aerobic and Anaerobic Glycolysis

Saliyu I\*


Department of Biochemistry, Federal University Dutse, Nigeria

\*Corresponding author: Ibrahim Saliyu Ismail, Department of Biochemistry, Federal University Dutse, Nigeria, Tel: +2348036316108; E-mail: Saliyu.ibrahim@fud.edu.ng

Received date: December 05, 2016; Accepted date: December 12, 2016; Published date: December 16, 2016

Copyright: © 2016 Saliyu I. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Image Article



**Figure 1:** Carbonic anhydrase between aerobic and anaerobic glycolysis.

Carbonic anhydrase converts CO<sub>2</sub> and H<sub>2</sub>O generated from aerobic oxidation of glucose in the mitochondria to HCO<sub>3</sub><sup>-</sup> + H<sup>+</sup>, the HCO<sub>3</sub><sup>-</sup> is

either transported in the red blood cells for expiration via the lungs, serve as a chemical buffer or is transported into the liver to serve as a substrate for pyruvate carboxylase for gluconeogenesis. Under condition of exclusively anaerobic glycolysis glucose produces lactate as an end product. Lactate is a metabolic dead end and it has to be shuttled out of the cell to prevent intracellular lactate accumulation. Carbonic anhydrase facilitates lactate transport in and out of the cells through monocarboxylate transporters. It facilitates transport of lactate from muscle and red blood cells into the liver where it serves as a substrate for gluconeogenesis. Both exogenous and endogenous glucose metabolism result in the production of this two metabolic dead end products (CO<sub>2</sub> and lactate) which must be transported out of the cells to prevent intracellular accumulation, failure of which result in metabolic acidosis. Inhibition of carbonic anhydrase has long been found to cause metabolic acidosis. Saliyu's cycle provides a means of recycling these end products through 'carbonic anhydrase' to prevent intracellular accumulation and hence increase the amount of energy needs of the body through continues ATP production (Figure 1). The question is how does this glycolytic metabolic shift affects energy imbalance especially in diabetes and cancer?