Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Commentary

Cellular Senescence Signatures Associated with Cancer Evolution and Therapeutic Responses

Daniela Moretti*

Department of Translational Cancer Sciences, Mediterranean Biomedical University, Florence, Italy

DESCRIPTION

Cellular senescence has emerged as one of the most extensively investigated biological processes in modern cancer research because of its complex relationship with malignant transformation, tissue homeostasis, aging, and responses to therapy. Initially described as a permanent state of proliferative arrest occurring after repeated cellular division, senescence is now recognized as a highly dynamic biological program involving extensive metabolic adaptation, altered cellular communication, chromatin reorganization, mitochondrial remodeling, and secretion of numerous biologically active molecules. Rather than representing a passive endpoint of cellular aging, senescent cells remain metabolically active and continuously interact with neighboring stromal populations, vascular structures, extracellular matrix components, and inflammatory cells. These interactions influence tissue organization and contribute to variability observed during cancer progression and therapeutic intervention. Consequently, cellular senescence has become an important subject within cancer research and immuno-oncology because of its participation in multiple stages of malignant disease.

Healthy tissues generate senescent cells under several physiological circumstances. During embryonic development, temporary senescence contributes to normal tissue remodeling and organ formation. In adult life, senescence occurs following repeated cellular replication, oxidative stress, metabolic imbalance, telomere shortening, radiation exposure, and activation of oncogenic signaling pathways. By permanently limiting cellular proliferation, senescence serves as a protective biological mechanism that reduces expansion of damaged cells. Under normal conditions, these cells are gradually removed through coordinated interactions with surrounding stromal and inflammatory populations, allowing tissue homeostasis to be maintained.

Malignant transformation introduces additional complexity to senescence biology. Cancer cells frequently develop molecular alterations that allow continued proliferation despite signals that normally induce growth arrest. At the same time, surrounding fibroblasts, endothelial cells, epithelial cells, and additional stromal populations may undergo senescence after prolonged exposure to inflammatory mediators, oxidative stress, altered metabolism, or therapeutic intervention. Consequently, malignant tissues often contain mixtures of proliferating cells and senescent populations occupying different anatomical regions. This coexistence creates a biologically diverse microenvironment characterized by continuous cellular communication and structural remodeling.

One of the defining characteristics of senescent cells is the senescence-associated secretory phenotype. This biological program involves production of cytokines, chemokines, extracellular matrix proteins, growth factors, proteolytic enzymes, lipids, metabolites, and extracellular vesicles that influence neighboring cells. Through secretion of these molecules, senescent populations actively modify tissue architecture and regulate communication among malignant cells, fibroblasts, endothelial cells, macrophages, dendritic cells, lymphocytes, and additional stromal components. The composition of secreted factors varies according to tissue type, initiating stimulus, metabolic status, and disease stage, demonstrating that senescence is a heterogeneous rather than uniform biological process.

Dendritic cells similarly respond to signaling molecules released by senescent populations. Alterations in antigen uptake, migration, cytokine production, and communication with lymphocyte populations may occur following exposure to senescence-associated secretory factors. Such biological responses influence adaptive cellular activity and contribute to variability in inflammatory regulation among different malignancies. Continued investigation seeks to clarify relationships between senescence and antigen presentation across multiple forms of cancer.

Lymphocyte populations also interact extensively with senescent cells. Cytotoxic T lymphocytes and natural killer cells participate in elimination of senescent populations under healthy

Correspondence to: Daniela Moretti, Department of Translational Cancer Sciences, Mediterranean Biomedical University, Florence, Italy, E-mail: daniela.moretti@mbu-italy.edu

Received: 01-Jun-2026, Manuscript No. JCRIO-26-43988; **Editor assigned:** 03-Jun-2026, PreQC No. JCRIO-26-43988(PQ); **Reviewed:** 17-Jun-2026, QC No. JCRIO-26-43988; **Revised:** 24-Jun-2026, Manuscript No. JCRIO-26-43988 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2684-1266.26.12.288

Citation: Moretti D (2026). Cellular Senescence Signatures Associated with Cancer Evolution and Therapeutic Responses. J Cancer Res Immunooncol. 11:288.

Copyright: © 2026 Moretti D. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

physiological conditions. However, chronic accumulation of senescent cells within malignant tissues modifies communication with lymphocyte populations through continuous secretion of cytokines, chemokines, and extracellular vesicles. These interactions influence migration, metabolic adaptation, and functional activity of neighboring lymphocytes, contributing to regional differences in cellular composition observed throughout malignant lesions.

Spatial transcriptomics has further strengthened these investigations by preserving anatomical relationships while measuring molecular activity within intact tissues. Researchers can determine where senescent populations are located relative to malignant cells, blood vessels, connective tissue, and inflammatory infiltrates. Such analyses reveal that local environmental conditions strongly influence senescence-associated molecular programs. Regional variation within individual lesions contributes to substantial biological diversity and affects communication among neighboring cellular populations.

Artificial intelligence has become increasingly valuable for integrating molecular, imaging, histopathological, and clinical information related to cellular senescence. Machine learning algorithms identify complex molecular signatures associated with disease progression and therapeutic responsiveness while analyzing digital pathology specimens containing senescent stromal populations. These computational approaches support development of predictive models capable of improving individualized patient assessment.

CONCLUSION

Future progress in cancer research and immuno-oncology will increasingly depend upon comprehensive characterization of senescence-associated molecular programs across diverse cellular populations. Cellular senescence influences extracellular matrix remodeling, vascular biology, metabolism, intercellular communication, and tissue organization through continuous interactions with fibroblasts, endothelial cells, macrophages, dendritic cells, lymphocytes, and malignant cells. Continued integration of single-cell technologies, spatial molecular profiling, metabolomics, computational biology, and clinical investigation will improve understanding of senescence heterogeneity while supporting development of refined biomarkers and innovative therapeutic strategies for patients affected by a wide range of human malignancies.