OPEN ACCESS Freely available online

Journal of Cancer Research and Immuno-Oncology

Commentary

Cellular Senescence within the Tumor Microenvironment and its Influence on Cancer Oncology

Mirela Novak*

Department of Molecular Oncology, Danubia Institute of Medical Sciences, Zagreb, Croatia

DESCRIPTION

Cancer development involves continuous interactions among malignant cells, stromal components, immune populations, vascular structures, and extracellular signaling molecules. Rather than functioning as isolated cellular masses, tumors represent highly organized biological systems in which every cellular population contributes to disease behavior. One area receiving increasing scientific attention is cellular senescence, a biological condition characterized by permanent withdrawal from the cell division cycle while maintaining metabolic activity. Although senescence originally attracted attention because of its association with aging, current research demonstrates that senescent cells also influence tissue repair, developmental processes, chronic diseases, and malignant disorders. Within cancer immuno-oncology, these cells have become an important subject because they modify immune communication, extracellular signaling, and tissue architecture throughout disease progression.

Within malignant tissues, however, this clearance process frequently becomes less efficient. Senescent cells may accumulate in both cancerous and non-cancerous compartments of the tumor microenvironment. Fibroblasts, endothelial cells, epithelial cells, immune populations, and even certain malignant cells can enter senescence following exposure to biological stress. As these cells remain metabolically active, they continuously release cytokines, chemokines, growth factors, extracellular vesicles, enzymes, and matrix-modifying proteins that influence neighboring cellular populations. These secreted molecules collectively create a dynamic signaling environment capable of modifying tumor behavior through multiple biological pathways.

One characteristic frequently associated with senescent cells involves alterations in their secretory profile. These signaling molecules influence cellular communication throughout malignant tissues by affecting immune cell recruitment, inflammatory activity, vascular development, extracellular matrix composition, and metabolic regulation. The composition of these secreted factors varies according to tissue type, initiating stimulus, cellular origin, and disease stage. Consequently, senescent cells demonstrate considerable biological diversity rather than representing a uniform population.

The relationship between senescence and malignant transformation appears highly dependent upon timing and biological context. During the earliest phases of tumor development, senescence may restrict expansion of damaged cells by preventing further proliferation. This response reduces opportunities for accumulation of additional molecular abnormalities. As disease progresses, persistent accumulation of senescent cells may gradually alter the surrounding microenvironment through continuous secretion of biologically active molecules. These changes may influence immune regulation, tissue remodeling, and interactions between malignant cells and surrounding stromal populations.

Natural killer cells are particularly important during recognition of senescent cells because they identify stress-associated molecules expressed on altered cellular surfaces. Successful interaction promotes removal of senescent populations and contributes to maintenance of tissue stability. Within malignant tissues, however, continuous exposure to suppressive signaling molecules may reduce natural killer cell activity. Such alterations permit prolonged survival of senescent cells, increasing their influence on surrounding biological processes.

Fibroblasts undergoing senescence contribute additional changes to the extracellular matrix. Healthy fibroblasts normally maintain structural integrity by producing collagen, elastin, and other matrix proteins. Senescent fibroblasts gradually alter matrix composition through secretion of remodeling enzymes and inflammatory mediators. These modifications influence tissue stiffness, cellular migration, vascular organization, and immune cell infiltration. Altered matrix architecture also affects diffusion of therapeutic agents within tumors, creating regional variation in treatment exposure.

Cancer therapy itself frequently induces senescence in both malignant and surrounding normal cells. Radiation therapy, cytotoxic chemotherapy, targeted agents, and certain biological treatments generate sufficient cellular stress to initiate

Correspondence to: Mirela Novak, Department of Molecular Oncology, Danubia Institute of Medical Sciences, Zagreb, Croatia, E-mail: mirela.novak@dims-croatia.edu

Received: 27-Feb-2026, Manuscript No. JCRIO-26-43978; **Editor assigned:** 02-Mar-2026, PreQC No. JCRIO-26-43978(PQ); **Reviewed:** 16-Mar-2026, QC No. JCRIO-26-43978; **Revised:** 23-Mar-2026, Manuscript No. JCRIO-26-43978 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2684-1266.26.12.277

Citation: Novak M (2026). Cellular Senescence within the Tumor Microenvironment and its Influence on Cancer Oncology. J Cancer Res Immunooncol. 11:277.

Copyright: © 2026 Novak M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

permanent cell cycle arrest. Therapy-induced senescence may contribute to temporary disease control by preventing additional malignant cell division. Nevertheless, persistence of senescent populations following treatment introduces additional biological complexity because these cells continue releasing signaling molecules capable of modifying the tumor microenvironment long after completion of therapy.

This observation has generated interest in distinguishing beneficial from unfavorable effects of therapy-induced senescence. Short-term senescence accompanied by effective immune-mediated clearance may contribute to improved disease control. Conversely, prolonged persistence of senescent cells may support chronic inflammatory activity, tissue remodeling, and altered immune communication. Clinical investigators continue examining factors that determine whether therapy-induced senescence produces favorable or unfavorable biological outcomes across different malignancies.

Several experimental therapeutic approaches seek to modify senescent cell populations. Some investigations evaluate compounds that selectively eliminate senescent cells after completion of conventional therapy, whereas others attempt to alter their secretory activity without affecting cell survival. Additional studies focus on improving immune-mediated clearance of senescent populations through modulation of natural killer cells or macrophage function. These approaches remain under active investigation across multiple preclinical and clinical settings.

Digital pathology combined with artificial intelligence provides additional opportunities to evaluate senescence within routine tissue specimens. Machine learning algorithms can analyze microscopic features associated with senescent morphology while integrating molecular staining patterns and spatial distribution. These computational approaches may improve identification of biologically significant senescent populations and assist pathologists in evaluating tissue heterogeneity across different malignancies.

CONCLUSION

Future progress in cancer immuno-oncology will likely depend upon comprehensive understanding of how senescent cells interact with malignant cells, immune populations, fibroblasts, endothelial cells, and extracellular matrix components. Rather than functioning as isolated biological entities, senescent cells participate in extensive communication networks that influence disease evolution through multiple interconnected mechanisms. Continued investigation into these interactions may contribute to improved biomarker development, refined therapeutic strategies, and enhanced understanding of individual differences observed among patients receiving cancer treatment. The expanding knowledge surrounding cellular senescence demonstrates that permanent cell cycle arrest represents only one aspect of a much broader biological process influencing immune regulation, tissue organization, therapeutic response, and long-term clinical outcomes across diverse forms of cancer.