

Cereblon as a Predictive Biomarker for Imid Therapy Sensitivity

Adel Gouri^{2*}, Amel Bouacha¹, Aoulia Dekaken³, Faiza Mehiddine⁴, Ahmed Aimen Bentorki², Amina Yakhlef⁵ and Mehdi Beleilli⁶

¹Department of Hematology, IBN ROCHD Public Hospital, Souk Ahras 41000, Algeria

²Laboratory of Medical Biochemistry, IBN ZOHR Public Hospital, Guelma 24000, Algeria

³Department of Internal Medicine; Pr EL OKBI Public Hospital, Guelma 24000, Algeria

⁴Department of Pneumology, IBN ZOHR Public Hospital, Guelma 24000, Algeria

⁵Department of Hematology, IBN ZOHR Public Hospital, Guelma 24000, Algeria

⁶Laboratory of Pharmaceutical Chemistry, College of Medicine, Annaba 2300, Algeria

*Corresponding author: Adel Gouri, Laboratory of Medical Biochemistry, IBN ZOHR Public Hospital, Guelma 24000, Algeria, Tel: 00213 666 088 ; E-mail: pharmagor@gmail.com

Received date: Apr 22, 2014, Accepted date: Apr 24, 2014, Published date: Apr 28, 2014

Citation: Gouri A, Bouacha A, Dekaken A, Mehiddine F, Bentorki AA, et al. (2014) Cereblon as a Predictive Biomarker for Imid Therapy Sensitivity. J Hematol Thrombo Dis 2: e112. doi: 10.4172/2329-8790.1000e112.

Copyright: © 2014 Gouri A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

Immunomodulatory drugs (IMiDs) are highly active in the treatment of hematological malignancies, including multiple myeloma (MM), non-Hodgkin lymphoma and chronic lymphocytic leukaemia, but the mechanisms of action are still incompletely understood. In multiple myeloma, myelodysplastic syndrome and lymphomas, thalidomide and/or the structurally analogous compound lenalidomide, and pomalidomide; are antiproliferative, increase immune surveillance and decrease stromal cell support.

Although several mechanisms have been proposed to explain the activity of these drugs in multiple myeloma (MM), the precise cellular targets and molecular mechanisms have only recently become clear. Cereblon (CRBN) a member of the cullin 4 ring ligase complex (CRL4) has been identified as the primary target of thalidomide teratogenicity and moreover, as an essential requirement for response to IMiD therapy [1]. CRBN is also implicated in several effects of IMiDs, such as down-regulation of tumor necrosis factor- α (TNF- α) and T cell immunomodulatory activity [2].

Cereblon's central role as a target of lenalidomide and pomalidomide suggests potential utility as a predictive biomarker of response or resistance to IMiD therapy; Low CRBN expression was found to correlate with poor drug response in MM cell lines and primary MM [3].

Recently, Steven et al. demonstrated a significant correlation between low CRBN expression and decreased survival outcomes in homogenously cohort of MM patients treated with single agent pomalidomide and dexamethasone [4].

More recently, Huang et al. suggested that expression of CRBN protein in myeloma cells assessed using immunohistochemistry is a practical approach to predict the response of IMiDs in MM patients [5]. CRBN expression is therefore a potential predictive biomarker for response and survival in MM, decreases expression developed resistance to IMiDs. In addition, understanding the CRBN-dependent basis of the anti-myeloma effects of IMiDs may lead to development of novel agents with antitumor activity distinct from the teratogenic effects associated with currently available IMiD [6].

The discovery of cereblon as a target of thalidomide and the IMiDs has enabled a pursuit of better understanding of the molecular

mechanism of action by these agents. In parallel, there is significant clinical interest to investigate the possible role of cereblon as a biomarker for the IMiD compounds response or resistance. Nevertheless, measurement of CRBN protein is associated with a number of assay limitations. The requirement for high-quality clinical samples, such as myeloma cells enrichment by cell sorting, limits the validation of such quantified transcriptional expression of the CRBN gene method to every MM patient. In addition, the lack of a consensus protocol to amplify the CRBN gene is a crucial problem.

The positive predictive value of high CRBN expression is less robust and the interpretation of CRBN expression is complicated by the presence of multiple CRBN isoforms.

Therefore, further analysis is required to validate the putative predictive effect of CRBN for IMiDs sensitivity and standardized assays for measuring CRBN expression accurately are more than needed.

References

1. Schuster SR, Kortuem KM, Zhu YX, Braggio E, Shi CX, et al. (2014) The clinical significance of cereblon expression in multiple myeloma. *Leuk Res* 38: 23-28.
2. Lopez-Girona A, Mendy D, Ito T, Miller K, Gandhi AK, et al. (2012) Cereblon is a direct protein target for immunomodulatory and antiproliferative activities of lenalidomide and pomalidomide. *Leukemia* 26: 2326-2335.
3. Broyl A, Kuiper R, van Duin M, van der Holt B, el Jarari L, et al. (2013) High cereblon expression is associated with better survival in patients with newly diagnosed multiple myeloma treated with thalidomide maintenance. *Blood* 121: 624-627.
4. Schuster SR, Kortuem KM, Zhu YX, Braggio E, Shi CX, et al. (2014) The clinical significance of cereblon expression in multiple myeloma. *Leuk Res* 38: 23-28.
5. Huang SY, Lin CW, Lin HH, Yao M, Tang JL, et al. (2014) Expression of cereblon protein assessed by immunohistochemical staining in myeloma cells is associated with superior response of thalidomide- and lenalidomide-based treatment, but not bortezomib-based treatment, in patients with multiple myeloma. *Ann Hematol* 14: 2063-2067.
6. Gandhi AK, Mendy D, Waldman M, Chen G, Rychak E, et al. (2014) Measuring cereblon as a biomarker of response or resistance to lenalidomide and pomalidomide requires use of standardized reagents and understanding of gene complexity. *Br J Haematol* 164: 233-244.