

Clinical Importance of Chirality in Drug Design and Pharmaceuticals

Rose RK*

Department of Chemistry, PSL University, CNRS, Institut de Recherche Chimie Paris (IRCP), Paris, France

DESCRIPTION

Chiral molecules are a class of organic molecules that possess chirality, which is a property of handedness or asymmetry. In simple terms, chiral molecules are non-superimposable mirror images of each other, much like a left and right hand. These two mirror-image forms are called enantiomers. Chirality arises from the spatial arrangement of atoms around a central carbon atom or a chiral center (chiral carbon). The two enantiomers of a chiral drug can have different pharmacological activities and biological effects in the human body, despite having the same chemical formula. Examples of chiral drugs include:

- Ibuprofen: The S-enantiomer is the active form responsible for pain relief.
- Albuterol: The R-enantiomer is the active form used to treat asthma.
- L-DOPA(L-3,4-Dihydroxyphenylalanine): The L-enantiomer is used to treat Parkinson's disease.

Steps involved in drug design

Selective target binding: Chiral molecules often interact with biological targets (e.g., enzymes, receptors) in a highly selective manner. The binding affinity and specificity of each enantiomer to its target can vary significantly. Drug designers aim to develop compounds with high affinity for the target and minimal off-target interactions.

Stereochemistry: The three-dimensional arrangement of atoms in a chiral molecule can affect its ability to bind to a biological target. The differences in stereochemistry can result in vastly different biological activities. Drug designers use computational modeling and structure-activity relationships to improve stereochemistry for target binding.

Regulatory approval: Regulatory agencies, such as the FDA and EMA, require thorough characterization of chiral drugs, including testing the safety and efficacy of each enantiomer separately. This process is essential for obtaining regulatory approval for a drug.

Chiral resolution: Drug design often involves the separation and purification of enantiomers to obtain the desired pharmacologically

active form. Techniques like chiral chromatography and asymmetric synthesis are used to prepare single enantiomers.

Patent protection: Pharmaceutical companies may seek patents for the use of a specific enantiomer of a drug to protect their intellectual property and market exclusivity.

Personalized medicine: Chirality allows for personalized medicine approaches, customizing drug therapy to an individual's metabolism and responsiveness to specific enantiomers. This can optimize treatment outcomes and minimize adverse effects.

Pharmacological activity

The role of chirality in drug design is important, because many drugs are chiral molecules, meaning they exist as enantiomers, which are mirror-image forms of each other. Chirality can have a profound impact on a drug's pharmacological activity, efficacy, safety, and metabolism. Enantiomers of a chiral drug can have different pharmacological activities. One enantiomer may be therapeutically active, while the other may be inactive or exhibit different biological effects. Understanding the activity of each enantiomer is significant in designing drugs with the desired therapeutic effect.

Metabolism

The human body can metabolize the two enantiomers of a drug differently. One enantiomer may be metabolized more rapidly or converted into different metabolites compared to the other. Understanding metabolic pathways helps design drugs with desired pharmacokinetic properties. One enantiomer of a chiral drug may be associated with greater toxicity or adverse side effects compared to the other. Drug designers work to minimize the potential for unwanted side effects by selecting or designing the less toxic enantiomer.

CONCLUSION

Chirality is a critical consideration in drug design because it can profoundly influence a drug's pharmacological properties, safety, and efficacy. Understanding the stereochemistry and biological activity of enantiomers is essential for the rational design of

Correspondence to: Rose RK, Department of Chemistry, PSL University, CNRS, Institut de Recherche Chimie Paris (IRCP), Paris, France, E-mail: roserk978@sl.eu

Received: 18-Jul-2023, Manuscript No. OCCR-23-27158; **Editor assigned:** 21-Jul-2023, Pre QC No. OCCR-23-27158 (PQ); **Reviewed:** 04-Aug-2023, QC No. OCCR-23-27158; **Revised:** 11-Aug-2023, Manuscript No. OCCR-23-27158 (R); **Published:** 18-Aug-2023, DOI: 10.35841/2161-0401.23.12.332.

Citation: Rose RK (2023) Clinical Importance of Chirality in Drug Design and Pharmaceuticals. *Organic Chem Curr Res.* 12:332.

Copyright: © 2023 Rose RK. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

pharmaceuticals with the desired therapeutic effects and minimal side effects. Chiral molecules have significant implications in fields like chemistry, pharmacology, and biochemistry. In drug development, for example, understanding the chirality of active pharmaceutical ingredients is important to make sure that the

desired therapeutic effect is achieved while minimizing potential side effects associated with the inactive enantiomer. Chiral molecules are essential in the study of molecular symmetry and the origin of homochirality in biological systems.