

Comparative Genomic Characteristics of Lichen Forming Fungi across Diverse Climatic Regions

Serena Volkov*

Department of Environmental Mycology, Eastland Scientific University, Brno, Czech Republic

DESCRIPTION

Lichens are complex biological associations involving fungi and photosynthetic partners such as algae or cyanobacteria. Although the visible structure of a lichen appears to be a single organism, it is actually a consortium of multiple species that interact closely throughout their life cycles. Within these associations, the fungal partner contributes the majority of the physical structure and plays a major role in determining growth form, habitat preference, and ecological performance. Recent developments in fungal genomics have provided opportunities to investigate how lichen-forming fungi have evolved and adapted to environments that range from humid tropical forests to arid deserts and polar landscapes.

Lichen-forming fungi belong predominantly to the phylum Ascomycota, although some Basidiomycota species also participate in lichen associations. For many years, investigations relied primarily on morphology and limited molecular markers to classify these organisms. Whole-genome sequencing has transformed understanding of their diversity by revealing patterns of genetic variation that were previously difficult to detect. Comparative analyses have demonstrated that lichen-forming fungi possess genomic characteristics that distinguish them from closely related non-lichenized species.

One notable feature involves the genetic basis of symbiotic interaction. Lichen-forming fungi maintain long-term associations with photosynthetic organisms that provide carbon compounds through photosynthesis. Genomic studies have identified specialized gene families associated with cellular communication, nutrient exchange, and recognition mechanisms. These genes contribute to the establishment and maintenance of stable symbiotic relationships. Their distribution varies among species occupying different climatic regions, suggesting adaptation to local environmental conditions and partner availability.

The ability of lichens to colonize exposed and nutrient-poor habitats has attracted considerable scientific interest. Many species thrive on rock surfaces, tree bark, soil crusts, and other challenging substrates where few organisms can persist.

Comparative genomic investigations indicate that lichen-forming fungi possess diverse metabolic capabilities that support survival under such conditions. Genes involved in nutrient scavenging, mineral interaction, and organic compound utilization frequently exhibit expansion compared with related fungi lacking symbiotic lifestyles.

Secondary metabolites constitute another prominent aspect of lichen biology. Lichens synthesize numerous chemical compounds that influence ecological interactions and environmental tolerance. Many of these substances possess antimicrobial, antioxidant, or protective properties. Genomic investigations have identified extensive collections of biosynthetic gene clusters responsible for metabolite production. Species from different climatic regions frequently exhibit distinct biosynthetic repertoires, reflecting adaptation to local ecological pressures.

Solar radiation presents an important challenge for exposed lichen communities. Ultraviolet radiation can damage cellular structures and genetic material. Comparative genomic analyses indicate that fungi inhabiting high-elevation or low-latitude environments often possess enhanced genetic capacity for protective pigment production and Deoxyribonucleic Acid (DNA) repair. These features contribute to survival under conditions characterized by elevated radiation exposure.

Reproductive strategies also influence genomic diversity. Some lichen-forming fungi reproduce primarily through sexual spores, while others rely heavily on vegetative propagules containing both fungal and photosynthetic partners. Comparative analyses suggest that reproductive mode affects patterns of genetic variation, recombination, and population connectivity. Species emphasizing sexual reproduction generally exhibit greater opportunities for genetic reshuffling and adaptation to changing environmental conditions.

Chromosomal organization among lichen-forming fungi displays considerable diversity. Genome sequencing has revealed variation in chromosome number, repetitive DNA content, and structural arrangement. Mobile genetic elements contribute to genomic flexibility by generating mutations and rearrangements.

Correspondence to: Serena Volkov, Department of Environmental Mycology, Eastland Scientific University, Brno, Czech Republic, E-mail: serena.volkov.efg@esu-czmail.org

Received: 27-Feb-2026, Manuscript No. FGB-26-42477; **Editor assigned:** 02-Mar-2026, PreQC No. FGB-26-42477 (PQ); **Reviewed:** 16-Mar-2026, QC No. FGB-26-42477; **Revised:** 23-Mar-2026, Manuscript No. FGB-26-42477 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2165-8056.26.16.303

Citation: Volkov S (2026). Comparative Genomic Characteristics of Lichen Forming Fungi across Diverse Climatic Regions. Fung Genom Biol. 16:303.

Copyright: © 2026 Volkov S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Although once considered relatively static, lichen fungal genomes are now recognized as dynamic systems capable of evolutionary change over time.

Comparisons with non-lichenized relatives have provided insights into the origins of lichen symbiosis. Certain gene families appear reduced or absent in lichen-forming species, whereas others have expanded significantly. These patterns suggest long-term specialization associated with symbiotic lifestyles. Evolutionary analyses indicate that multiple independent origins of lichenization have occurred throughout fungal history, leading to diverse genomic solutions for similar ecological challenges.

Transcriptomic studies complement genome sequencing by examining gene activity under specific environmental conditions. Analyses of lichen fungi exposed to dehydration, temperature stress, or altered light conditions reveal dynamic changes in expression patterns. These responses provide insight into the molecular mechanisms supporting survival across diverse habitats.

The availability of high-quality genome assemblies continues to improve rapidly. Long-read sequencing technologies and advanced computational methods now permit more detailed examination of repetitive regions, chromosome structure, and evolutionary history. As additional species are sequenced,

comparative analyses become increasingly informative, allowing broader assessment of genomic diversity among lichen-forming fungi.

Research involving lichen fungal genomics contributes to multiple scientific disciplines, including ecology, evolution, conservation biology, and biotechnology. Understanding how these organisms persist in challenging environments may provide insights relevant to stress tolerance, natural product discovery, and ecosystem function. Their capacity to inhabit environments ranging from tropical forests to polar rock surfaces reflects a long history of genetic adaptation and symbiotic specialization.

CONCLUSION

The comparative study of lichen-forming fungal genomes reveals a remarkable diversity of evolutionary strategies shaped by climate, substrate, symbiotic interaction, and environmental pressure. Through analyses of gene content, genome organization, metabolic potential, and population structure, researchers continue to expand understanding of how these unique fungi have adapted to some of the most varied habitats on Earth. As genomic resources increase, future investigations will further clarify the biological processes that contribute to the success and persistence of lichen-forming fungi.