Immunogenetics: Open Access

OPEN ACCESS Freely available online

Perspective

Complement System Diversity and Its Contribution to Host Immune Protection

Natalie Sørensen*

Department of Immunogenetics and Molecular Medicine, Northern Biomedical Research Institute, University of Copenhagen, Copenhagen, Denmark

DESCRIPTION

The human immune system relies on several interconnected defense mechanisms that work together to identify and eliminate harmful microorganisms while preserving healthy tissues. One of the earliest protective responses is provided by the complement system, which consists of numerous circulating proteins that become activated in a carefully coordinated sequence after infection or tissue injury. This system assists in destroying microorganisms, enhancing the activity of immune cells, removing damaged cells, and supporting communication between different components of immunity. Although the complement system functions in every healthy individual, inherited variation within complement system genes contributes to considerable differences in immune performance, susceptibility to infectious diseases, inflammatory conditions, and autoimmune disorders. The study of these inherited differences has become an important area of immunogenetics because it explains why individuals often respond differently to similar biological challenges.

The complement system includes more than thirty proteins produced primarily by the liver, although several other tissues also contribute to their production. These proteins circulate in an inactive state until they encounter signals indicating infection, damaged tissue, or immune complexes. Once activated, they participate in a sequence of carefully regulated reactions that amplify immune activity while limiting unnecessary injury to healthy cells. Every stage of this process depends upon proteins encoded by inherited genes, making genetic variation an important determinant of complement system efficiency.

Activation of the complement system occurs through three principal pathways. The classical pathway begins when antibodies bind to foreign material and activate complement proteins. The lectin pathway starts after specialized recognition molecules attach to carbohydrate structures commonly found on the surfaces of microorganisms. The alternative pathway remains continuously available at a low level of activity and becomes rapidly amplified when microorganisms are detected. Although each pathway begins differently, all three eventually produce common biological effects, including attraction of immune cells, enhancement of microbial destruction, and removal of cellular debris.

One of the most thoroughly investigated complement proteins is complement component three. This protein occupies a central position because all activation pathways eventually converge upon its activation. Once activated, complement component three produces fragments that attach to microbial surfaces, making harmful organisms easier for immune cells to recognize and remove. Additional fragments stimulate inflammatory responses by attracting protective immune cells toward sites of infection. Individuals carrying inherited variants that influence complement component three activity may display measurable differences in resistance to bacterial, viral, or fungal infections.

Complement component five also plays an important biological role following activation of the complement cascade. Activation produces fragments that recruit inflammatory cells and contribute to destruction of microorganisms through formation of membrane attack complexes. These complexes create openings within microbial membranes, allowing destruction of susceptible organisms. Inherited variation affecting complement component five or proteins participating in membrane attack complex formation may influence the efficiency of this protective mechanism.

Regulation of complement activity is equally important because uncontrolled activation may damage healthy tissues. Numerous regulatory proteins limit complement activation by preventing unnecessary amplification and protecting normal cells from accidental injury. Genes encoding these regulatory proteins demonstrate inherited diversity that contributes to differences in immune regulation. Reduced activity of regulatory proteins may permit excessive complement activation, increasing susceptibility to inflammatory diseases, whereas excessive regulation may weaken protection against infection.

Complement factor H represents one of the major regulatory

Correspondence to: Natalie Sørensen, Department of Immunogenetics and Molecular Medicine, Northern Biomedical Research Institute, University of Copenhagen, Copenhagen, Denmark, E-mail: natalie.sorensen@nbri-copenhagen.dk

Received: 24-Feb-2026, Manuscript No. IGOA-26-43864; **Editor assigned:** 26-Feb-2026, PreQC No. IGOA-26-43864(PQ); **Reviewed:** 12-Mar-2026, QC No. IGOA-26-43864; **Revised:** 19-Mar-2026, Manuscript No. IGOA-26-43864 (R); **Published:** 26-Mar-2026, DOI: 10.35248/IGOA.26.11.293

Citation: Sørensen N (2026). Complement System Diversity and Its Contribution to Host Immune Protection. Immunogenet Open Access. 11:293.

Copyright: © 2026 Sørensen N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

proteins responsible for controlling activation of the alternative pathway. This protein distinguishes healthy human cells from microorganisms by recognizing specific molecular patterns located on cellular surfaces. Once recognition occurs, complement factor H suppresses complement activation on healthy tissues while allowing continued activity against invading microorganisms. Inherited variation affecting complement factor H has been associated with several inflammatory and degenerative disorders involving the kidneys, eyes, and blood vessels. These observations demonstrate how inherited regulation contributes to maintenance of immune balance.

Complement factor I provides another important regulatory function by breaking down activated complement proteins after their protective role has been completed. Without adequate regulation by complement factor I, excessive complement activation may continue beyond the initial immune response, increasing tissue injury. Individuals carrying inherited variants affecting complement factor I activity may experience altered susceptibility to inflammatory disorders depending upon the nature of the inherited change.

The complement system works closely with antibody-mediated immunity. Antibodies identify foreign structures with remarkable specificity, while complement proteins strengthen the effectiveness of antibody binding by promoting destruction of microorganisms and improving recognition by immune cells. Because inherited variation influences both antibody production and complement activity, overall immune performance depends upon coordinated interaction between these biological systems rather than any single inherited characteristic.

CONCLUSION

Future investigation in immunogenetics will continue integrating genomic sequencing, protein analysis, cellular biology, computational modeling, and population research to improve understanding of inherited complement regulation. Continued international collaboration will expand available genomic resources while supporting investigation of rare inherited variants that cannot be adequately studied within individual populations alone. As scientific knowledge continues to increase, inherited variation within complement system genes will remain an important area of immunogenetics because it provides valuable insight into individual differences in immune protection, inflammatory regulation, susceptibility to infection, autoimmune disease development, and overall maintenance of immune homeostasis.