

## Cpap Pressure by Cpap Titration or Prediction Formula

Pattarapong Makarawate, Vichai Senthong and Kittisak Sawanyawisuth\*

Department of Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand

### Case Report

Obstructive sleep apnea (OSA) is a common disease in clinical practice and can lead to several cardiovascular complications such as death, heart failure, stroke, or atrial fibrillation [1]. The estimated prevalence is about 5-10% of general population [2]. Successful treatment with continuous positive airway pressure machine (CPAP) significantly reduces cardiovascular events [2,3]. Appropriate CPAP pressure can be determined by two methods: in hospital and home CPAP titration [4].

There are advantages and disadvantages of both methods. In hospital CPAP titration needs sleep technicians and hospitalization. Expenses and time consuming are two main disadvantages. In contrast, home automatic CPAP titration is more available, more convenient, less expensive, and less time consuming. Some issues on home CPAP titration are existed including high leakage of CPAP therapy or other troubleshooting problems on CPAP therapy while using CPAP [4,5]. Split night polysomnography or half night polysomnography with half night CPAP titration is an alternative way to save costs and times for in hospital CPAP titration. However, this modality may be appropriated only in severe OSA. To know the appropriate CPAP pressure is the goal of CPAP titration. In resource limited or busy sleep facilities, CPAP treatment while waiting for a titration test may be justified. Prediction of CPAP pressure by using a predictive formula is beneficial and equivalent to the in hospital standard method [6,7].

A CPAP prediction formula for American patients by Loredo et al. was published in 2007 [8]. Factors in the formula included respiratory disturbance, nadir saturation, and mean saturation. CPAP (pressure in cmH<sub>2</sub>O) = 30.8 + RDI × 0.03 - nadir saturation × 0.05 - mean saturation × 0.2 [8]. In 2012, a report from Turkey showed that CPAP pressure prediction can be calculated by using neck circumference and oxygen desaturation index [9]. Later on, Lee et al. reported a CPAP prediction model for Asian OSA patients [10]. The predictive equation was: predicted pressure (cm H<sub>2</sub>O) = 6.656 + 0.156 × (body mass index [kg/m<sup>2</sup>]) - 0.071 × (minimal SpO<sub>2</sub> [%]) + 0.041 × (respiratory disturbance index) + 0.094 × (score of Epworth Sleepiness Scale). Note that ESS was not in the US model.

The CPAP prediction model may be worthwhile but may need to be validated in different study population (Table 1). Ethnicity may be an important factor that affects the CPAP prediction model.

There are two types of CPAP machines for OSA treatment; auto

CPAP and fixed pressure CPAP machine. Both types have pros and cons. The auto CPAP is more comfortable and provides therapy with variable range of CPAP pressure. Using auto CPAP machine therefore may not need the in hospital CPAP titration. The 90<sup>th</sup> or 95<sup>th</sup> percentile of CPAP pressure is given. A meta-analysis showed that the auto CPAP may have better outcomes on patient preference, compliance, and sleep architecture [11]. Some disadvantages of the auto CPAP include higher cost and limited clinical data [12,13]. The effect of auto CPAP on long-term cardiovascular outcomes needs further studies. One report showed that the auto CPAP may have residual apnea events compared with in hospital CPAP titration [14]. In resource limited setting, treatment of OSA with fixed CPAP machine using predictive model to identify the CPAP pressure may be a justified option. This strategy may reduce cost and facilitate faster treatment of OSA.

### Conclusion

In conclusion, CPAP pressure prediction is beneficial particularly in resource-limited facilities. The formula may need to be validated in one's own setting.

### References

- Gottlieb DJ, Yenokyan G, Newman AB, O'Connor GT, Punjabi NM, et al. (2010) Prospective study of obstructive sleep apnea and incident coronary heart disease and heart failure: the sleep heart health study. *Circulation* 122: 352-360.
- Young T, Palta M, Dempsey J, Peppard PE, Nieto FJ, et al. (2009) Burden of sleep apnea: rationale, design, and major findings of the Wisconsin Sleep Cohort study. *WMJ* 108: 246-249.
- Punjabi NM, Caffo BS, Goodwin JL, Gottlieb DJ, Newman AB, et al. (2009) Sleep-disordered breathing and mortality: a prospective cohort study. *PLoS Med* 6: e1000132.
- Masa JF, Jiménez A, Durán J, Capote F, Monasterio C, et al. (2004) Alternative methods of titrating continuous positive airway pressure: a large multicenter study. *Am J Respir Crit Care Med* 170: 1218-1224.
- Yu CC, Hua CC, Tseng JC, Liu YC (2006) Comparison optimal pressure between automatic titrating and predicting continuous positive airway pressure. *Chang Gung Med J* 29: 583-589.
- Hertegonne KB, Volna J, Portier S, De Pauw R, Van Maele G, et al. (2008) Titration procedures for nasal CPAP: automatic CPAP or prediction formula? *Sleep Med* 9: 732-738.
- Lopez-Campos JL, Garcia Polo C, Leon Jimenez A, Gonzalez-Moya E, Arnedillo A, et al. (2007) CPAP titration: Different methods for similar clinical results. *Eur J Intern Med* 18: 230-234.

| Model                    | Population/Year | Formula   |
|--------------------------|-----------------|---|
| Loredo et al. [8]        | US/2007         | 30.8 + RDI × 0.03 - nadir saturation × 0.05 - mean saturation × 0.2   |
| Basoglu and Tasbakan [9] | Turkey/2012     | (0.148 × NC) + (0.038 × ODI)  |
| Lee et al. [10]          | Korea/2013      | 6.656 + 0.156 × (body mass index [kg/m <sup>2</sup> ]) - 0.071 × (minimal SpO <sub>2</sub> [%]) + 0.041 × (respiratory disturbance index) + 0.094 × (score of Epworth Sleepiness Scale) |

Note: RDI: Respiratory Disturbance Index; NC: Neck Circumference; ODI: Oxygen Desaturation Index

**Table 1:** Summary of CPAP pressure prediction model in three studies.

\*Corresponding author: Dr. Kittisak Sawanyawisuth, Srinagarind hospital, Department of Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand, Tel: 66-43-363664; Fax: 66-43-348399; E-mail: [kittisak@kku.ac.th](mailto:kittisak@kku.ac.th)

Received May 21, 2013; Accepted June 15, 2013; Published June 20, 2013

**Citation:** Makarawate P, Senthong V, Sawanyawisuth K (2013) Cpap Pressure by Cpap Titration or Prediction Formula. *J Sleep Disorders Ther* 2: 125. doi:10.4172/2167-0277.1000125

**Copyright:** © 2013 Makarawate P, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

8. Loredó JS, Berry C, Nelesen RA, Dimsdale JE (2007) Prediction of continuous positive airway pressure in obstructive sleep apnea. *Sleep Breath* 11: 45-51.
9. Basoglu OK, Tasbakan MS (2012) Determination of new prediction formula for nasal continuous positive airway pressure in Turkish patients with obstructive sleep apnea syndrome. *Sleep Breath* 16: 1121-1127.
10. Lee GH, Kim MJ, Lee EM, Kim CS, Lee SA (2013) Prediction of Optimal CPAP Pressure and Validation of an Equation for Asian Patients With Obstructive Sleep Apnea. *Respir Care* 58: 810-815.
11. Xu T, Li T, Wei D, Feng Y, Xian L, et al. (2012) Effect of automatic versus fixed continuous positive airway pressure for the treatment of obstructive sleep apnea: an up-to-date meta-analysis. *Sleep Breath* 16: 1017-1026.
12. Vennelle M, White S, Riha RL, Mackay TW, Engleman HM, et al. (2010) Randomized controlled trial of variable-pressure versus fixed-pressure continuous positive airway pressure (CPAP) treatment for patients with obstructive sleep apnea/hypopnea syndrome (OSAHS). *Sleep* 33: 267-271.
13. Patruno V, Aiolfi S, Costantino G, Murgio R, Selmi C, et al. (2007) Fixed and autoadjusting continuous positive airway pressure treatments are not similar in reducing cardiovascular risk factors in patients with obstructive sleep apnea. *Chest* 131: 1393-1399.
14. Denotti AL, Wong KK, Dungan GC 2nd, Gilholme JW, Marshall NS, et al. (2012) Residual sleep-disordered breathing during autotitrating continuous positive airway pressure therapy. *Eur Respir J* 39: 1391-1397.