

Current Trends in Sporotrichosis

Roberto Christ Vianna Santos^{1*}, Rodrigo de Almeida Vaucher¹ and Sydney Hartz Alves²

¹Microbiology Laboratory, Health Sciences, Franciscan University Center – UNIFRA, Brazil

²Mycological Research Laboratory, Department of Microbiology and Parasitology, Universidade Federal de Santa Maria (UFSM), Santa Maria, Rio Grande do Sul, Brazil

Sporotrichosis is a sub-acute or chronic infection that affects the dermis and subcutaneous tissue, with a worldwide distribution, being more common in tropical and subtropical areas. In South America, the incidence is 48 to 60 cases per 100,000 inhabitants [1]. The classification schemes of several clinically important fungi have suffered major modifications in recent decades, due to advancement of molecular techniques, which allows easier access to sequences of DNA. However, in 2006, modifications occurred in the Sporothrix classification. [2] Marimon has suggested that *S. schenckii* could not be considered the only species, causing sporotrichosis. They described four new species of Sporothrix complex: *S. globosa*, *S. brasiliensis*, *S. Mexicana*, and *S. luriei*, formerly *S. schenckii* var. *luriei*.

Currently, the available data seems to indicate that the new species have different geographic distributions. *S. globosa* has a worldwide distribution [3], *S. brasiliensis* related epidemics of sporotrichoses in Rio de Janeiro, Brazil [4] and *S. Mexicana* appears limited to Mexican territory [2]. Only the correct identification of these new species will allow us to address the question, that the different clinical manifestations of Sporotrichosis, are produced by different species. Until now, the different species composing the complex have different susceptibility rates to antifungal drugs [5,6].

Molecular studies have demonstrated the existence of a high level of intraspecific variability, in which isolates are mainly grouped according to their geographical origin established [7-9], on the basis of Internal Transcribed Spacer (ITS) region sequence analysis, that more than one species could exist, within *S. schenckii*.

Several researches on the genomic DNA base composition rendered an average guanine and cytosine content, of about 54.7 mol% [10]. *S. schenckii* presents six to eight chromosomes of 460 to 6,200 kb, at a total genome size of about 28 Mbp [11]. Maybe such differences regard to either the different species recently described [2], or to underestimation in the methods adopted for genome size determination. It has also been reported that *S. schenckii*, a diploid organism, bearing around 50 fg DNA per cell, in both the filamentous and yeast phases. Aneuploidy, a state in which most of the chromosomes are disomic, cannot be excluded though [12]. Diploidy is considered as essential for thermal dimorphism in *Cryptococcus neoformans*, and similarities in life cycle between this fungus and other dimorphic fungi, including *S. schenckii*, might happen [13].

Mitochondrial DNA (mtDNA) analysis of Restriction Length Polymorphisms (RFLP) with the restriction enzyme Hae III, is usually the basis for studies on identification, typing, and epidemiology of sporotrichosis. It has been initially cited 24 mtDNA types [14], and more recently, 25 to 30 [15] and 31 to 32 [16] were introduced. Such analyses have been adopted in several studies, with *S. schenckii* strains from different geographical origins and also environmental isolates [17,18] being deposited to date in Gene Bank, nucleotide sequences of these fungi 914 being the most observed species *S. schenckii*, followed by *S. brasiliensis*, *S. globosa*, *S. mexicana*, *S. Schenkivarluriei* among others, respectively.

References

- da Rosa AC, Scroferneker ML, Vettorato R, Gervini RL, Vettorato G, et al. (2005) Epidemiology of sporotrichosis: a study of 304 cases in Brazil. J Am Acad Dermatol 52: 451-459.
- Marimon R, Cano J, Gené J, Sutton DA, Kawasaki M, et al. (2007) Sporothrix brasiliensis, *S. globosa*, and *S. mexicana*, three new Sporothrix species of clinical interest. J Clin Microbiol 45: 3198-3206.
- Madrid H, Cano J, Gené J, Bonifaz A, Toriello C, et al. (2009) Sporothrix globosa, a pathogenic fungus with widespread geographical distribution. Rev Iberoam Micol 26: 218-222.
- Oliveira, M. M. E. 2009. M.Sc. thesis. Fundação Oswaldo Cruz Rio de Janeiro, Brazil.
- Marimon R, Serena C, Gené J, Cano J, Guarro J. (2008) In vitro antifungal susceptibilities of five species of Sporothrix. Antimicrob Agents Chemother 52: 732-734
- Barros,MB, de Almeida Paes R, Schubach AO (2011) Sporothrix schenckii and Sporotrichosis. Clin Microbiol Rev 24: 633-654.
- Ishizaki H (2003) Mitochondrial DNA analysis of Sporothrix schenckii. Nihon Ishinkin Gakkai Zasshi 44: 155-157.
- Ishizaki H, Kawasaki M, Aoki M, Matsumoto T, Padhye AA, et al. (1998) Mitochondrial DNA analysis of Sporothrix schenckii in North and South America. Mycopathologia 142: 115-118.
- de Beer ZW, Harrington TC, Vismar HF, Wingfield BD, Wingfield MJ (2003) Phylogeny of the Ophiostoma stenoceras-Sporothrix schenckii complex. Mycologia 95: 434-441.
- Mendonça-Hagler LC, Travassos LR, Lloyd KO, Phaff HJ (1974) Deoxyribonucleic acid base composition and hybridization studies on the human pathogen Sporothrix schenckii and Ceratocystis species. Infect Immun 9: 934-938.
- Tateishi T, Murayama SY, Otsuka F, Yamaguchi H (1996) Karyotyping by PFGE of clinical isolates of Sporothrix schenckii. FEMS Immunol Med Microbiol 13: 147-154.
- Torres-Guerrero H (1999) Ploidy study in sporothrix schenckii. Fungal Genet Biol 27: 49-54.
- Sia RA, Lengeler KB, Heitman J (2000) Diploid strains of the pathogenic basidiomycete Cryptococcus neoformans are thermally dimorphic. Fungal Genet Biol 29: 153-163.
- Lin J, Kawasaki M, Aoki M, Ishizaki H, You G, et al. (1999) Mitochondrial DNA analysis of Sporothrix schenckii clinical isolates from china. Mycopathologia 148: 69-72.
- Mora-Cabrera M, Alonso RA, Ulloa-Arvizu R, Torres-Guerrero H (2001) Analysis of restriction profiles of mitochondrial DNA from Sporothrix schenckii. Med Mycol 39: 439-444.

*Corresponding author: Roberto Christ Vianna Santos, Microbiology Laboratory, Health Sciences Center Universitário Franciscano - UNIFRA, Brazil, Tel: +55 55 3220-1200; E-mail: robertochrist@gmail.com

Received September 14, 2012; Accepted September 16, 2012; Published September 23, 2012

Citation: Santos RCV, de Almeida Vaucher R, Alves SH (2012) Current Trends in Sporotrichosis. Fungal Genom Biol 2:e110. doi:10.4172/2165-8056.1000e110

Copyright: © 2012 Santos RCV, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

-
16. Ishizaki H, Kawasaki M, Anzawa K, Mochizuki T, Chakarabarti A, et al. (2009) Mitochondrial DNA analysis of *Sporothrix schenckii* in India, Thailand, Brazil, Colombia, Guatemala and Mexico. *Nihon Ishinkin Gakkai Zasshi* 50:19-26.
17. Arenas R, Miller D, Campos-Macias P (2007) Epidemiological data and molecular characterization (mtDNA) of *Sporothrix schenckii* in 13 cases from Mexico. *Int J Dermatol* 46: 177-179.
18. Xu TH, Lin JP, Gao XH, Wei H, Liao W, et al. (2010) Identification of *Sporothrix schenckii* of various mtDNA types by nested PCR assay. *Med Mycol* 48: 161-165.