

Depression among orphan children residing in orphanages in Nepal: a cross-sectional study

Bhatt Kumari Bandana

Ministry of Health and Population, Nepal



Abstract

Background: Children living in orphanages are vulnerable groups of the population separated from the care and love of their parents. Lack of parental support during the early years of growth and development can lead to develop and manifest different levels of depression. This study aimed to determine the prevalence of depression and its associated factors among orphan children residing in orphanages in Nepal.

Methods: An institutional-based cross-sectional study was conducted in 20 orphanages of Nepal. The orphanages were selected by simple random sampling method and 13-17 years children were invited to participate in the study. A structured self-administered questionnaire and Standard Beck Depression Inventory-III (BDI-II) tools were used for data collection whereas logistic regression was used for identifying associated factors with depression at the significant level $\alpha = 0.05$.

Results: A total of 322 orphans participated in the study; 55.4% were males, the average age was 14.7 years. Most of the participants (91.3%) studied in secondary level and most (32.8%) belonged from indigenous ethnic group. The overall prevalence of depression among orphans was 32.2%, with higher preponderance among the female population (39.2%). In the multivariate model, sex, health problems, social support, and being bullied were significantly associated with depression in orphans. Females were 2.30 times more likely to develop depression than males (95% CI=1.38-3.82). Those who had health problems were 2.08 times more likely to develop depression than those who did not (95% CI=1.21-3.57). Those who had low social support were 2.48 times more likely to develop depressive symptoms than those who had high social support (95% CI=1.24-4.95), and those who had been bullied were 2.59 times more likely to develop depressive symptoms than those who were not bullied (95% CI=1.46-4.59).

Conclusion: This study showed a high prevalence of depression among children living in the orphanages in Nepal particularly among girls. Therefore, measures should be taken to identify, prevent and eliminate any practices within orphanages that may be reproducing discriminatory and stigmatizing behaviors that

exist toward girl's which research has shown to impact adversely on the mental health and wellbeing of children and which can manifest in depression.



Biography:

Kumari Bandana Bhatt is a Senior Public Health officer in the Department of Health Services under Ministry of Health and Population of Nepal. In her role, she is responsible for the planning, preparedness and management of epidemic/outbreak, vector borne and Neglected Tropical Diseases as well as planning of the program and budget. She is also involved in researches associated with public health issues. Moreover, providing trainings, their monitoring and supervision is also the major role in her job.

She has earned the degree of the Master of Science in Public Health (Epidemiology) and the Master of Public Administration (MPA). She has conducted seven independent study; 3 of them have been published in international Journals.

[9th International Conference on Epidemiology & Public Health](#); Webinar- September 24-25, 2020.

Abstract Citation:

Bhatt Kumari Bandana, Depression among orphan children residing in orphanages in Nepal: a cross-sectional study, Epidemiology Summit 2020, 9th International Conference on Epidemiology & Public Health; Webinar- September 24-25, 2020.

(<https://publichealth-epidemiology.conferenceseries.com/abstract/2020/depression-among-orphan-children-residing-in-orphanages-in-nepal-a-cross-sectional-study>)