

Research Article

Description of *Myxidium tetraodoni* Sp. Nov., *Myxidium anisocapsularis* Sp. Nov. and *Myxobolus magai* Sp. Nov. (Myxosporea: Bivalvulida) Infecting Some Freshwater Fishes in Cameroon (Central Africa)

Deli Arnaud, Lekeufack Folefack Guy Benoit^{*} and Fomena Abraham

Department of Animal Biology and Physiology, Laboratory of Parasitology and Ecology, University of Yaounde I, Cameroon

*Corresponding author: Lekeufack Folefack Guy Benoit, Department of Animal Biology and Physiology, Laboratory of Parasitology and Ecology, University of Yaounde I, Cameroon, Tel: +00237677887294; E-mail: leguyzo@yahoo.fr

Received date: November 21, 2017; Accepted date: December 06, 2017; Published date: December 13, 2017

Copyright: © 2017 Deli A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Examination of some Teleost fishes captured in the Maga detention lake located in the Far North Region of Cameroon, revealed the presence of three new species of Myxosporidia of the genera *Myxidium* Bütschli, 1882 and *Myxobolus* Bütschli, 1882 of which complete description is given in the present study. These species are: *Myxidium tetraodoni* sp. nov., parasite of the urinary bladder of *Tetraodon lineatus* Linnaeus, 1758 (Tetraodontidae) that form ellipsoidal spores with a turgid middle part and rounded ends. They measured 11.6 (10.5-12.5) µm long × 8.2 (7.2-9.6) µm broad; the spherical polar capsules are of equal size and measure 3.7 (3.0-4.3) µm. *Myxidium anisocapsularis* sp. nov., a parasite of the gall bladder of *Distichodus engycephalus* Günther, 1964 (Distichodontidae) form spindle-shaped and elongated spores, that measure 15.2 (14.0-16.2) µm long × 6.0 (5.0-6.7) µm broad; its polar capsules are quite unequal and respectively measure 6.0 (5.0-6.5) × 3.3 (3.0-3.8) µm for the smaller. *Myxobolus magai* sp. nov., a gill parasite of *Labeo batesii* Boulenger, 1911 (Cyprinidae) form ovoid spores with the anterior end larger with small protuberance, that measure 10.6 (9.0-12.0) × 6.3 (5.5-7.0) µm. Its polar capsules measure 2.8 (2.4-3.4) × 2.3 (2.0-3.0) µm.

Keywords Myxozoa; *Myxidium tetraodoni* sp. nov.; *Myxidium anisocapsularis* sp. nov.; *Myxobolus magai* sp. nov.; Parasite; Freshwater fish; Cameroon

Introduction

Due to the increasing needs for animal protein production worldwide, fish industry is a major activity in human populations today [1]. However, fishes are vulnerable to various parasitic infections, including Myxosporidiosis. Myxosporidia can infect numerous fish organs and cause a wide variety of damages [2]. The extent of damage to host tissues and organs depend on the parasite species, its life cycle, infection intensity, and host response [3]. Many pathogenic species can be responsible of severe epizootics in fish farms and hatcheries. Indeed, in natural and artificial environments, Myxosporidia can cause the loss of production and death of fishes. Some parasitized fishes are unsightly and considered unfit for human consumption. Human health can be affected by Myxosporidia when immunodepressed people are contaminated with infected fish [4].

Knowledge of fish parasites is a prerequisite for a rapid diagnosis of the pathogen responsible for epizootic diseases. Early diagnosis can lead to preventive measures that remain the best way to reduce epidemics [5]. Thus it is necessary to record and report new parasites and pathological conditions when they are discovered because such information may be useful in the future as a baseline data for assessing the ecosystems health in the face of global warming [6]. Based on the morphological and metric characteristics of the spores, the number of Myxosporidia described around the world is estimated at 2200 species belonging to 64 genera and 17 families [7]. Eiras et al. [8,9] provided a synopsis of 905 species of *Myxobolus* Bütschli [10]; this genus has the largest global distribution and the high number of Myxosporidia species. With more than 232 species, the genus *Myxidium* Bütschli [10] is numerically the second genus of Myxosporidia [11]. Species of the genus *Myxidium* are predominantly coelozoic (rarely histozoic) infecting the gall bladder, the urinary bladder and the urinary tubules of the kidneys [2,12,13]. The number of Myxosporidia parasites of African freshwater fish is estimated at about 270 species [14,15]. Those of the freshwater fishes in Cameroon are represented by about 80 species, with forty of them belonging to the genus *Myxobolus* and nine to the genus *Myxidium* [14,16,17]. Taxonomic studies and description of new species of Myxosporidia throughout the world suggested that the diversity of species in this group of parasites is largely underestimated.

During a study of Myxosporidia parasites of teleosts of great importance of food and economic in Cameroon, we found three new species whose complete descriptions are given in the present paper. These species are: *Myxidium tetraodoni* sp. nov. a parasite of *Tetraodon lineatus* Linnaeus, 1758 (Tetraodontidae), *Myxidium anisocapsularis* sp. nov. found in *Distichodus engycephalus* Günther, 1964 (Distichodontidae) and *Myxobolus magai* sp. nov. a gill parasite of *Labeo batesii* Boulenger, 1911 (Cyprinidae).

Page 2 of 10



Figure 1: Microphotograph (optical microscopy) of a Trophozoite of *Myxidium tetraodoni* sp. nov. with a granular endoplasm, the spores are formed in pairs within a pansporoblast. Scale bar: $5 \,\mu$ m.



Figure 3: Microphotograph (optical microscopy) of spores of *Myxidium tetraodoni* sp. nov. stained with May-Grünwald-Giemsa. Scale bar: 5 μm.



Figure 2: Microphotograph (optical microscopy) of fresh spore of *Myxidium tetraodoni* sp. nov. : polar capsules are spherical. Scale bar: $5 \mu m$.



Figure 4: Microphotograph (optical microscopy) of fresh spore of *Myxidium anisocapsularis* sp. nov. : polar capsules are unequal. Scale bar: 5 µm.



Figure 5: Microphotograph (optical microscopy) of fresh spores of *Myxidium anisocapsularis* sp. nov. showing unequal polar capsules. Scale bar: $5 \mu m$.



Figure 6: Microphotograph (optical microscopy) of spore of *Myxidium anisocapsularis* sp. nov. stained with May-Grünwald-Giemsa. Scale bar: $5 \mu m$.

Material and Methods

Fish examined were sampled in the Maga detention lake from March to May 2016. Maga is a small village of the Mayo-Danay Division (situated between latitude 10° and 13° North and longitude 14° and 16° East), in the Far North Region of Cameroon. The Maga reservoir lake is located at 77 km from the city of Maroua and covers an area of 39000 hectares. The climate in this locality is the Sahelo-Sudanian type, which is characterized by a long dry season from October to April and a short rainy season from May to September. The average temperature is 28°C [18]. Maga is located in a floodplain and its vegetation consists of the shrub savannah (weakly wooded) and the herbaceous steppe with grass. Its soil, predominantly clay-sandy, take a clayey-silty shade around the lake [18].

Fish were captured using gill nets. In the field, once the fish were caught, a buttonhole was made on the ventral surface of each host individual. The latter were immediately immersed in a 10% formalin solution contained in a plastic container, before being brought to the laboratory. Systematic position of sampled fishes was taken from Lévêque et al. [19,20]. The species captured are: *Tetraodon lineatus* (Tetraodontidae), *Distichodus engycephalus* (Distichodontidae) and *Labeo batesii* (Cyprinidae).



Figure 7: Microphotograph (optical microscopy) of fresh spores of *Myxobolus magai* sp. nov.; The spores are ovoid with a wider anterior end. Scale bar: $5 \mu m$.



Figure 8: Microphotograph (optical microscopy) of fresh spore of *Myxobolus magai* sp. nov.; The anterior end of the spore is hillock. Scale bar: $5 \mu m$.



Figure 9: Microphotograph (optical microscopy) of spores of *Myxobolus magai* sp. nov. stained with May-Grünwald-Giemsa; polar capsules are equal. Scale bar: $5 \mu m$.

In the laboratory, the fish were firstly examined with naked eye (eyes, fins, operculum, scales, skin) and then with the Olympus BO61 binocular stereoscopic microscope for the presence of cysts. After dissection of the fish hosts, internal organs such as gills, heart, liver, kidneys, spleen, gallbladder, gonads, intestine and urethra were taken individually and examined. The contents of the cysts were identified with the objective 100X of an IVYMEN light microscope. The content of the gall bladder, urinary bladder and swim bladder was also examined between glass and glass cover slide under the microscope. Smears of kidneys, spleen, liver, gonads, heart and urethra were carefully examined at the 40X objective of the microscope. Spore smears were fixed with methanol and stained with May-Grünwald-Giemsa. Drawings of fresh spores were performed using a Wild M-20 microscope equipped with a camera Lucida. Measurements were carried out on 50 spores using an objective micrometer. Variables taken into account are those proposed by Lom et al. [21]. Microphotographs of spores were performed using an Olympus BH-2 microscope equipped with a microphotograph device.





Results

Myxidium tetraodoni sp. nov.

Vegetative form: Circular and polysporous plasmodia are found in the urinary bladder of the host. Ectoplasm is clear and smooth. In the granular endoplasm, spores are formed in pairs within pansporoblasts (Figure 1).

Spores: Of small size (11.6 μ m length on average), spores are ellipsoidal with a bulging middle part and rounded ends (Figures 2-11). Shell valves striated, one can count 9 to 12 fine striations on each (Figure 11). Polar capsules are spherical and equal (Figures 3 and 11). Within each of them, 4 to 5 coils of polar filament can be found (Figure 10). A granular sporoplasm is located between polar capsules.

Measurements of the spore: (Table 1).

Host: Tetraodon lineatus Linnaeus, 1758 (Tetraodontidae).

Page 5 of 10

Location: Maga (in the reservoir lake) in Cameroon (Central Africa).

Location in the host: Urinary bladder

Prevalence: 76.9% (20 individuals fish parasitized out of 26 examined).

Myxidium species	Host	Site of infection	Country	LS	ws	PC	LPC	WPC	FC	StV	Reference
M. tetraodoni	Tetraodon lineatus	urinary bladder	Cameroon	11.6 (10.5– 12.5)	8.2 (7.2– 9.6)	=	3.7 (3– 4.3)	3.7 (3– 4.3)	4–5	9–12	Present study
M.brienomyri	Brienomyrus brachyistus	gall bladder	Cameroon	13.7 (12.2– 16.2)	6.5 (5.5–9)	=	4.2 (3.5–5)	4.2 (3.5–5)	3–5	6–12	Fomena et al. [22]
M. latesi	Lates niloticus	gall bladder	Chad	15.4 (15– 16)	8.3 (8– 9)	=	3.3 (3– 3.5)	3.3 (3– 3.5)	-	±	Kostoïngué et al. [23]
M. macrocapsulare	Scardinius erythrophthalmus Grunnian Aplodinotus	gall bladder	Europe America	10–12	6	=	3–4	3–4	-	±	Auerbach [24]
M. macrocheili	Catostomus macrocheilus	gall bladder	North America	11.7 (10.0– 14.4)	6.6 (5.5– 8.0)	=	4.0 (3.0– 5.5)	3.5 (2– 4.5)	4–6	9–11	Mitchell [26]
M. moxostomatis	Myxostoma sp.	gall bladder	USA	8.5–10.5	5–6	=	3	3	-	10	Kudo [25]
M. nyongensis	Barbus jae, B. aspilus, B. guirali, B. martorelli	gall bladder	Cameroon	12.3 (10.8– 14.4)	6.4 (4.7– 9.4)	=	3.2 (2.0– 4.5)	3.2 (2.0– 4.5)	-	12	Fomena et al. [22]
Averages of the paran	neters measured are followe	d by minimal and r	maximal values i	in brackets.							

LS: length of the spore; WS: width of the spore; PC: relative length of the polar capsules (=: equal); LPC: length of the polar capsules; WPC: width of the polar capsules; FC: number of polar filament coils; StV: Number of striations on shell valves;-: absent or not reported in the species description; ±: shell valves striations are present, but the number was not reported in the species description.

Table 1: Comparative description of Myxidium tetraodoni sp. nov. with morphologically similar species (measurements in micrometre).

Myxidium anisocapsularis sp. nov.

Vegetative form: Not observed; spores, sometimes abundant, were found free in the bile (Figure 12).

Spores: Of medium size (15.2 μ m long on average), they are spindleshaped, elongated (2.5 times longer than wide), with a turgid medial part and acuminate extremities (Figures 4 and 5). Each valve carries 8 to 11 longitudinal striations (Figures 5 and 13). The suture line is straight. The polar capsules are piriform and quite unequal (Figures 6 and 13). The larger polar capsule contains 5 to 6 loops of the filament and the smaller 4 to 5 (Figure 12). A finely granular sporoplasm is located in the extra-capsular space.

Measurements of the spore: (Table 2).

Host: Distichodus engycephalus Günther, 1964 (Distichodontidae).

Location: Maga (in the reservoir lake) in Cameroon (Central Africa).

Location in the host: Gall bladder.

Prevalence: 15.38% (04 parasitized fish out of 26 examined).

Myxidium species	Host	Site of infection	Country	LS	ws	PC	LPC	WPC	FC	StV	Reference
M. anisocapsularis	Distichodus engycephalus	gall bladder	Cameroon	15.2 (14.0– 16.2)	6.0 (5.0–	¥	6 (5.0– 6.5)*	3.3 (3.0– 3.8)*	5–6*	8–11	Present study
				10.2)	0.7)		4.7 (4.0– 5.5)**	3. (2.3– 3.3)**	4–5**		
M. aydai	Caesio suevicus	gall bladder	Egypt	23.0 (22.0– 24.0)	5.6 (5.0– 6.0)	=	7.2 (7.0– 8.0)	3.4 (3.0 – 4.0)	8–9	-	Abdel-Baki [29]

Page 6 of 10

M. birgii	Aphyosemiom bivittatum	gall bladder	Cameroon	20.0 (17.7– 22.5)	8.9 (7– 11)	=	8.1 (7.1– 9.1)	3.7 (2.5– 4.8)	7–9	8–14	Fomena et al. [22]
M. camerounensis	Neolebias ansorgei	gall bladder	Cameroon	22 (19.2– 25.5)	6.4 (5– 8.4)	=	8.9 (7– 10.9)	3.4 (2.8 – 3.9)	7–9	6–10	Fomena et al. [22]
M. distichodi	Distichodus engycephalus	gall bladder	Chad	16.3 (16– 17)	6.5 (6– 7)	=	4.9 (4.5– 5.5)	3.2 (3– 3.5)	-	±	Kostoïngué et al. [23]
M. mendehi	Barbus guirali, B. martorelli	Kidneys	Cameroon	9.9 (7.8– 13.2)	4.1 (3.1– 4.9)	=	3.4 (2.7– 4.5)	2.3 (1.8 – 3.1)	5–7	5	Fomena et al. [28]
M. parachannae	Parachanna obscura	gall bladder	Benin	23.4 (21– 25)	4.3 (3– 5)	=	5.8 (5–7)	2.2 (1.5– 3)	-	-	Sakiti [27]
M. petrocephali	Petrocephalus simus	gall bladder	Cameroon	24.1 (21.5–27)	8 (6.5– 9.8)	=	10.3 (8.5 –11.6)	4.3 (3.8– 5)	13	12	Fomena et al. [22]
Averages of the parameters measured are followed by minimal and maximal values in brackets.											
LS: length of the spore; WS: width of the spore; PC: relative length of the polar capsules (=: equal; ≠: unequal); LPC: length of the polar capsules; WPC: width of the polar capsules; FC: number of polar filament coils; StV: Number of striations on shell valves; –: absent or not reported in the species description; ±: shell valves striations are present, but the number was not reported in the species description;											

*:Referring to the largest polar capsule;

**:Referring to the smallest polar capsules.

Table 2: Comparative description of Myxidium anisocapsularis sp. nov. with morphologically similar species (measurements in micrometre).

Myxobolus magai sp. nov.

Vegetative form: This Myxosporidia forms whitish and ovoid cysts, located between the secondary gill lamellae. These plasmodia are polysporous and measure $172-300 \times 116 \mu$ m-200 μ m. one can find 10 to 132 cysts per individual fish host (Figures 2-11).

Spores: of small size (10.6 μ m long on average), mature spores are ovoid with the anterior end larger and hillocky, the posterior end being narrow (Figures 7, 8, 14 and 15). The wider part of the spore is observed at the mid length of the polar capsules. Polar capsules are ovoid and symmetrical (Figures 7 and 8). They occupy the 1/3 of the spore cavity, converge and open at the anterior end of the spore (Figure

8). Each of them contains 4 to 5 coils of polar filament (Figure 14). The iodinophilous vacuole is found in a developped sporoplasm (Figures 7 and 15).

Measurements of the spore: (Table 3).

Host: Labeo batesii Boulenger, 1911 (Cyprinidae).

Location: Maga (in the reservoir) in Cameroon (Central Africa).

Location in the host: gills.

Prevalence: 28.5% (04 parasitized fish out of 14 examined).

	1	1		1					1		1
species	Host	Site of infection	Country	LS	ws	PC	LPC	WPC	FC	IP	Reference
Myxobolus magai	Labeo batesii	Gills	Cameroon	10.6 (9.0– 12.0)	6.3 (5.5– 7.0)	=	2.8 (2.4– 3.4)	2.3 (2.0– 3.0)	4–5	A	Present study
Myxobolus cichlidarum	Oreochromis niloticus ; Sarotherodon galilaeus	Gills, fins, eye, kidneys, spleen, liver	chad	15.0 (13.5–16)	9.6 (8.5– 10.8)	=	5.8 (5.2– 6.5)	3,2 (2,8– 3,5)	4–5	Р	Abakar- ousman [31]
Myxobolus eirasi	Cirrhina mrigala	caudal fin	India	8.4–8.8 (8.6)	6.5–6.9 (6.7	=	3.1–3.3 (3.2)	1.4–1.7 (1.5)	3–4	A	Kaur et al. [32]
Myxobolus galilaeus	Sarotherodon galilaeus	Kidneys, spleen	Israel	11.9 (10.3– 13.1)	9.1 (7.9– 10.0)	=	3.5 (3.1– 4.0)	2.8 (2.3– 3.1)	4–5	A	Landsberg [30]
Myxobolus gandiolensis	Tilapia guineensis	kidneys	Senegal	11.3 (10– 12)	10.3 (9– 12)	=	3.8 (3-5)	3.8 (3– 5)	-	A	Fall et al. [33]
Myxobolus nounensis	Sarotherodon galilaeus, Tilapia mariae	Spleen, kidneys	Cameroon	14.3 (13– 15)	12.8 (11.5–14)	=	5.8 (5– 6.5)	4.5 (4– 5)	4–5	Р	Fomena et al. [35]

Page 7 of 10

Myxobolus sarigi	Oreochromis aureus x O. niloticus	Spleen	Israel	11.3 (9.9 –13.1)	8.4 (7.9– 9.6)	=	4.5 (4.1– 5.2)	3.2 (2.9– 4.0)	4–5	A	Landsberg [30]
Myxobolus yogendrai	Cirrhina mrigala	under scales	India	9–9.5	7.2	=	2.8–3.6	2.8–3.6	-	A	Landsberg et al. [34]
Averages of the parameters measured are followed by minimal and maximal values in brackets.											
LS: length of the spore; WS: width of the spore; PC: relative length of the polar capsules (=: equal); LPC: length of the polar capsules; WPC: width of the polar capsules; FC, number of polar filament coils (-: not reported in the species description); IP: intercapsular process (A: absent; P: present).											

Table 3: Comparative description of Myxobolus magai sp. nov. with morphologically similar species (measurements in micrometre).

Discussion

Myxidium tetraodoni sp. nov.

By the spore shape, this myxosporidia come close to some species previously described in Africa and other continents. *Myxidium brienomyri* Fomena et al. [22], develops large trophozoites in the gall bladder of *Brienomyrus brachyistus* (Mormyridae) in Cameroon. Spores of this species are longer (13.7 μ m on average) and less wide (6.47 μ m on average); although spherical, the polar capsules are more developed (4.2 μ m diameter on average).

Myxidium nyongensis Fomena et al. [22], develop circular and large trophozoites in the gall bladder of *Barbus jae*, *B. aspilus*, *B. guirali* and *B. martorelli* (Cyprinidae) in Cameroon. Its spores are ellipsoidal with rounded ends but longer (12.38 μ m on average) and less wide (6.47 μ m on average) compared to those of the species in description.

The spores of *Myxidium latesi Kostoïngué*, Faye et al. [23] (host: *Lates niloticus* in Chad) are oval, with pointed ends. They are considerably more developed (15.44 μ m long × 8.33 μ m wide) compared to those of the present species.

Myxidium macrocapsulare Auerbach [24] (host: *Scardinius erythrophthalmus* in Europe and *Grunnian Aplodinotus* in America) forms free spores in the gall bladder with slightly curved ends pointed in the opposite directions. These spores are narrower (5.3 μ m-6.8 μ m) with polar capsules rather ovoid (4.6 μ m × 3.8 μ m on average).

The shape of spores of *Myxidium moxostomatis* Kudo [25], recalls that of the species we are describing; however, this parasite of the gall bladder of *Myxostoma* sp. in America forms much smaller spores (8.5 μ m-10.5 μ m × 5-6 μ m).

The spores of *Myxidium macrocheili* Mitchell [26] (host: *Catostomus macrocheilus* in North America) have comparable length with those of our parasite (11.7 μ m long on average); however, they are less wide (6.6 μ m on average) with ovoid polar capsules which measure 4 × 3.5 μ m on average.

The parasite of *Tetraodon lineatus*, which differ from other species by numerous characteristics, is probably new. We propose the name *Myxidium tetraodoni* sp. nov. refering to the generic name of the fish host. This is the first time a Myxosporidia is described in *Tetraodon lineatus*.



Fish Aqua J, an open access journal ISSN:2150-3508

Figure 13: Line drawing of a spore of *Myxidium anisocapsularis* sp. nov. showing longitudinally striated shell valves.

Myxidium anisocapsularis sp. nov.

The only *Myxidium* species so far described in African Distichodontidae is *Myxidium distichodi* Kostoïngué, [23], a parasite of the gall bladder of *Distichodus engycephalus* in Chad. The spores dimensions of *M. distichodi* (16.3 (16-17) μ m × 6.5 (6-7) μ m) are comparable to those of the species being described but their polar capsules are equal (constant character).

The general form of spores of the species in description recalls that of *Myxidium parachannae* Sakiti [27], parasite of the gall bladder of *Parachanna obscura* in Benin. However, the spores of *M. parachannae* are much longer ($21 \mu m$ - $25 \mu m$) and less wider ($3 \mu m$ - $5 \mu m$).

Myxidium birgii Fomena et al. [22], found in the gall bladder of *Aphyosemiom bivittatum* (Cyprinodontidae) in Cameroon form fusiform spores, with striated valves and a bulging medial part, but longer (17.7 μ m-22.5 μ m long). The spores shape of *Myxidium mendehi* Fomena et al. [28], recalls that of the species we are describing; however, this parasite of the kidneys of *Barbus guirali* and *B. martorelli* (Cyprinidae) form smaller spores (9.9 μ m × 4.1 μ m in average).

The spores of *Myxidium camerounensis* Fomena et al. [22], although fusiform with a bulging medial part, are longer (22.04 μ m on average), with equal polar capsules containing 7 to 9 loops of polar filament.

The spores of *Myxidium petrocephali* Fomena et al. [22] (host: *Petrocephalus simus* in Cameroon) are slightly arched, more developed (24.14 μ m × 8.05 μ m on average) with symmetrical and larger polar capsules (10.3 μ m × 4.3 μ m on average).

Myxidium aydai Abdel-Baki [29], parasite of *Caesio suevicus* in Egypt produces spores measuring 23 μ m long on average with symmetrical polar capsules measuring 7.2 μ m on average.

Out of all Myxosporidian species of the genus *Myxidium* described so far in the world, no one form spores with unequal polar capsules. We believe that the parasite of *Distichodus engycephalus* is new and propose the name *Myxidium anisocapsularis* sp. nov., refering to the asymmetry observed on its polar capsules.





Myxobolus magai sp. nov.

By the spore's general shape, the present parasite resemble some species of Myxosporidia previously described.

Myxobolus galilaeus Landsberg [30], forms diffuse spores in melano-macrophage centers of the kidneys and spleen of *Sarotherodon galilaeus* in Israel. These spores are ovoid with the anterior end flattened. They are larger (11.9 μ m × 9.1 μ m on average) and 3 to 12 folds are found on the suture line. *Myxobolus sarigi* Landsberg [30], a systemic parasite of Cichlidae in Israel, forms spores with the anterior end wider. These spores are wider (8.4 μ m on average) with larger polar capsules (4.5 μ m × 3.2 μ m on average).

Myxobolus cichlidarum Abakar-ousman [31], parasitizes fins, gills, eyes, spleen, kidneys and liver of Oreochromis niloticus and

Page 9 of 10

Sarotherodon galilaeus in Chad. Although ovoid with a wider anterior end, the spores of this species, differ from those of our species by the following characteristics: larger size (15 μ m × 9.6 μ m on average), more developed polar capsules (5.8 μ m × 3.2 μ m on average), presence of an intercapsular appendix.

The spores of *Myxobolus eirasi* Kaur et al. [32] (parasite of *Cirrhina mrigala* in India), are morphologically comparable to those of our parasite, but shorter ($8.4 \mu m$ - $8.8 \mu m$) with smaller polar capsules ($1.4 \mu m$ - $1.75 \mu m$).

Fall et al. [33] described *Myxobolus gandiolensis* in the kidneys of *Tilapia guineensis* in Senegal. This Myxosporidia differs from the species in description by its wider spores (10.3 (9-12) μ m) and its spherical polar capsules.

In India, *Myxobolus yogendrai* (Triparthi, 1952) Landsberg et al. [34], parasitize *Cirrhina mrigala*; its spore, although ovoid with a wider anterior end, possess an intercapsular triangle and four valvular folds.

Myxobolus nounensis Fomena et al. [35], is a parasite of the spleen and kidneys of *Sarotherodon galilaeus* and *Tilapia mariae* (Cichlidae) in the Noun River in Cameroon. Spores of this species are more developed (14.3 μ m × 12.8 μ m on average), same as its polar capsules (5.8 μ m × 4.5 μ m on average) and a well-developed intercapsular appendix is present.

These differences lead to think that the parasite of *Labeo batesii* is a new species. We propose the name *Myxobolus magai* sp. nov., referring to the locality of Maga where the fish host were captured.

Conclusion

Although Africa has one of the most diverse ichthyological fauna, and the genus Myxidium is the second major group of Myxosporidia, the description of Myxidium tetraodoni sp. nov. and M. anisocapsularis sp. nov. Only brings up to 18 the number of species of the genus Myxidium described in freshwater fishes of this continent, and to 11 the number of species of the genus described in freshwater fishes of Cameroon. M. tetraodoni is the first Myxosporidia described in fishes of the family Tetraodontidae and M. anisocapsularis is the only species of the genus Myxidium hitherto found with unequal polar capsules. Species of the genus Myxidium are generally coelozoic and rarely histozoic in fish host. The two new species of Myxidium described in the present work are coelozoic. These coelozoic Myxosporidia are generally less pathogenic than histozoic species, their pathogenic action seems to be less evident apart from the occlusion of bile and urinary ducts. In Africa, the species of Myxidium that are identified so far affect hosts belonging to 12 families. In decreasing importance order, these host families can be classified as: Cyprinidae (5 species); Claroteidae (2); Distichodontidae (2); Mochokidae (2); Mormyridae (2); Anabantidae (1); Aplocheilidae (1); Channidae (1); Cichlidae (1); Clariidae (1); Latidae (1); Tetraodontidae (1). Myxosporidia of the genus Myxobolus are generally known as histozoic parasites in freshwater fish. They form cysts of varying size in various organs in their host (gills, fins, stomach wall, intestinal wall, skin, heart, liver, operculum, eyes ...). Eiras et al. [8,9] estimated at about 905 species the number of Myxobolus described in fishes around the world. 33.25% of these species affect the gills in their hosts. The description of M. magai sp. nov. in the gills of Labeo batesii confirms the preference of this organ by the Myxosporidia of the genus Myxobolus.

Compliance with Ethical Standards

Animals used followed a protocol approved and authorized by Institutional Animal Care and Use Committee at Animals Biology and Physiology Department, Faculty of Science, University of Yaounde I, Cameroon.

Acknowledgement

The authors are thankful to the Faculty of Science, University of Yaounde I, Cameroon, for providing all the facilities to complete this work.

Conflict of Interest Statement

The authors declared: There is no conflict of interest.

References

- Abdel-Ghaffar F, Bashtar AR, Mehlhorn H, Al-Rasheid K, Al-Olayan E, et al. (2009) Ultrastructure and host parasite relationships of Kudoa pagrusi (Myxozoa) infecting the heart muscles of sea bream Pagrus pagrus (L.) from the Red Sea. Parasitology Research 106: 121-129.
- Lom J, Dykova I (2006) Myxozoan genera: definition and notes on taxonomy, life- cycle terminology and pathogenic species. Folia Parasitologica 53: 1-36.
- Lom J, Dykova I (1992) Myxosporidia (Phylum Myxozoa). In: Protozoan parasites of fish, Elsevier, Amsterdam-London-New York-Tokyo (ed.) pp: 159-235.
- Hessen EM, Zamzame ML (2004) Myxobolus sp. a possible new opportunistic parasite in immunocompromized patients in Ismailia. J Egypt Soc Parasitol 34: 925-930.
- Abdel-Ghaffar F, Morsy K, Bashtar AR, El-Ganainy S, Gamal S (2012) Thelohanellus niloticus sp. nov. (Myxozoa: Myxosporea) a parasite of the Nile carp Labeo niloticus from the River Nile, Egypt. Parasitology Research 112: 379-383.
- Harvell CD, Mitchell CE, Ward JR, Altizer S, Dobson A, et al. (2002) Climate warming and disease risks for terrestrial and marine biota. Science, 296, 2158-2162.
- Fiala I, Bartosova-Sojkova P, Whipps CM (2015) Classification and Phylogenetics of Myxozoa. Myxozoan Evolution, Ecology and Development pp: 85-110.
- Eiras JC, Molnar K, Lu YS (2005) Synopsis of the species of Myxobolus Bütschli, 1982 (Myxozoa: Myxosporea: Myxobolidae). Systematic Parasitology 61: 1-46.
- Eiras JC, Zhang J, Molnar K (2014) Synopsis of the species of Myxobolus Bütschli, 1882 (Myxozoa: Myxosporea, Myxobolidae) described between 2005 and 2013. Systematic Parasitology 88: 11-36.
- Butschli O (1882) Myxosporidia. Bronn's Klass. Ordn. Protozoa 1: 590-603.
- Eiras JC, Saraiva A, Cruz CF, Santos MJ, Fiala I (2011) Synopsis of the species of Myxidium Bütschli, 1882 (Myxozoa: Myxosporea: Bivalvulida). Systematic Parasitology 80: 81-116.
- 12. Diamant A, Lom J, Dykova I (1994) Myxidium leei n. sp., a pathogenic myxosporean of cultured sea bream Sparus aurata. Diseases of Aquatic Organisms 20: 137-141.
- 13. Jirku M, Bolek MG, Whipps CM, Janovy JJR, Kent ML et al. (2006) A new species of Myxidium (Myxosporea: Myxidiidae), from the western chorus frog, Pseudacris triseriata triseriata, and Blanchard's cricket frog, Acris crepitans blanchardi (Hylidae), from eastern Nebraska: morphology, phylogeny, and critical comments on amphibian Myxidium taxonomy. Journal of Parasitology 92: 611-619.
- 14. Lekeufack Folefack GB (2010) Faunistic and biology of Myxosporidia (Myxozoa: Myxosporea) parasites of some teleosts in the Sange river

(Wouri sub-tributary) in Cameroon. Ph.D. Thesis, Department of Animal Biology and Physiology, University of Yaounde I, Yaounde, Cameroon.

- 15. Nchoutpouen E (2015) Myxosporidia (Myxozoa: Myxosporea) parasites of some teleosts of the Noun Basin (West Region, Cameroon): taxonomy and biology of species parasitizing Oreochromis niloticus Linneaus, 1758 and Labeo parvus Boulenger. Department of Animal Biology and Physiology, University of Yaounde I, Yaounde, Cameroon.
- Fomena A, Lekeufack Folefack GB, Bouix G (2010) Two new species of Myxidium (Myxosporea: Myxidiidae) parasites of fresh water fish in Cameroon. Parasite 17: 9-16.
- Nchoutpouen E, Fomena A (2011) Description of three new species of Myxobolus (Myxosporea: Myxobolidae) parasites of Labeo parvus Boulenger, 1902, Cyprinid fish in Cameroon. Journal of Applied Bioscience 38: 2508-2517.
- Leumbe O, Bitom D, Mamdem L, Tiki D, Ibrahim A (2015) Mapping of flooding risk zones in the Sudano-Sahelian zone: case of Maga and its surroundings in the Far north Cameroon. Afrique Science 11: 45-61.
- 19. Leveque C, Paugy D, Teugels GG (1992) Fauna of freshwater and brackish fish from West Africa. Tome 2. O.R.S.T.O.M. (ed). Paris pp: 389-902.
- 20. Stiassny MLG, Teugels GG, Hopkins CD (2007) Freshwater and Brackish Fish from Lower Guinea, Western Central Africa. Volume 1. In: Tropical Fauna and Flora Collection, IRD (Ed), Paris.
- 21. Lom J, Arthur JR (1989) A guideline for the preparation of species descriptions in Myxosporea. Journal of Fish Diseases 12: 151-156.
- 22. Fomena A, Bouix G (1986) Contribution to the study of Myxosporidia parasites of freshwater fishes of Cameroon. I. New species of the genus Myxidium Bütschli, 1882. Acta Tropica 43: 319-333.
- 23. Kostoingue B, Faye N, Toguebaye BN (1998) New species of Myxosporidia of genus Myxidium Bütschli, 1882 and Myxobolus Bütschli, 1882 (Myxozoa: Myxosporea) of freshwater fishes of Chad (Central Africa). Journal of African Zoology 112: 249-259.
- 24. Auerbach M (1910) Zwei neue Cnidosporidien aus cyprinoiden Fischen. Zoologischer Anzeiger 36: 440-441.
- 25. Kudo RR (1921) On some Protozoa parasitic in freshwater fishes of New York. Journal of Parasitology 7: 166-74.
- Mitchell LG (1967) Myxidium macrocheili n. sp. (Cnidospora: Myxidiidae) from the largescale sucker Catostomus macrocheilus Girard,

and a synopsis of the Myxidium of North American freshwater vertebrates. Journal of Protozoology 14: 415-424.

- 27. Sakiti GN (1997) Myxosporidia and Microsporidia of freshwater fishes of Benin: faunistic, ultrastructure, biology. Ph.D Thesis, National University of Benin.
- Fomena A, Bouix G (1994) New Myxosoridea species (Myxozoa) from freshwater Telosts in Southern Cameroon (Central Africa). Journal of African Zoology 108: 481-491.
- 29. Abdel-Baki AZ (2009) Two new Myxidium species (Myxosporea: Myxidiidae) infecting the gallbladder of African flying fish, Cheilopogon nigricans and Suez fusilier, Caesio suevicus from the Red Sea, Egypt: a morphological and morphometric study. Parasitology Research 105: 513-518.
- Landsberg JH (1985) Myxosporean infections in cultured tilapias in Israel. Journal of Protozoology 32: 194-201.
- 31. Abakar-Ousman (2006) Myxosporidia (Myxozoa: Myxosporea) parasites of Chad freshwater fish: fauna and biology of the parasites species of Oreochromis niloticus (Linne, 1758) and Sarotherodon galilaeus (Linné, 1758) (Cichlidae). Ph.D. Thesis, Department of Animal Biology and Physiology, University of Yaounde I, Yaounde, Cameroon.
- 32. Kaur H, Singh R (2009) A new myxosporean species, Myxobolus eirasi sp. nov., a known species M. venkateshi Seenappa, Manohar, 1981 from the Indian major carp fish Cirrhina mrigala (Ham). Protistology 6: 126-130.
- 33. Fall M, Fomena A, Kostoingue B, Diebakate C, Faye N, et al. (2000) Myxosporidia (Myxozoa, Myxosporea) parasites of Cichlidae fish from Cameroon, Senegal and Chad, with the description of two new species. Annales des Sciences Naturelles 21: 81-92.
- Landsberg JH, Lom J (1991) Taxonomy of the genera of the Myxobolus/ Myxosoma group (Myxobolidae: Myxosporea), current listing of species and revision of synonyms. Systematic Parasitology 18: 165-186.
- 35. Fomena A, Bouix G (2000) Henneguya mbakaouensis sp. nov., Myxobolus nounensis sp. nov. and M. hydrocyni Kostoingue & Toguebaye, 1994 (Myxosporea, Myxozoa) parasites of Contropomidae, Cichlidae and Characidae (Teleosts) of Sanaga basin in Cameroon (Central Africa). Parasite 7: 209-214.

Page 10 of 10