

Alcohol Consumption is affected by a Protein Linked to the Circadian Rhythm

Keisuke Mizuo*

Assistant Professor, School of Medicine, Sapporo Medical University, Japan

It's a commonly heard question after New Year's: "Why do we drink the way we do?" A group of researchers has found that at least some of it has to do with a particular protein in the part of the forebrain that regulates, among other things, decision-making and reward perception.

It, the researchers announce that the presence of the *Bmal1* gene in the striatum affects alcohol consumption in both male and female mice ~ but in a sexually dimorphic manner. Male mice without the protein consumed more alcohol than those that had it, while female mice without the protein consumed less than females with it.

Bmal1 is also an integral element in the suprachiasmatic nucleus, the master circadian clock found in all mammals that regulates the sleep-wake cycle. Previous association analyses of clock genes revealed a potential role for *Bmal1* in alcohol-drinking behaviour. Expanding on this ~ and given evidence of sex differences in alcohol consumption and in some functions of clock genes ~ the researchers hypothesized that *Bmal1* may affect alcohol intake in a sex-dependent manner.

RISK IN FEMALES, PROTECTION IN MALES

The researchers created two lines of mice, using molecular biology methods to delete or "knock out" the *Bmal1* gene from the striatum's medium spiny neurons in one of them. The gene remained present in other parts of the body, since it plays a critical role in the circadian clock. The other line was used as a control.

Males who had the *Bmal1* gene deleted from the striatum were found to consume more alcohol than the ones that did not have it

deleted, while in the females, the results were the opposite: those without *Bmal1* consumed less alcohol than those that had it. (Normally, female rodents tend to consume more alcohol per body weight than males.)

"The main conclusion we can draw from this is that in females, *Bmal1* in the striatum confers risk, since they consume more alcohol when the gene is present," Amir says. "In males, the gene is protective, as they drink less alcohol. The sex differences you see in normal mice are eliminated when the gene is taken out of the striatum."

Amir notes that neither the sugar consumption nor circadian rhythms is affected by the deletion of the gene.

"It seems that striatal *Bmal1* plays a causal role in the control of alcohol consumption and makes an important contribution to sex differences in alcohol intake," he explains.

A BASIS FOR SEX-BASED TREATMENT?

The researchers believe this discovery can help in treating addiction in humans. For instance, while women report lower alcohol use and dependency than men, they suffer more adverse consequences of alcohol use and dependency.

"So far, the limited biological and pharmacological treatments for alcohol dependence don't distinguish between males and females, even though there are major differences in alcohol drinking behaviour and addiction between the sexes," he says. "By discovering sexually dimorphic mechanisms, addiction treatment specialists could ultimately use this knowledge to develop sex-based treatment."

*Correspondence to: Keisuke Mizuo, Assistant Professor, School of Medicine, Sapporo Medical University, Japan, E-mail: nagamine@sapmed.ac.jp

Received: December 06, 2021; Accepted: December 20, 2021; Published: December 24, 2021

Citation: Mizuo K (2021) Alcohol Consumption is Affected by a Protein Linked to the Circadian Rhythm. *J Alcohol Drug Depend* 9: 373. doi: 10.35248/2329-6488.21.9.373.

Copyright: ©2021 Mizuo K. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.