

Effect of Maize Prolamins on Peripheral Blood Mononuclear Cells from Celiac Disease Patients

Juan Pedro Ortiz Sánchez and Ana María Calderón de la Barca*

Departamento de Nutrición y Metabolismo, Centro de Investigación en Alimentación y Desarrollo, A.C. Carr. La Victoria, km 0.6, P. O. Box 1735. Hermosillo 83304, Sonora, Mexico

*Corresponding author: Ana María Calderón de la Barca, Departamento de Nutrición y Metabolismo, Centro de Investigación en Alimentación y Desarrollo, A.C. Carr. La Victoria, km 0.6, P. O. Box 1735, Hermosillo 83304, Sonora, Mexico, Tel: +52-662289 24 00; E-mail: juanpedroo@gmail.com

Received date: March 26, 2016; Accepted date: April 28, 2016; Published date: May 02, 2016

Copyright: © 2016 Sánchez JPO, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Celiac disease (CD) is an enteropathy induced by wheat prolamins (gliadins) and in some rare cases by maize prolamins (zeins) possibly due to a similar immune response. The aim was to study the cellular immune response to zeins in comparison to gliadins of peripheral blood mononuclear cells (PBMC) from CD patients. Isolated PBMC from two treated CD patients and three non-CD controls were challenged *in vitro* with gliadins or zeins and released gamma-interferon (IFN- γ) in culture medium was measured. PBMC were stimulated with gliadin or zein immunogenic peptides or their digested fractions (3-5 kDa). The gliadin peptide G33-mer induced an expected IFN- γ releasing of PBMC from both patients 1 and 2, with a higher level between days 0 and 6 for patients 1 and no differences para patient 2. The zein peptide Z34-mer induced a higher increase of IFN- γ in both CD patients at 0 day, even higher that it of G33-mer for patient 2, and both of them were highly decreased at day 6. Finally, the zein digested fraction induce an IFN- γ release similar to that of gliadin digested fraction in both cases, although negligible for patient 1 and significant for patient 2. In conclusion, the cellular response to zeins was partially similar to it of gliadins after an *in vitro* challenge.

Key words:

Celiac disease, T-cell response, Maize prolamins, Zeins

Abbreviations

CD: Celiac Disease; PBMC: Peripheral Blood Mononuclear Cells; HLA: Human Leucocyte Antigen; Ttg: Tissue Transglutaminase; PHA: Phytohaemagglutinin A; PT: Pepsin-Trypsin; Gd: Gliadin; G33-mer, Immunogenic Peptide of α -gliadin; Z34-mer, Immunogenic Peptide of α -zein

Introduction

CD is an immunologically mediated systemic disorder developed in genetically predisposal individuals, exacerbated by wheat and related cereals as barley and rye. Disease symptoms are promoted by inflammation of the intestinal mucosa, inducing gastrointestinal and/or extra-intestinal manifestations [1]. CD is a lifelong condition and gluten-free diet is the only treatment. One of the most important alternative cereals used for the gluten-free bakery products is maize; additionally, its prolamins have been used as a negative control in different studies on CD. By chance, in some of those studies maize prolamins have demonstrated adverse effects [2,3] inducing doubt about the maize use for dietary treatment of CD patients. The response to maize prolamins could be due to similarities between maize (zein) and wheat (gliadin) prolamins both with a high percentage of glutamine able to be deamidated by transglutaminase and proline residues that hinders a full digestion by gastrointestinal proteases [3].

The proposed pathogenesis of CD highlights the role of T-cells, after peptide presentation by dendritic cells to Th1 cells via the HLA-DQ2/8 context, activating them and consequently releasing cytokines, mainly

IFN- γ [4]. Therefore, IFN- γ is a marker of cellular response to different gluten peptides by *in vitro* assays; its advantage is that promotes tissue inflammation and has no autocrine effect on other PBMC, like monocytes [5-7]. Gluten specific T-cells producing IFN- γ can be found in peripheral blood of CD patients in gluten-free diet after a short gluten challenge [5]. Isolation and subsequent *in vitro* stimulation of these T-cells with a wide variety of dietary peptides, generates a reliable tool to evaluate the cellular response to gluten-free foods [6]. The aim of this study was to evaluate the T-cell response *in vitro* to maize prolamins in comparison to wheat prolamins of peripheral blood mononuclear cells (PBMC) from CD patients and PBMC from non-CD individuals as controls, after gluten-free diet followed by a three-day gluten challenge.

Materials and Methods

Patients

Patients underwent gluten-free diet for at least one month, and a three days challenge with at least 50 g/day gluten was made and blood samples were taken at day 0 and day 6. The ethical committee of the Centro de Investigación en Alimentación y Desarrollo (CIAD A.C.) approved the study and all samples were taken under informed written consent. Whole blood was taken (14 mL) from each patient by venipuncture into Vacutainer tubes (BD Medical Systems, USA). DNA was extracted from 200 μ L whole blood by the QIAamp DNA Blood Mini Kit (QIAGEN, USA) and genotyping of HLA-DQ2/DQ8 was done by real time PCR (Step One Plus, Applied Biosystems) using specific primers [8]. Isolation of peripheral blood mononuclear cells (PBMC's) from 12 mL blood was done using Ficoll-Paque PLUS (Amersham-Biosciences, Sweden) density gradient centrifugation technique. Plasma anti-gliadin (Gd) IgG, anti-Gd IgA, anti-zein IgA

and anti-transglutaminase (TG) IgA antibodies were analyzed by a direct enzyme-linked immunosorbent assay (ELISA), as previously reported [9]. IgA anti-gliadin and/or zeins and IgA anti-TG were expressed as an index value and it was calculated based on the mean of absorbance values of control individuals as reported before [9] and index values of 1.0 and above were considered as positive.

Peptide preparation

The immunogenic peptides α -gliadin 33-mer (LQLQFPQPPELPYPQPPELPYPQPPE; MW = 3914.51 Da), later referred to as G33-mer, and α -zein 34-mer (LQQAIAASNIPLSPLLFQSPALSLVQSLVQTIR; MW = 3646.32 Da), later referred to as Z34-mer, were supplied by United Biosystems (USA) with purities of 97.54% and 95.66%, respectively. Gliadins from wheat and zeins from maize (Sigma Chem Co, St. Louis, MO USA) were subjected to pepsin-trypsin (PT) digestion, as previously described [2]. All immunogenic peptides and digested prolamins were treated with transglutaminase (TG) from guinea pig liver (Sigma-Aldrich, St Louis, MO USA) 5 μ g/500 mg of protein in CaCl₂ 2 mM for 60 min at 37°C and then placed on ice. Separation of TG was performed by ultrafiltration (UF cell, Amicon Inc. Beverly, MA. USA.), with a 30 kDa cut-off membrane and peptides were recovered in sterile water.

Cell culture and cytokine assays

Isolated PBMC were incubated at a final concentration of 2 x 10⁵ cells/mL on culture plates and cultured in Dulbecco's Modified Medium (D-MEM) containing 10% fetal calf serum (FCS), 100 U/mL penicillin and 100 μ g/mL streptomycin (Gibco, USA) at 37°C in a 5% CO₂ atmosphere. The immunogenic peptides were used in the experiments at final concentration of 50 μ g/mL and the digested prolamins at 100 μ g/mL. Phytohemagglutinin A (PHA) (Sigma Aldrich, USA) was used as positive control at concentration of 25 μ g/mL. After 20 h, supernatants were collected and frozen at -70°C prior to cytokine evaluation. ELISA kits were used for IFN- γ (Mabtech, Sweden) detection according to manufacturer.

Statistical analysis

Experiments were performed in triplicate, results are given as mean values that were compared after ANOVA. Statistical significance among days 0 and 6 was compared by Student's one sample T-test and statistical significance among treatments by Tukey-Kramer multiple comparison test using the statistical software NCSS, version 2001. The p-values of 0.05 or less were considered as statistically significant.

Results

The characteristics of the three control subjects and the two celiac patients are described in Table 1. All the control individuals showed negative indexes (<1.0) of anti-Gd IgG, anti-Gd IgA and anti-TG IgA antibodies. Celiac patient 1 presented positive indexes (>1.0) for anti-Gd IgG and anti-TG IgA antibodies, while patient 2 had for anti-Gd IgG, anti-Gd IgA, anti-TG IgA and anti-Zein IgA antibodies (Table 1).

Production of IFN- γ in PBMC of control individuals was not stimulated with any of gliadin or zein immunogenic peptides or PT-digested fractions compared to untreated cells and the poor response was averaged to simplify the results graphically (Figure 1). As expected, in both CD patients, the α -gliadin immunogenic peptide

(G33-mer) increased release of IFN- γ in PBMC respect to controls (p<0.005) at days 0 and 6. Additionally, on patient 1 IFN- γ release was higher at day 6 compared to day 0 (p<0.05), while for patient 2 the IFN- γ increase was similar for both days.

Interestingly, an increase in IFN- γ release by stimulation with Z34-mer peptide was observed at day 0, mostly on patient 2 (p<0.0005) respect to controls, but stimulation diminishes at day 6 in both cases (patients 1 and 2), remaining higher than controls just for patient 2 (p<0.05), as it is shown in Figure 1. Both zeins and gliadins fractions ZFIII and GFIII induced a similar IFN- γ release in PBMC at 0 or 6 days; however, such increase was not significant (p>0.05) respect to controls for patient 1, while it was significant (p<0.005) as compared with IFN- γ releasing for PBMC from patient 1 or controls, for patient 2.

Discussion

Both CD patients described as patient 1 and 2, reported extra-intestinal and intestinal symptoms that were alleviated after a gluten-free diet. They also showed some positive indexes for antibodies anti-gliadins, anti-transglutaminase and patient 2 against zeins (Table 1). The deamidation of gluten peptides in lamina propria by tissue transglutaminase is the first step in CD pathogenesis, and after activation of the immune response, IgA anti-TG autoantibodies are induced; they characterize CD [1]. Additionally, in active CD there are antibodies against gliadins, the exogenous antigen. Interestingly, only patient 2 had a positive index for anti-zeins IgA antibodies, as it was previously found in some CD patients by Cabrera-Chávez et al. [9]. Peripheral blood effector T-cells reactive to gliadins were found in both patients before the *in vivo* gluten challenge and this result agrees with those found by Liu et al. [7] who detected higher levels of IFN- γ in CD patients that carried both haplotypes HLA-DQ2 and/or HLA-DQ8. Furthermore, they also observed that the stimulation of peripheral blood T-cells proliferation is possible without a previous *in vivo* challenge. Indexes of anti-transglutaminase and anti-gliadin IgA antibodies remained positive, especially on patient 2 (Table 1), since half-life of IgA antibodies last for about 4 months [10], patients possibly did not follow a strict gluten-free diet. Therefore, the *in vivo* gluten challenge was not effective.

Subject	Age (years)	Haplotype or alleles	Index of antibodies				Symptoms
			IgG anti-Gd	IgA anti-Gd	IgA anti-TG	IgA anti-Zn	
Control 1	30	DQA1*501, DQA1*0301	0.899	0.773	0.744	0.768	None
Control 2	30	DQA1*501, DQB1*302/3	0.695	0.451	0.356	0.353	None
Control 3	27	DQA1*0301	0.796	0.728	0.668	0.619	None
Patient 1	31	HLA-DQ2	1.316	0.748	1.046	0.938	Migraine, fatigue and bloating
Patient 2	46	DQA1*0501	1.217	1.238	1.280	1.68	Anemia, constipation, bloating

-
9. Cabrera-Chávez F, Iameti S, Miriani M, Calderón de la Barca A, Mamone G, et al. (2012) Maize prolamins resistant to peptic-tryptic digestion maintain immune-recognition by IgA from some celiac disease patients. *Plant Food Hum Nutr* 67: 24-30.
 10. Cerutti A (2010) Immunology. IgA changes the rules of memory. *Science* 328: 1646-1647.