

Effects of Bee Bread Supplementation on Isokinetic Knee Extension and Flexion of Lower Limb

Chee Ping Fadzel Wong^{1*}, Chee Keong Chen², Foong Kiew Ooi², Mahaneem Mohamed³

¹Department of Nutrition, Universiti Malaysia Sabah, Kota Kinabalu, Sabah, Malaysia; ²Department of Sport Sciences, Universiti Sains Malaysia, Kelantan, Malaysia; ³Department of Physiology, Universiti Sains Malaysia, Kelantan, Malaysia

ABSTRACT

A nutritional strategy is the common types of ergogenic aid used by athlete to enhance their sport performance. Athlete usually consumes bee products before a sports competition to make sure they have adequate nutrition, maximize their energy storage and enhance their sport performance. Scientific data on the ergogenic effects of bee products consumed at pre exercise on isokinetic knee extension and flexion of lower limb is scanty. This study investigated the effects of 8 weeks of bee bread supplementation at pre exercise on isokinetic knee extension and flexion of lower limb among athletes. Twelve volunteer athletes were recruited in this scientific study. Subjects consumed either bee bread at a dosage of 20 g.d-1 or placebo for 8 weeks prior to the experimental trial. Subjects' isokinetic knee extension and flexion of lower limb were measured before and after 8 weeks of experimental period. Statistical analyses were performed using ANOVA with repeated measures. This present study found that bee bread supplementation showed some positives impact on few measured parameters of isokinetic knee extension and flexion of lower limb.

Keywords: Isokinetic; Extension; Flexion; Lower limb; Sport performance

INTRODUCTION

Bee products have been become popular ergogenic supplement among athletes. Several studies investigated bee products and reported bee products have antioxidant, anti-inflammatory antitumor, anti-allergic, and antimicrobial. There are limited scientific researches on effect of bee products supplementation on sport performance [1]. For instances, found that supplementation of honey at a dosage of 15 g every 16 km during a simulated 64 km cycling time trial was significantly faster to complete time trial compared to the placebo trial reported that combination of aerobic dance exercise and honey supplementation (20 g of honey diluted in 300 mL of plain water for 8 weeks) elicited more beneficial effects on bone health and immune function compared to the aerobic dance exercise or honey supplementation alone [2]. It found that ingestion of 500 ml of honey drink one hour before trial and 3 ml per kg body weight of honey drink every 20 minutes during the running trial

was as good as a sports drink in improving running time trial performance.

Indicated that combination of jumping exercise and honey supplementation may elicit beneficial effects on tibia and femur bone generally when compared to either jumping exercise or honey supplementation alone in young female rats [3]. However, found that acute supplementation of honey beverage at a dosage of 1 g.kg body weight before and during soccer stimulation test did not significantly improve Progressive Shuttle-Run (PSR) test to exhaustion compared to the placebo trial. To date, there is scanty information on the possible beneficial effects of bee bread supplementation on isokinetic knee flexion and extension [4]. Therefore, the aim of this study was to investigate effects bee bread supplementation on isokinetic knee extension and flexion of lower limb among athletes.

METHODOLOGY

Twelve Malaysian male athletes were recruited in this study. Subjects were asked to refrain from exercise for 24-hour before

Corresponding Author: Chee Ping Fadzel Wong, Department of Nutrition, Universiti Malaysia Sabah, Kota Kinabalu, Sabah, Malaysia, Tel: 1151845358; E-mail: fadzel@ums.edu.my

Received: June 23, 2021; **Accepted:** July 7, 2021; **Published:** July 14, 2021

Citation: Wong CPF, Chen CK, Ooi FK, Mohamed M (2021) Effects of Bee Bread Supplementation on Isokinetic Knee Extension and Flexion of Lower Limb. J Yoga Phys Ther. 11:355.

Copyright: © 2021 Wong CPF, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the tests to ensure that they have adequate rest before the isokinetic knee extension and flexion tests. Their food diary and physical activity diary for the last 72 hours were also collected. Subjects recorded their 3-day food diary prior to the first trial and repeated the same diet over 3 days before the days of consecutive test to minimize the differences in muscle glycogen between the trials [5]. Isokinetic dynamometer (Multi Joint System 3 Pro: Biodex Medical System, USA) was used to measure subjects' isokinetic knee extension and flexion of lower limb. Testing angular velocities were set at $180^{\circ} \cdot s^{-1}$ and $300^{\circ} \cdot s^{-1}$ to measure knee extension and flexion muscular strength and power of subjects prior and post 8 weeks of experimental period. These testing angular velocities were used to measure the isokinetic right and left knee extension and flexion peak torque and average power of the subjects. Subject was required to do a quadriceps and hamstring stretching exercise before the exact testing to prevent injuries. Subject familiarised with the isokinetic dynamometer protocol before the actual testing for the measurements for knee extension and flexion [6].

To begin the test, subjects were seated while leaning against a backrest tilted 85° from the horizontal position. The chair and dynamometer were rotated to 90° . Knee attachment was moved attached to the dynamometer. Subject's knee axis of rotation was aligned with dynamometer shaft. The seat was raised or lowered or the subject was moved towards or away from the dynamometer to fine adjust the testing position [7]. Knee attachment was adjusted so that it was proximal to medial malleoli. Subject's knee, shoulder, waist and thigh were strapped to minimise body movement and gave stabilisation position during the test. Shoulder straps were applied diagonally across the chest to prevent excessive upper body movement. The lateral femoral epicondyle was palpated and used as a bony landmark for matching the axis rotation of the knee joint and the axis rotation of the dynamometer shaft [8]. All subjects were fully informed about the procedures to determine the isokinetic muscular strength and power test before performing these tests. There were asked to do 5 repetitions for the $180^{\circ} \cdot s^{-1}$ angular to velocity and 10 repetitions for the $300^{\circ} \cdot s^{-1}$ angular velocity for the each lower limb. The subjects were given sixty seconds to rest between each angular velocity. All the statistical analyses were computed by using the Statistical Programme for the Social Sciences (SPSS) version 26.0 (SPSS Incorp, United States). The level of significance for all analysis was set at $p < 0.05$. All the collected data were expressed in mean and standard deviation (mean $\pm$ SD). ANOVA with repeated measures was used to determine the differences of the measured isokinetic knee extension and flexion parameters over time and between trials [9]. Bonferroni adjustment for multiple comparisons was used to locate the differences when repeated measures analysis of variance revealed a significant main effect of time.

RESULTS

Twelve volunteer athletes were recruited in this scientific study. The mean maximum oxygen consumption (VO_{2max}) reflected that the subjects had good cardio-respiratory fitness [10]. Mean Body Mass Index (BMI) of the subjects was categorised as healthy (Table 1).

Physical characteristics of subjects	Mean $\pm$ SD
Age (years)	24.0 $\pm$ 1.8
VO_{2max} (mL. kg ⁻¹ . min ⁻¹)	52.0 $\pm$ 2.8
Height (cm)	173.7 $\pm$ 6.1
Body mass (kg)	67.4 $\pm$ 6.2
Body mass index (kg. m ⁻¹)	22.3 $\pm$ 1.3

Table 1: Mean and SD of physical characteristics of the subjects.

Figure 1 showed mean values of isokinetic right and left knee extension peak torque and average powers at $180^{\circ} \cdot s^{-1}$. There was a significant main effect of time on isokinetic right knee extension peak torque and average power at $180^{\circ} \cdot s^{-1}$ in the bees bread trial ($p < 0.05$) [11]. Isokinetic right knee extension peak torque and average powers at $180^{\circ} \cdot s^{-1}$ was significantly higher the post test compared to pre test in the bee bread trial ($p < 0.05$). Figure 1 showed mean values of isokinetic right and left knee flexion peak torque and average power at $180^{\circ} \cdot s^{-1}$. There was significant main effect of time on isokinetic right knee flexion peak torque and average power at $180^{\circ} \cdot s^{-1}$ in the bee bread trials ($p < 0.05$). Isokinetic right knee flexion peak torque and average powers at $180^{\circ} \cdot s^{-1}$ was significantly higher at post test compared to pre test in the bee bread trial ($p < 0.05$).




Figure 1: Right and left knee extension and flexion peak torque and average powers at $300^{\circ} \cdot s^{-1}$ in the placebo and the bee bread trials. Data are presented as mean $\pm$ SD.

The mean values of isokinetic right and left knee extension peak torque and average power at $300^{\circ} \cdot s^{-1}$ are presented in the Figure 2. There was no significant main effect of time on isokinetic right and left knee extension peak torque and average power at $300^{\circ} \cdot s^{-1}$ in both the trials ($p > 0.05$) [12]. There was no analysis of significant difference in isokinetic right and left knee extension peak torque and average power at $300^{\circ} \cdot s^{-1}$ between pre and post tests in both the trials ($p > 0.05$). Figure 2 showed the mean values of isokinetic right and left knee flexion peak torque and average power at $300^{\circ} \cdot s^{-1}$. There was no significant main effects of time on isokinetic right and left knee flexion peak torque and average powers at $300^{\circ} \cdot s^{-1}$ in both the trials ($p > 0.05$). There was no significant difference in isokinetic right and left knee flexion peak torque and average power at $300^{\circ} \cdot s^{-1}$ between pre and posts tests in both the trials ($p > 0.05$).




Figure 2: Right and left knee extension and flexion peak torque and average powers at 300°/s in the placebo and the bee bread trials. Data are presented as mean $\pm$ SD.

DISCUSSION

Isokinetic assessment is the most common method used to measure isokinetic extension and flexion of lower limb muscles such as quadriceps and hamstring muscles. This present study found that there were significant increases in isokinetic right knee extension peak torque at 180°/s-1, right knees extension the average power at 180°/s-1, right and left knee flexion peak torque 180°/s-1 and right and left knee flexion average powers at 180°/s-1 between pre and post test after 8-week supplementation in the bee bread trial. To our knowledge, this is the first study exploring the effect of bee bread supplementation on muscular strength and power [13]. However, there was a study that investigated the effect of combined aerobic dance exercise and honey supplementation on muscular strength and power. This previous study reported that combined of aerobic dance exercise and honey supplementation for 8 weeks elicited more beneficial effects on muscular strength and power compared to the aerobic dance exercise or honey supplementation alone.

Bee bread contains high protein, i.e. essential amino acid and non-essential amino acid and it is speculated that protein contained in this supplement enhanced the muscular strength and power compared to placebo in this present study. Protein supplementation is one of the most common nutritional supplements used by athlete to increase the muscular strength and power. Skeletal muscle is undergoing constant remodelling through the continuous and stimulation process of muscle protein synthesis and muscle protein breakdown. Hence, it is speculated that bee bread supplementation was able to increase new or repair damaged muscle fibres through a cellular process where it fuses muscle fibres together to form new muscle myofibrils compared to the placebo in the present study. Skeletal muscle is composed of thread-like myofibrils and sarcomeres that form a muscle fibre and are the basic units of contraction. These new synthesis or repaired myofibrils increase in size of a muscle and number of myofibrils to create muscle become hyperplasia and hypertrophy and lead to increased muscular strength and power with bee bread supplementation [14].

Many previous studies have reported that protein supplementation play an important role to repair and remodelling process of skeletal muscle fibres to increase muscular strength and power. Protein has important

implications for improving both muscle size and strength found that supplementation of whey protein at a dosage of 1.2 g/kg/day for 6 weeks significantly increased the peak torque compared to the placebo group. Based on these previous study findings, it seems that bee bread supplementation in this study was more advantageous compared to the placebo in terms of enhancing muscular strength and power. This present finding was similar with previous study finding by who also found that 20 g of honey supplementation alone for 8 weeks only improved few measured parameters of muscular and strength. Based on this previous finding, it is demonstrated that specific training of aerobic dance exercise with honey supplementation was crucial to increase muscular strength and power compared to the aerobic dance exercise or honey supplementation alone. It is speculated that only with bee bread supplementation alone without any specific training program in this present study was not sufficient to increase all the measured parameters of muscular and strength power.

CONCLUSION

Supplementation of bee bread showed some positives impact on few measured parameters of isokinetic knee extension and flexion of lower limb.

CONFLICT OF INTEREST

The author declare no conflict of interest.

REFERENCES

- Nagai T, Nagashima T, Myoda T, Inoue R. Preparation and functional properties of extracts from bee bread. *Nahrung*. 2004;48(3):226-229.
- Nagai T, Sakai M, Inoue R, Inoue H, Suzuku N. Antioxidative activities of some commercially honeys, royal jelly and propolis. *Food Chem*. 2001;75(2):237-240.
- Nagai T, Nagashima T, Suzuku N, Innoue R. Antioxidant activity and angiotensin I-converting enzyme inhibition by enzymatic hydrolysates from bee bread. *Z Naturforsch C J Biosci*. 2005;60(2):133-138.
- Nakajima Y, Tsuruma K, Shimazawa M, Mishima S, Hara H. Comparison of bee product based on assays of antioxidant capacities. *BMC Complement Alter Med*. 2009;9(4):1-9.
- Yang X, Guo D, Zhang J, Wu M. Characterization and antitumor activity of pollen polysaccharide. *Int Immunopharmacol*. 2007;7(4):427-434.
- Medeiros KCP, Figueiredo CAV, Figueredo TB, Freire KRL, Santos FAR, Alcantara-Neves NM, et al. Anti-allergic effect of bee pollen phenolic extract and myricetin in ovalbumin-sensitized mice. *J Ethnopharmacol*. 2008;119(1):41-46.
- Abouda Z, Zerdani I, Kalalou I, Faid M, Ahami MT. The antibacterial activity of moroccan bee bread and bee-pollen (fresh and dried) against pathogenic bacteria. *Res J of Microbiol*. 2011;6(4):376-384.
- Earnest CP, Lancaster SL, Rasmussen C J, Kerkisck CM, Lucia A, Greenwood MC, Almada AL, Cowan PA, Kredier RB (2004) Low Vs high glycemic index carbohydrate gel ingestion during

- stimulated 64 km cycling time trial performance. *J Strength Cond Res.* 18(3): 466-472.
9. Tavafzadeh SS, Ooi FK, Krasilshchikov O, Sulaiman SA. Effect of a combination of jumping exercise and honey supplementation on the mass, strength and physical dimensions of bones in young female rats. *J ApiProduct ApiMedic Sci.* 2011;3(1):26-32.
 10. Abbey EL, Rankin JW. Effect of ingesting of honey sweetened beverage on soccer performance and exercise induced cytokine response. *Int J Sport Nutr Metab.* 2009;19(6):659-672.
 11. Schenk TL, Costley CD. When food becomes a drug: Nonanabolic nutritional supplement use in athletics. *Amer J Sports Med.* 2002;30(6):907-916.
 12. Ratamess NA, Kraemer WJ, Volek JS, Rubin MR, Gomez AL, French DN, et al. The effects of amino acid supplementation on muscular performance during resistance training overreaching. *J Strength Cond Res.* 2003;17(2):250-258.
 13. Tarnopolsky MA, Atkinson SA, MacDougall JD, Chesley A, Phillips S, Swarcz HP, et al. Evaluation of protein requirement for trained strength athletes. *J Appl Physiol.* 1985;73(5):1986-1995.
 14. Burke DG, Chilibeck PD, Davison KS, Candow DG, Farthing J, Smith-Palmer T, et al. The effect of whey protein supplementation with and without creatine monohydrate combined with resistance training on lean tissue mass and muscle strength. *Int J Sports Nutr Exerc Metab.* 2001;11(3):349-364.