

## Effects of Cultural Conditions on the Production of Extracellular Protease by *Streptomyces albolongus* and *Streptomyces aburaviensis*

Naznin Akhtar<sup>1\*</sup>, Abu Sayeed Mohammad Mahmud<sup>2</sup>, Muhammad Shahjalal Khan<sup>3</sup>, Tarannum Taznin<sup>4</sup>, Muhammad Ehteshamul Haque<sup>5</sup>, Sharmin Sultana<sup>5</sup> and Sharmin Sultana<sup>5</sup>

<sup>1</sup>Tissue Banking and Biomaterial Research Unit (TBBRU), Atomic Energy Research Establishment (AERE), Savar, Dhaka, Bangladesh

<sup>2</sup>Industrial Microbiology Research Division, BCSIR Laboratories, Bangladesh

<sup>3</sup>Department of Microbiology, University of Chittagong, Bangladesh

<sup>4</sup>Department of Microbiology, Jessore Science and Technology University, Bangladesh

<sup>5</sup>Department of Microbiology, University of Chittagong, Bangladesh

### Abstract

Proteolytic activity of two isolates of actinomycetes, *Streptomyces albolongus* and *Streptomyces aburaviensis* was investigated on the basis of their ability to hydrolyze skimmed milk casein, egg albumin and gelatin. Both isolates were found to have potential for extracellular proteases production. Effects of culture conditions for the production of extracellular protease from *S. albolongus* and *S. aburaviensis* were determined. Highest protease yield from *S. albolongus* was obtained after 5 days of incubation with an initial pH 7, at static state when inoculated in medium composed of 1% glucose, 2% beef extract, 0.2% yeast extract, 0.1% KH<sub>2</sub>PO<sub>4</sub>, 0.3% K<sub>2</sub>HPO<sub>4</sub>, and trace MgSO<sub>4</sub>·7H<sub>2</sub>O. Optimum incubation conditions for *S. aburaviensis* were 4 days, with an initial medium pH 8 at shaking condition (100 rpm). *S. aburaviensis* preferred 1.5% lactose and 1.5% tryptone as a carbon and nitrogen source. Both the isolates showed maximum protease yield at 37°C. The result of the present study might be helpful for large-scale production of extracellular protease from these *Streptomyces* spp.

**Keywords:** *Streptomyces albolongus*; *Streptomyces aburaviensis*; Extracellular proteases; Culture conditions

### Introduction

Proteases play a colossal role in biotechnology and are widely used in tanning industry, in the manufacturing of biological detergents, meat tenderization, peptide synthesis, food industry, pharmaceutical industry, and in bioremediation processes [1-3]. Although protease can be obtained from several sources, including plant, animal and microorganisms, microbial proteases are preferred in view of their rapid growth, ease of cultivation purification and genetic manipulation. A large number of microorganisms have been reported for protease production [4-8].

Microbial proteases can be produced using many processes like solid-state fermentation and submerged fermentation [9,10]. Cultural conditions (physical, chemical and nutritional factors) play significant role in the production of extracellular proteases by microorganisms [11-13]. Physical factors include aeration, temperature, pH and incubation time. In addition to these physical factors, nutritional factors such as the sources of carbon and nitrogen also significantly affect protease production.

Researchers have examined bacteria and fungi from various habitats to obtain suitable proteases. At present, majority of commercially available proteases are secreted by *Bacillus* spp., although there have been increasing reports of the potential use of proteases of fungal origin [14-16]. Actinomycetes as a source of naturally occurring extracellular protease is overlooked and information on proteases from actinomycetes has been limited. However, the industrial demands of proteolytic enzymes stimulate the search of new enzyme sources with extended range of applications. Among the actinomycetes, several species of the *Streptomyces* are among the most important industrial microorganisms because of their capacity to produce numerous bioactive molecules, particularly antibiotics. *Streptomyces* species are heterotrophic feeders, which can utilize both complex and simple

molecules as nutrients. In addition to antibiotics, *Streptomyces* species liberate several extra cellular enzymes [17]. They produce variety of extra cellular proteases that have been related to aerial mycelium formation and sporulation [18]. In view of the above, the present work was undertaken to study the effects of culture conditions on the production of extracellular proteases by *S. albolongus* and *S. aburaviensis*, and also aimed at optimization of media composition, which has been predicted to play a significant role in enhancing the production of protease.

### Materials and Methods

#### Microorganism

Two isolates of actinomycetes, *S. albolongus* (A<sub>5</sub>) and *S. aburaviensis* (RB<sub>20</sub>) were collected from the laboratory stock culture of the Department of Microbiology, University of Chittagong. For the growth and preservation of the isolates, nutrient agar was used.

#### Screening of the isolate for proteolytic activity

The organisms' ability to produce extracellular protease enzyme were determined by growing the isolates in solid medium containing protein sources like gelatin, skimmed milk casein and boiled egg albumin. Secondary screening of the isolates was done by measuring the protease activity in liquid medium by quantitative method. For

**\*Corresponding author:** Naznin Akhtar, Scientific Officer, Tissue Banking and Biomaterial Research Unit (TBBRU), Atomic Energy Research Establishment (AERE), Savar, Dhaka, Bangladesh, E-mail: [naznin.akhtar@baec.org.bd](mailto:naznin.akhtar@baec.org.bd)

**Received** February 05, 2013; **Accepted** February 10, 2013; **Published** February 20, 2013

**Citation:** Akhtar N, Mahmud ASM, Khan MS, Taznin T, Haque ME, et al. (2013) Effects of Cultural Conditions on the Production of Extracellular Protease by *Streptomyces albolongus* and *Streptomyces aburaviensis*. Enz Eng 2: 110. doi:10.4172/2329-6674.1000110

**Copyright:** © 2013 Akhtar N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

this purpose, the isolates were inoculated into three liquid medium, viz. (i) Peptone yeast extract-dextrose broth (Yeast Extract 1%, Peptone 2%, Dextrose 2%) [19] (ii) Tryptone-yeast extract-dextrose broth (Tryptone 1%, Dextrose 0.1%, Yeast extract 0.5%) [20] (iii) Gelatin-yeast extract- glucose broth (Gelatin 1%, glucose 1%, Yeast Extract 0.2%,  $K_2HPO_4$  0.3%,  $KH_2PO_4$  0.1%,  $MgSO_4 \cdot 7H_2O$  trace) [21] at  $p^H$  7.0 and were incubated at  $35 \pm 2^\circ C$  for 5 days [22].

### Measurement of enzyme activity

After incubation, the broth cultures were filtered with Whatman grade 1 filter paper. Then the filtrates were centrifuged at 8,000 rpm for 15 minutes at  $4^\circ C$ . The supernatant was used as crude enzyme. The enzymes were stored at  $4^\circ C$  with few drops of toluene/Sodium Azide to avoid bacterial contamination. Protease assay was done by using the method described by Meyers and Ahearn [23], which is a modified method of Hayashi et al. [24]. Briefly, 3 ml of crude enzyme, 3 ml of citrate phosphate buffer and 3 ml of 1% (w/v) casein was taken in a 25 ml test tube, and the tube was placed in a water bath at  $35^\circ C$  for 1 hour. Enzyme-substrate reaction was stopped by adding 5 ml of 20% (w/v) trichloroacetic acid (TCA). After one hour, the solution was filtered by Whatman grade 540 (ash less) filter paper. From the filtrate, 1 ml enzyme-substrate mixture was taken into a test tube and 2 ml of 20%  $Na_2CO_3$  was added to it. To this mixture, 1 ml of Folin-Ciocalteu reagent was added and the contents of the tube were mixed well immediately. After 30 minutes, 6 ml distilled water was added to the tube and absorbance of the solution was measured at 650 nm in a Vis-UV spectrophotometer (LaboMedInp). The amounts of amino acids released were determined using a standard curve plotted from known concentration of tyrosine. The enzyme activity was expressed in Unit. One unit of enzyme was defined as the amount of enzyme that releases 1 g of tyrosine from substrate (casein) per hour, under assay conditions.

### Biomass yield

The actinomycetes cultures were filtered through Whatman grade 1 filter paper. The filter paper was dried in oven at  $80^\circ C$  to a constant weight. The amount of biomass was calculated by subtracting the weight of filter paper. The yield was expressed as mg/g of protein.

### Optimization of culture conditions

Broth cultures were carried out at various culture conditions such as temperature (10, 27, 37,  $45^\circ C$ ), initial  $p^H$  of the culture media (4, 5, 6, 7, 8, 9), incubation time (1, 2, 3, 4, 5, 6, 7 days), to optimize the culture conditions for the production of extracellular proteases. To determine the effects of aeration, inoculated mediums were incubated at both stationary and shaking condition (100 rpm), keeping other experimental conditions at optimum.

### Effects of carbon and nitrogen sources

The production of extracellular proteases under different carbon and nitrogen availability was studied in the liquid culture medium. Four carbon sources (glucose, lactose, fructose, galactose), five organic and two inorganic nitrogen sources (Gelatin, Yeast Extract, peptone, Beef Extract, Pulse,  $KNO_3$ ,  $(NH_4)_2HPO_4$ ) were added to the medium, and the effect of this carbon and nitrogen sources on the production of protease was recorded. To ascertain optimum percent of carbon and nitrogen sources, the study was carried out with 0.5 to 2.5% carbon and 0.5 to 2.5% nitrogen sources, keeping other experimental conditions at optimum.

## Results

Two isolates of actinomycetes, *Streptomyces albolongus* and *Streptomyces aburaviensis* were tested for proteolytic activity. Primary screening was done by the boiled egg albumin degradation, gelatin hydrolysis and skimmed milk casein hydrolysis method. Both the isolates showed clear zone of hydrolysis in gelatin agar plate and casein agar plate after 3 days of incubation at  $37^\circ C$  (Figures 1A and 1B). Complete degradation of egg albumin was observed after 7 and 9 days of incubation at  $37^\circ C$  for *S. albolongus* and *S. aburaviensis*, respectively (Figure 1C). The isolates were allowed to grow in three liquid media and maximum enzyme activity were found in Gelatin-yeast extract- glucose broth and tryptone-dextrose-yeast extract broth for *S. albolongus* and *S. aburaviensis*, respectively (Table 1).

### Effects of incubation period on the production of proteases

*S. albolongus* showed maximum enzyme production (789.14 U/ml) after 5 days of incubation and highest protease production (567.99 U/ml) by *S. aburaviensis* was recorded after 4 days of incubation (Figure 2). For both isolates, highest biomass yield (140 mg/g substrate for *S. albolongus* and 150 mg/g substrate for *S. aburaviensis*) were observed after 6 days of incubation, Table 2 associated with surface and sedimentary growth with white sporulation in case of *S. albolongus* and grayish sporulation in case of *S. aburaviensis*. The  $p^H$  of the culture filtrates was ranged from 5.26 to 6.14 and 6.10 to 6.90 for *S. albolongus* and *S. aburaviensis*, respectively.

### Effects of medium pH on the production of protease

Figure 3 represents the protease activities of the isolate *S. albolongus* and *S. aburaviensis* at different initial  $p^H$  of culture medium. *S. albolongus* showed highest enzyme production (795.32 U/ml), having medium  $p^H$  7.0, but highest biomass yield (160 mg/g substrate) was recorded with medium  $p^H$  8.0 (Table 3). The  $p^H$  of the culture filtrate ranged from 5.12 to 7.88. The isolate exhibited surface and sedimentary





Figure 1: Photograph showing clear zone of hydrolysis A. casein hydrolysis by *S. albolongus* (A-1) and *S. aburaviensis* (A-2), B. Gelatin hydrolysis by *S. albolongus* (B-1) and *S. aburaviensis* (B-2) and C. Egg albumin degradation.

Culture medium	Protease activity (U/ml) Isolates of actinomycetes	
	<i>S. albolongus</i>	<i>S. aburaviensis</i>
Yeast Extract 1%, Peptone 2%, Dextrose 2%	173.07	160.02
Tryptone 1%, Dextrose 0.1% Yeast extract 0.5%	326.92	474.58
Gelatin 1%, glucose 1%, Yeast Extract 0.2%, $K_2HPO_4$ 0.3%, $KH_2PO_4$ 0.1% $MgSO_4$ trace	631.86	253.43

Enzyme-substrate reaction  $p^H$  and temperature was 6.5 and  $35^\circ C$ , respectively.


Table 1: Protease activities of the isolates in different liquid medium.



**Figure 2:** Effects of incubation period on the production of protease by *S. albolongus* and *S. aburaviensis*. Enzyme production was carried out in broth culture and incubation temperature was  $35 \pm 2^\circ\text{C}$ , medium pH 7.0, enzyme-substrate reaction pH and temperature was 6.5 and  $35^\circ\text{C}$ , respectively.

Incubation Period (in days)	Biomass yield (mg/gm of substrate)	
	<i>S. albolongus</i>	<i>S. aburaviensis</i>
1	4	12
2	40	30
3	60	36
4	80	102
5	136	134
6	140	150
7	132	121

**Table 2:** Effects of incubation period on biomass yield.



**Figure 3:** Effects of medium pH on the production of protease by *S. albolongus* and *S. aburaviensis*. Enzyme production was carried out in broth culture and incubation temperature was  $35 \pm 2^\circ\text{C}$ , enzyme-substrate reaction pH and temperature was 6.5 and  $35^\circ\text{C}$ , respectively.

growth with white sporulation at medium pH 7.0. On the other hand, *S. aburaviensis* showed highest enzyme production (612.63 U/ml) and maximum biomass yield (180 mg/g substrate) at pH 8.0. The pH of the culture filtrates ranged from 5.19 to 7.93. Biomass characteristics of the isolate were varied with the pH of the medium.

### Effects of incubation temperature on production of proteases

Both *S. albolongus* and *S. aburaviensis* showed maximum protease production (747.24 U/ml and 619.50 U/ml respectively) (Figure 4) and maximum biomass yield (152 mg/g substrate and 174 mg/g substrate respectively), at  $37^\circ\text{C}$  associated with surface and sedimentary growth with sporulation. Table 4 represents biomass yields of the isolates at different incubation temperature. The pH of the culture filtrates was ranged from 7.10 to 7.89 for *S. albolongus* and 7.52 to 7.79 for *S. aburaviensis*.

### Effects of stationary and shaking conditions on the production of proteases

Stationary and shaking conditions have marked influence on protease production. *S. albolongus* showed maximum protease activity (683.60 U/ml) at stationary condition, whereas shaking at 100 rpm significantly increased protease production (847.28 U/ml) by *S. aburaviensis* (Figure 5).

### Effects of carbon and nitrogen sources on the production of proteases


To investigate the effects of various carbon and nitrogen sources, the isolates were allowed to grow in different media containing four different carbon sources and five organic and two inorganic nitrogen sources. *S. albolongus* exhibited highest enzyme activity (673.76 U/ml) in glucose and beef extract containing media (Figure 6). Maximum protease (734.95 U/ml) was released when 1% glucose, and 2% beef extract were used as a carbon and nitrogen source in the growth medium (Figure 7). The strain *S. aburaviensis* showed maximum enzyme activity (484.62 U/ml) in lactose and tryptone containing medium (Figure 8). The isolate *S. buraviensis* showed maximum protease activity (571.27 U/ml), when 1.5% lactose and 1.5% tryptone were used as carbon and nitrogen source, respectively (Figure 9).

### Discussion

The results of primary screening indicated that both *S. albolongus* and *S. aburaviensis* have the ability to produce extracellular protease in solid medium under static incubation condition. Then the isolates were

Medium pH	Biomass yield (mg/gm of substrate)	
	<i>S. albolongus</i>	<i>S. aburaviensis</i>
4	44	9
5	80	29
6	122	115
7	141	155
8	160	180
9	106	127


**Table 3:** Effects of medium pH on biomass yield.




**Figure 4:** Effects of incubation temperature on the production of protease by *S. albolongus* and *S. aburaviensis*. Enzyme-substrate reaction pH and temperature were 6.5 and  $35^\circ\text{C}$ , respectively.

Temperature ( $0^\circ\text{C}$ )	Biomass yield (mg/gm of substrate)	
	<i>S. albolongus</i>	<i>S. aburaviensis</i>
10	38	34
27	72	130
37	152	174
45	114	118


**Table 4:** Effect of temperatures on biomass yield.




**Figure 5:** Effects of stationary and shaking condition on the production of protease by *S. albolongus* and *S. aburaviensis*. Enzyme-substrate reaction pH and temperature were 6.5 and 35°C, respectively.



**Figure 6:** Effects of carbon and nitrogen sources on the production of protease by *S. albolongus*. Incubation period 5 days, incubation temperature 37°C, medium pH 7.0, enzyme-substrate reaction pH and temperature 6.5 and 37°C, respectively.



**Figure 7:** Effects of percent of carbon and nitrogen sources on the production of protease by *S. albolongus*. Incubation period 5 days, incubation temperature 37°C, medium pH 7.0, enzyme-substrate reaction pH and temperature 6.5 and 37°C, respectively.



**Figure 8:** Effects of carbon and nitrogen sources on the production of protease by *S. aburaviensis*. Incubation period 4 days, incubation temperature 37°C, medium pH 8.0, enzyme-substrate reaction pH and temperature 6.5 and 37°C, respectively.

allowed to grow in three liquid media and maximum enzyme activity were found in Gelatin-yeast extract-glucose broth and tryptone-dextrose-yeast extract broth for *S. albolongus* and *S. aburaviensis*, respectively. There was a gradual increase of protease production up to day 4 for both the isolates. Maximum production was obtained after 5 days of incubation for *S. albolongus*, and it was 4 days for *S. aburaviensis*, but maximum biomass was obtained after 6 days for both isolates. Maximum production of extra cellular protease after 120


hours of incubation by *Streptomyces* sp. 594, which was not growth associated, was reported by Azeredo et al. [25]. Our results were in concurrence with them.

Microorganisms are very sensitive to the concentration of hydrogen ions present in the medium. So, pH is one of the most important factors that determine the growth and the production of protease by microbes. *S. albolongus* preferred medium with neutral pH for maximum enzyme production, but maximum protease activities *S. aburaviensis* was obtained in medium having initial pH 8.0. Production of microbial proteases in neutral and alkaline medium pH was also reported by other authors [25-27]. Temperature is an important environmental factor for growth and enzyme production. The best incubation temperature for both the isolates was 37°C. A wide range of temperature (30°C-55°C) has been reported for optimum growth and protease production by *Streptomyces* sp. [13,25]. Extracellular protease from *S. albolongus* was optimally produced when incubated at static condition. About nine fold greater protease activity from *Streptomyces* sp. in stationary culture than produced under shake flask condition was reported by Gibb et al. [28]. Our finding with *S. albolongus* was in accordance with them. Shaking at 100 rpm markedly increase proteases production by *S. aburaviensis*. This finding was also reflected in other studies [24,26,28,29].

Microorganisms show a considerable variation in their nutrient requirements. Carbon and nitrogen sources are important variables that affect the growth and products of microbes [30]. Many authors had reported variability of carbon and nitrogen sources with different microorganisms [31-33]. Shafee et al. [30] had reported that maximum protease was produced in medium containing glucose and beef extract, which was similar to our findings with *S. albolongus*. In case of *S. aburaviensis*, highest protease production was observed in medium containing lactose and tryptone. Both the isolates produced proteases in response to both organic and inorganic forms of nitrogen, but preferred organic nitrogen for better production. Similar result was reflected in other studies [22,31]. Concentration of carbon and nitrogen sources in the production media significantly affects protease [34-37]. Variability in the production of protease with different percent of carbon and nitrogen sources was also observed in the present investigation. Glucose concentration higher than 1.0 % was found to reduce protease yield from *S. albolongus*, when 2.0% beef extract was used as nitrogen source. In case of *S. aburaviensis* protease production was gradually increased with the increase of lactose concentration, and 1.5% was found optimum when 1.5% tryptone was used as nitrogen source.

## Conclusion

*Streptomyces albolongus* and *Streptomyces aburaviensis* are potential microbes for extracellular protease production, and various



**Figure 9:** Effects of percent of carbon and nitrogen sources on the production of protease by *S. aburaviensis*. Incubation period 4 days, incubation temperature 37°C, medium pH 8.0, enzyme-substrate reaction pH and temperature 6.5 and 37°C, respectively.

environmental and nutritional factors have significant effects on their growth and protease production. In the present investigation, we have determined the optimum conditions for maximum production of extracellular proteases. The optimized media composition and cultural conditions might be implemented in large scale for the production of extracellular proteases by *Streptomyces albolongus* and *Streptomyces aburaviensis*.

## References

1. Anwar A, Saleemuddin M (2000) Alkaline protease from *Spilosoma obliqua*: Potential applications in bioformulation. *Biotechnol Appl Biochem* 2: 85-89.
2. Gupta MN, Roy I (2002) Applied biocatalysis: An overview. *Indian J Biochem Biophys* 39: 220-228.
3. Lin AN, Ding AY, Chen J, Liu SA, Zhang M, et al. (2007) Purification and characterization of two thermostable proteases from the thermophilic fungus *Chaetomium thermophilum*. *J Microbiol Biotechnol* 17: 624-631.
4. Nadeem M, Qazi JI, Baig S (2009) Effect of aeration and agitation rates on alkaline protease production by *Bacillus licheniformis* UV-9 mutant. *Turk J Biochem* 34: 89-96.
5. Allpress JD, Mountain G, Gowland PC (2002) Production, purification and characterization of an extracellular keratinase from *Lysobacter* NCIMB 9497. *Lett Appl Microbiol* 34: 337-342.
6. Sindhu R, Suprabha GN, Shashidhar S (2009) Optimization of process parameters for the production of alkaline protease from *Penicillium godlewskii* SBSS 25 and its application in detergent industry. *Afr J Microbiol Res* 3: 515-522.
7. Thumar JT, Singh SP (2007) Secretion of an alkaline protease from a salt-tolerant and alkaliphilic, *Streptomyces clavuligerus* strain MIT-1. *Braz J Microbiol* 38: 766-772.
8. Vishalakshi N, Lingappa K, Amena S, Prabhakar M, Dayanand A (2009) Production of alkaline protease from *Streptomyces gulbargensis* and its application in removal of blood stain. *Ind J Biotechnol* 8: 280-285.
9. Kumar CG, Takagi H (1999) Microbial alkaline proteases: from a bioindustrial viewpoint. *Biotechnol Adv* 17: 561-594.
10. Haki GD, Rakshit SK (2003) Developments in industrially important thermostable enzymes: A review. *Bioresour Technol* 89: 17-34.
11. Kim IS, Kim YB, Lee KJ (1998) Characterization of the leupeptin-inactivating enzyme from *Streptomyces exfoliatus* SMF13 which produces leupeptin. *Biochem J* 331: 539-545.
12. Patke Dey (1998) Proteolytic activity from a thermophilic *Streptomyces measporus* strain SDP4. *Lett Appl Microbiol* 26: 171-174.
13. Seong CN, Jo JS, Choi SK, Kim SW, Kim SJ, et al. (2004) Production, purification, and characterization of a novel thermostable serine protease from soil isolate, *Streptomyces tendae*. *Biotechnol Lett* 26: 907-909.
14. Tunga R, Shrivastava B, Banerjee R (2003) Purification and characterization of a protease from solid state cultures of *Aspergillus parasiticus*. *Process Biochem* 38: 1553-1558.
15. Paranthaman R, Alagusundaram K, Indhumathi J (2009) Production of protease from rice mill wastes by *Aspergillus niger* in solid state fermentation. *World J Agric Sci* 5: 308-312.
16. Vishwantha KS, Appu-Rao AG, Singh SA (2010) Acid protease production by solid-state fermentation using *Aspergillus oryzae* MTCC 5341: optimization of process parameters. *J Ind Microbiol Biotechnol* 37: 129-138.
17. Gupta R, Saxena RK, Chaturvedi P, Viridi JS (1995) Chitinase production by *Streptomyces viridificans*: its potential in cell wall lysis. *J Appl Bacteriol* 78: 378-383.
18. Kim I, Lee K (1995) Physiological roles of leupeptin and extracellular protease in mycelium development of *Streptomyces exfoliatus* SMF 13. *Microbiology* 141: 1017-1025.
19. Klar AJ, Halvorson HO (1975) Proteinase activities of *Saccharomyces cerevisiae* during sporulation. *J Bacteriol* 124: 863-869.
20. Matta H, Punj V, Kanwar SS (1997) An immuno-dot blot assay for detection of thermostable protease form *Pseudomonas* sp. AFT- 36 of dairy origin. *Lett Appl Microbiol* 25: 300-302.
21. Shalinisen S, Satyanarayana T (1993) Optimization of alkaline protease production by thermophilic *Bacillus licheniformis* S-40. *Ind J Microbiol* 33: 43-47.
22. Meyers SP, Ahearn DG (1977) Extracellular proteolyses by *Candida lipolytica*. *Mycologia* 69: 646-651.
23. Hayashi K, Fukushima D, Mogi K (1967) Alkaline proteinase of *Aspergillus sojae*. Physicochemical properties, amino acid composition and molecular conformation. *Agric Biol Chem Tokyo* 31: 1237-1241.
24. Azeredo LAID, Freire DMG, Soares RMA, Leite SGF, Coelho RRR (2004) Production and partial characterization of thermophilic proteases from *Streptomyces* sp. isolated from Brazilian cerrado soil. *Enzyme Microbiol Technol* 34: 354-358.
25. Rahman RN, Geok LP, Basri M, Salleh AB (2005) Physical factors affecting the production of organic solvent-tolerant protease by *Pseudomonas aeruginosa* strain K. *Bioresour Technol* 96: 429-436.
26. Kazan D, Denizci AA, Oner MN, Erarslan A (2005) Purification and characterization of a serine alkaline protease from *Bacillus clausii* GMBAE42. *J Ind Microbiol Biotechnol* 32: 335-344.
27. Gibb GD, Ordaz DE, Strohl WR (1989) Overproduction of extracellular protease activity by *Streptomyces C<sub>5</sub>-A<sub>13</sub>* in fed-batch fermentation. *Appl Microbiol Biotechnol* 31: 119-124.
28. Elibol M, Moreira AR (2005) Optimizing some factors affecting alkaline protease production by a marine bacterium *Teredinobacter turnirae* under solid substrate fermentation. *Process Biochem* 40: 1951-1956.
29. Shafee N, Aris SN, Abd Rahman RNZ, Basri M, Salleh AB (2005) Optimization of environmental and nutritional conditions for the production of alkaline protease by a newly isolated bacterium *Bacillus cereus* strain 146. *J Appl Sci Res* 1: 1-8.
30. Beg QK, Saxena RK, Gupta R (2002) De-repression and subsequent induction of protease synthesis by *Bacillus mojavensis* under fed-batch operations. *Process Biochem* 37: 1103-1109.
31. Haab D, Hagspiel K, Szakmary K, Kubicek CP (1990) Formation of the extra cellular proteases from *Trichoderma reesei* QM 9414 involved in cellulase degradation. *J Biotechnol* 16: 187-198.
32. Ito ET, Pereira GV, Miyagui DT, Pinotti MHP, Neves PMOJ (2007) Production of extracellular protease by a Brazilian strain of *Beauveria bassiana* reactivated on coffee berry borer, *Hypothenemus hampei*. *Braz Arch Biol Technol* 50: 1516-1531.
33. Nguyen TT, Quyen DT (2011) Overproduction of an extracellular protease from *Serratia* sp. DT3 just using soyabain powder. *World J Agric Sci* 7: 29-36.
34. Sobita RR, Rahman M, Chowdhury N (1995) Extracellular proteolytic activity of *Sacharomyces cerevisiae* strain DSM 1848. *Bangladesh J Microbiol* 12: 122.
35. Nadeem M, Qazi JI, Baig S, Syed QA (2008) Effect of medium composition on commercially important alkaline protease production by *Bacillus licheniformis* N-2. *Food Technol Biotechnol* 46: 388-394.
36. Braaksma M, Smilde AK, Van der Werf MJ, Punt PJ (2009) The effect of environmental conditions on extracellular protease activity in controlled fermentations of *Aspergillus niger*. *Microbiology* 155: 3430-3439.