OPEN ACCESS Freely available online

Journal of Clinical Microbiology and Antimicrobials

Commentary

Epidemiology of Campylobacter Infections: Trends and Emerging Risks

Rosie Clara*

Department of Microbiology, Sichuan University, Sichuan, China

ABOUT TO STUDY

Campylobacter infections, primarily caused by *Campylobacter jejuni* and *Campylobacter coli*, are among the most common causes of bacterial gastroenteritis worldwide. These infections are predominantly foodborne, transmitted mainly through contaminated food, water, and animal contact, with poultry being the primary reservoir. As a leading cause of diarrhea and abdominal cramps, *Campylobacter* infections present a significant public health burden. In addition to causing gastroenteritis, *Campylobacter* has been linked to severe complications, including Guillain-Barré syndrome (GBS), a rare neurological disorder. Over recent decades, the epidemiology of *Campylobacter* infections has evolved, with shifting trends, emerging risks, and growing concerns over antimicrobial resistance. This article reviews the epidemiology of *Campylobacter* infections, examining trends in incidence, risk factors, and the emerging challenges posed by antimicrobial resistance.

Global burden of *Campylobacter* infections

Campylobacter jejuni and *Campylobacter coli* are responsible for the majority of Campylobacter infections globally. The World Health Organization (WHO) estimates that *Campylobacter* causes approximately 96 million cases of gastroenteritis annually, resulting in significant morbidity, especially in children under five years of age and in immunocompromised individuals. While many cases of *Campylobacter* infection are self-limiting, the infection can lead to prolonged symptoms, including bloody diarrhea, fever, and dehydration. In some instances, *Campylobacter* infections can trigger GBS, which can cause severe neurological complications.

The burden of *Campylobacter* infections is particularly high in low- and middle-income countries, where sanitation and food safety practices may be inadequate. In high-income countries, *Campylobacter* remains a leading cause of foodborne disease, with outbreaks linked to contaminated poultry, unpasteurized milk, and untreated water.

Trends in *Campylobacter* infections

Over the years, the incidence of *Campylobacter* infections has fluctuated, with noticeable variations based on geographic location, seasonality, and changes in food safety practices. Several key trends in *Campylobacter* epidemiology have emerged:

Increased incidence in developed countries: In high-income countries, there has been a relatively stable or slightly increasing trend in *Campylobacter* infections over the past few decades. Despite improvements in food safety standards, *Campylobacter* remains the leading cause of bacterial gastroenteritis. This is partly due to the high consumption of poultry, which remains a primary source of infection. In countries like the United States and the European Union, surveillance systems report significant numbers of *Campylobacter* cases each year. Poultry, especially undercooked or contaminated chicken, continues to be the most common source of infection.

Seasonality of infections: *Campylobacter* infections exhibit a marked seasonality, with higher rates typically observed during the warmer months of the year. This seasonal variation is thought to be linked to increased foodborne transmission, particularly through the handling and consumption of poultry, which is more common during summer months and holiday seasons. Additionally, environmental conditions such as higher temperatures can facilitate the growth of *Campylobacter* in the food chain.

Geographic variations in incidence: The incidence of *Campylobacter* infections varies by region. In some countries, particularly in Europe and North America, well-established surveillance systems track the prevalence of *Campylobacter* infections, revealing both seasonal peaks and year-on-year fluctuations. Conversely, in lower-income regions, *Campylobacter* remains underreported due to limited healthcare infrastructure and diagnostic capabilities.

Increased recognition of non-gastrointestinal manifestations: In addition to gastrointestinal symptoms, there is growing recognition of the link between *Campylobacter* infections and non-gastrointestinal complications, such as GBS. This association has drawn increased attention to the neurological impacts of

Correspondence to: Rosie Clara, Department of Microbiology, Sichuan University, Sichuan, China; E-mail: rosieclara@eua.edu

Received: 30-Dec-2024, Manuscript No. JCMA-24-36633; **Editor assigned:** 01-Jan-2025, PreQC No. JCMA-24-36633 (PQ); **Reviewed:** 15-Jan-2025, QC No. JCMA-24-36633; **Revised:** 10-Jul-2025, Manuscript No. JCMA-24-36633 (R); **Published:** 17-Jul-2025, DOI: 10.35248/ CMA.25.09.229

Citation: Clara R, (2025) Epidemiology of Campylobacter Infections: Trends and Emerging Risks. J Clin Microbiol Antimicrob. 9:229

Copyright: © 2025 Clara R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Campylobacter and led to more targeted research into the mechanisms by which *Campylobacter* can trigger autoimmune responses.

Risk factors for campylobacter infections

Several risk factors contribute to the transmission and spread of *Campylobacter* infections, including environmental, behavioral, and food-related factors.

Poultry consumption: The most significant risk factor for *Campylobacter* infection is the consumption of contaminated poultry. Poultry, particularly chicken, is a primary reservoir for *Campylobacter*, which is present in the intestines of infected animals. Cross-contamination during food handling, such as improper washing or undercooking of chicken, is a major contributor to the spread of the bacteria.

Contaminated water: Waterborne transmission is another important route of infection. Contaminated drinking water, especially in areas without adequate sanitation or in rural settings, is a well-documented source of *Campylobacter* outbreaks. Unpasteurized milk is another potential source of infection, particularly when dairy products are produced under suboptimal hygiene conditions.

Contact with animals: Direct contact with animals, especially farm animals such as cattle, pigs, and poultry, is another risk factor for *Campylobacter* infection. In addition, handling of animal products, such as raw meat, increases the likelihood of exposure. Occupational exposure to animals, such as among farmers, veterinarians, or slaughterhouse workers, also carries a higher risk of infection.

Travel and international outbreaks: Traveling to regions with less stringent food safety regulations increases the risk of *Campylobacter* infection. International travelers are particularly susceptible to acquiring foodborne illnesses, and *Campylobacter* is a common cause of "traveler's diarrhea." This risk is heightened by the consumption of local food and water, particularly in countries with inadequate sanitation.

Immunocompromised individuals: Individuals with weakened immune systems, such as those undergoing chemotherapy, organ transplant recipients, or individuals with HIV/AIDS, are at increased risk of severe *Campylobacter* infections. For these individuals, the infection may be more severe and require medical intervention.

Emerging risks and antimicrobial resistance

One of the most pressing concerns regarding *Campylobacter* infections is the increasing prevalence of Antimicrobial Resistance (AMR). Over the past few decades, *Campylobacter* strains, particularly *Campylobacter jejuni*, have shown resistance to several commonly used antibiotics, including fluoroquinolones, macrolides, and tetracyclines. This resistance has significant implications for treatment, as *Campylobacter* infections are typically managed with antibiotics in severe cases or when complications such as GBS occur.

Resistance to Fluoroquinolones: Fluoroquinolones, including ciprofloxacin, have been widely used to treat *Campylobacter* infections, but resistance to this class of antibiotics has emerged as a major concern. Resistance is often linked to the widespread use of fluoroquinolones in animal husbandry, particularly in poultry production. This has led to the development of strains that are more difficult to treat in humans.

Resistance to Macrolides: Macrolides, such as azithromycin, are often used to treat resistant *Campylobacter* infections. However, resistance to macrolides has also been increasing, particularly in *Campylobacter jejuni*. This is particularly concerning because macrolides are considered a first-line treatment for severe cases, including those involving GBS.

Transfer of resistance between animal and human: Populations The use of antibiotics in livestock, particularly in poultry farming, plays a crucial role in the development of resistant *Campylobacter* strains. These resistant strains can then be transmitted to humans through consumption of contaminated meat, exacerbating the challenge of treating infections.

Prevention and control strategies

Effective prevention and control strategies are essential to reducing the burden of *Campylobacter* infections. Key measures include:

Improved food safety and hygiene practices: Ensuring that poultry is properly cooked and handled is critical to preventing *Campylobacter* infections. Public health campaigns focusing on food safety, such as avoiding cross-contamination and cooking chicken to the appropriate internal temperature, can significantly reduce the risk of infection.

Antimicrobial stewardship in agriculture: Restricting the use of antibiotics in animal agriculture, particularly in poultry, can help reduce the development and spread of resistant *Campylobacter* strains. Several countries have already implemented regulations to limit the use of antibiotics in livestock farming, which has shown promise in reducing resistance levels.

Surveillance and monitoring: Ongoing surveillance of *Campylobacter* infections and resistance patterns is crucial for monitoring trends and informing public health interventions. Surveillance systems should also focus on foodborne outbreaks and emerging sources of infection, including waterborne transmission.

CONCLUSION

Campylobacter infections remain a major cause of foodborne illness worldwide, with significant public health implications. While the incidence of infections remains high in both developed and developing countries, the evolving epidemiology of *Campylobacter* infections including the emergence of antimicrobial resistance presents new challenges for prevention and treatment. Effective food safety practices, improved surveillance, and antimicrobial stewardship are essential to reducing the burden of *Campylobacter* infections and addressing the growing problem of antimicrobial resistance.

Through coordinated efforts, it is possible to mitigate the risks associated with *Campylobacter* and protect public health in the years to come.