

Epigenetic Drift in Human Oocytes during Ovarian Aging and its Consequences for Embryo Development in Assisted Reproduction

Helena Varga*

Department of Reproductive Epigenomics, Central European Institute of Fertility Sciences, Budapest, Hungary

DESCRIPTION

Ovarian aging is a progressive biological process characterized not only by a decline in oocyte quantity but also by deterioration in oocyte molecular quality. Among the most important molecular changes observed with advancing maternal age is epigenetic drift, a phenomenon describing gradual and cumulative alterations in epigenetic marks such as Deoxyribonucleic Acid (DNA) methylation, histone modification, and chromatin organization. In assisted reproductive cycles, these changes are increasingly recognized as significant contributors to reduced embryo developmental potential, altered implantation rates, and increased risk of developmental abnormalities.

Epigenetic regulation plays a fundamental role in oocyte development, beginning during fetal life and continuing through follicular maturation. DNA methylation patterns are established during oogenesis and are essential for proper genomic imprinting, gene silencing, and developmental regulation after fertilization. These epigenetic marks ensure that genes are expressed in a parent-of-origin-specific manner and that early embryonic development proceeds in a coordinated fashion.

With increasing maternal age, the stability of these epigenetic marks becomes compromised. Epigenetic drift refers to the gradual loss of precision in DNA methylation patterns, leading to aberrant gene expression profiles. This may result in improper activation or silencing of genes involved in meiosis, spindle formation, mitochondrial function, and early embryonic development. Such alterations may not prevent fertilization but can significantly affect post-fertilization developmental competence.

Histone modifications also undergo age-associated changes. Histones are proteins around which DNA is wrapped, and their chemical modifications regulate chromatin accessibility. In aged oocytes, alterations in histone acetylation and methylation patterns have been observed, potentially affecting chromatin condensation and transcriptional regulation. These changes may impair the oocyte's ability to properly organize its genetic material during meiotic division. Chromatin structure within

oocytes becomes increasingly unstable with age. This instability can contribute to errors in chromosome segregation during meiosis, leading to aneuploid embryos. Aneuploidy is a major cause of implantation failure and early miscarriage in assisted reproductive treatment. Epigenetic drift may indirectly increase aneuploidy risk by weakening chromatin cohesion and spindle assembly integrity.

Mitochondrial function is closely linked to epigenetic regulation. Declining mitochondrial efficiency in aged oocytes can influence the availability of metabolic substrates required for epigenetic maintenance. Reduced adenosine triphosphate levels may impair enzymes responsible for DNA methylation and histone modification, further contributing to epigenetic instability. This interaction between metabolism and epigenetics forms a feedback loop that exacerbates oocyte aging. Environmental and lifestyle factors also contribute to epigenetic drift. Exposure to toxins, poor dietary habits, chronic stress, and metabolic disorders can accelerate epigenetic alterations within oocytes. These factors may interact with natural aging processes, leading to earlier onset of molecular decline even in relatively younger women. Assisted reproductive patients with such risk factors may therefore exhibit oocyte quality variations independent of chronological age.

During assisted reproductive stimulation, multiple oocytes are retrieved, each potentially exhibiting different degrees of epigenetic stability. This heterogeneity may explain variability in embryo development outcomes within the same cycle. Some oocytes may retain relatively stable epigenetic patterns and support normal embryogenesis, while others may exhibit altered developmental trajectories despite normal fertilization. Early embryonic development relies heavily on maternal epigenetic information until embryonic genome activation occurs. If epigenetic drift is present in the oocyte, improper gene regulation may persist into early cleavage stages. This may manifest as delayed cell division, abnormal gene expression profiles, or reduced blastocyst formation rates. Such effects are often subtle and may not be detectable through standard morphological assessment alone.

Correspondence to: Helena Varga, Department of Reproductive Epigenomics, Central European Institute of Fertility Sciences, Budapest, Hungary, E-mail: helena.varga.ceifs@epigenlabmail.org

Received: 02-Jun-2025, Manuscript No. JFIV-25-41861; **Editor assigned:** 04-Jun-2025, PreQC No. JFIV-25-41861 (PQ); **Reviewed:** 18-Jun-2025, QC No. JFIV-25-41861; **Revised:** 24-Jun-2025, Manuscript No. JFIV-25-41861 (R); **Published:** 02-Jul-2025, DOI: 10.35841/2375-4508.25.13.412.

Citation: Varga H (2025). Epigenetic Drift in Human Oocytes during Ovarian Aging and its Consequences for Embryo Development in Assisted Reproduction. *J Fertil In Vitro IVF World w Reprod Med Gent Stem Cell Biol.* 13:412.

Copyright: © 2025 Varga H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Assisted reproductive technologies such as intracytoplasmic sperm injection bypass natural selection mechanisms, allowing fertilization of oocytes with varying epigenetic integrity. While this increases chances of fertilization, it also highlights the importance of understanding oocyte molecular quality beyond structural appearance. Epigenetic profiling has therefore become an area of increasing interest in reproductive medicine research. Single-cell epigenomic technologies have enabled detailed analysis of DNA methylation and chromatin states in individual oocytes. These methods have revealed that epigenetic variability increases with maternal age and is associated with changes in genes involved in developmental regulation and cellular metabolism. Such findings provide molecular evidence supporting the concept of epigenetic drift as a contributor to reproductive aging.

Interventions aimed at reducing oxidative stress and improving metabolic health have been investigated for their potential to stabilize epigenetic marks. Antioxidant supplementation, lifestyle modification, and metabolic regulation may indirectly support epigenetic maintenance by reducing cellular stress.

However, the extent to which these approaches can reverse established epigenetic drift remains uncertain. Emerging research also explores whether epigenetic modifications in oocytes can be partially restored through *in vitro* maturation techniques or optimized culture conditions. While experimental data suggest some degree of epigenetic plasticity, clinical application remains limited and requires further validation.

CONCLUSION

Current evidence indicates that epigenetic drift in oocytes is a significant factor influencing reproductive aging and embryo developmental variability in assisted reproductive cycles. Alterations in DNA methylation, histone modification, and chromatin structure collectively contribute to reduced developmental competence and increased reproductive challenges with advancing maternal age. Continued research integrating epigenetics, reproductive biology, and clinical embryology may improve understanding of oocyte aging and support development of more effective fertility treatment strategies.