Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Opinion Article

Exosomal Communication Networks Influencing Cancer Progression and Therapeutic Responses

Victoria Andersson*

Department of Cellular Communication in Oncology, Nordic Institute of Translational Biosciences, Stockholm, Sweden

DESCRIPTION

Exosomes have become one of the most actively investigated mediators of intercellular communication in modern oncology because they transport diverse biological molecules between neighboring and distant cells. These nanosized extracellular vesicles originate from the endosomal pathway and are released into the extracellular environment after fusion of multivesicular bodies with the plasma membrane. Nearly every cell type within the human body produces exosomes, including malignant cells, fibroblasts, endothelial cells, macrophages, dendritic cells, lymphocytes, adipocytes, and epithelial cells. Once released, these vesicles circulate through blood, lymphatic fluid, saliva, urine, cerebrospinal fluid, and additional biological fluids, delivering molecular cargo capable of modifying the physiology of recipient cells. Their participation in cellular communication has transformed current understanding of cancer biology by demonstrating that malignant progression depends not only upon direct cell-to-cell contact but also upon long-distance molecular exchange occurring throughout the body. Within cancer research and immuno-oncology, exosomal signaling has become an important area of investigation because of its influence on disease evolution, biological heterogeneity, biomarker development, and therapeutic monitoring.

Unlike soluble signaling molecules that often act through membrane receptors, exosomes provide protected transport of complex molecular cargo enclosed within lipid bilayers. Their membranes shield biological molecules from enzymatic degradation while facilitating movement through body fluids over considerable distances. Recipient cells internalize exosomes through endocytosis, membrane fusion, receptor-mediated uptake, or phagocytosis, allowing transferred cargo to influence intracellular signaling pathways, metabolism, protein synthesis, and transcriptional regulation. This mechanism permits efficient communication among cellular populations occupying different anatomical locations.

Endothelial cells responsible for blood vessel formation also exchange exosomes extensively with surrounding cellular populations. Endothelial-derived exosomes contain proteins regulating vascular integrity, lipid metabolism, intracellular signaling, and communication with neighboring stromal cells. Similarly, malignant cell-derived exosomes influence endothelial physiology by modifying migration, proliferation, and organization of vascular networks. These interactions contribute to regional variation in nutrient delivery and oxygen availability observed within malignant tissues.

Lymphocyte populations also generate exosomes containing diverse molecular cargo associated with cellular activation, metabolism, and intracellular signaling. Cytotoxic T lymphocytes, helper T cells, regulatory T cells, B lymphocytes, plasma cells, and natural killer cells each produce vesicles reflecting their functional status. Uptake of lymphocyte-derived exosomes by surrounding stromal populations modifies communication networks operating throughout malignant tissues. Likewise, lymphocytes respond to exosomes originating from malignant cells and additional stromal populations, illustrating the complexity of reciprocal extracellular communication.

Metabolic adaptation significantly affects exosomal communication. Changes in glucose utilization, amino acid metabolism, lipid processing, and oxidative phosphorylation alter intracellular availability of proteins, metabolites, and nucleic acids selected for vesicle packaging. Environmental conditions including oxygen limitation, nutrient deprivation, and oxidative stress further influence exosome composition. Consequently, extracellular vesicles reflect the metabolic status of their originating cells while simultaneously transmitting metabolic information to neighboring populations.

Single-vesicle analysis has recently emerged as an important technological advancement complementing conventional bulk exosome studies. Earlier investigations frequently examined mixed vesicle populations obtained from body fluids, limiting recognition of vesicle heterogeneity. New analytical methods permit characterization of individual extracellular vesicles according to size, surface proteins, nucleic acid content, and

Correspondence to: Victoria Andersson, Department of Cellular Communication in Oncology, Nordic Institute of Translational Biosciences, Stockholm, Sweden, E-mail: victoria.andersson@nitb-sweden.edu

Received: 01-Jun-2026, Manuscript No. JCRIO-26-43992; **Editor assigned:** 03-Jun-2026, PreQC No. JCRIO-26-43992(PQ); **Reviewed:** 17-Jun-2026, QC No. JCRIO-26-43992; **Revised:** 24-Jun-2026, Manuscript No. JCRIO-26-43992 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2684-1266.26.12.292

Citation: Andersson V (2026). Exosomal Communication Networks Influencing Cancer Progression and Therapeutic Responses. J Cancer Res Immunooncol. 11:292.

Copyright: © 2026 Andersson V. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

molecular composition. These approaches reveal remarkable diversity among exosomes released even by the same cellular population.

Spatial molecular technologies further enhance understanding of exosomal communication by preserving anatomical relationships while evaluating molecular activity within intact tissues. Combining spatial transcriptomics with vesicle analysis enables investigators to identify regions characterized by active extracellular communication and determine relationships between vesicle-producing cells and recipient populations. Such integrated analyses contribute to comprehensive characterization of communication networks operating within malignant environments.

Continued investigation has also demonstrated that exosomal communication extends beyond local tissues. Vesicles released into circulation reach distant organs where they influence epithelial cells, stromal populations, vascular structures, and resident inflammatory cells. This systemic communication illustrates that malignant disease affects biological processes throughout the body rather than remaining restricted to primary anatomical sites. Better understanding of long-distance vesicle transport may strengthen interpretation of disease progression and therapeutic monitoring.

CONCLUSION

Future progress in cancer research and immuno-oncology will increasingly depend upon comprehensive investigation of exosomal communication networks linking diverse cellular populations. Exosomes participate in coordinated exchange of proteins, nucleic acids, lipids, metabolites, and signaling molecules among malignant cells, fibroblasts, endothelial cells, macrophages, dendritic cells, lymphocytes, and additional stromal populations. Integration of advanced molecular technologies, single-vesicle analysis, artificial intelligence, liquid biopsy, spatial biology, and computational modeling will continue improving knowledge of extracellular communication while supporting development of informative biomarkers and innovative therapeutic strategies across numerous forms of human malignancy.