

Exploring Novel Isocratic HPLC Method for Quantitative Determination of Cinnarizine and Piracetam in Their Capsule Preparations

El-Adl SM*, El-sadek ME and Hasan MH

Department of Medicinal Chemistry, Faculty of Pharmacy, Zagazig University, Zagazig, Egypt

Abstract

A novel isocratic HPLC method had been developed for rapid simultaneous separation and determination of cinnarizine and Piracetam in pure form or in pharmaceutical formulations within less than 10 minutes. Separation was carried out on a Hypersil gold C₁₈ (10µm, 100x4.6mm) column. Effect of pH and composition of mobile phase in addition to flow rate was studied. Calibration was obeyed in the range of (10-80) µg/ml for cinnarizine or (160-960) µg/ml for Piracetam. The method was applied for the simultaneous determination of these drugs in both bulk and pharmaceutical forms and this is the first published method that uses isocratic HPLC method for cinnarizine and piracetam determination in combined formulation and the method were validated according to ICH parameters.

Keywords: Cinnarizine; Piracetam; CinnarizineR; HPLC

Introduction

Cinnarizine is antihistaminic, ca-channel blocker and sedative drug [1]. It is described for the treatment of motion sickness, nausea and vertigo. The main problem with cinnarizine administration is drowsiness [2]. British pharmacopoeia states a non- aqueous titration method for cinnarizine assay. Cinnarizine have been determined by Several other reported methods, including spectrophotometric methods [3-7] and high-performance liquid chromatography (HPLC) [8-10].

Piracetam is a nootropic and psychopharmacological drug [11]. It is a synthetic cyclic derivative of GABA. It is described as a myoclonus and neuroprotective agent. The mechanism of action depends on eliminating of calcium chloride those results in a decrease of the rhythm rate and an increase of the contraction amplitude [12]. Manufacturing occur by condensing ethyl chloroacetate with 2-pyrrolidinone in the presence of a metal hydride and converting the ester into an amide with ammonia [13]. British pharmacopoeia states a chromatographic method for piracetam determination. Several methods have been developed for piracetam determination, including spectrophotometric methods [14-17], high-performance liquid chromatography (HPLC) [18-22] and electrochemical method [23].

Review of literature revealed that only limited reports have been published on the chromatographic assay of cinnarizine and piracetam in the combined formulation [24,25]. However, there is no report on the isocratic RP-HPLC method for assay these two drugs. The main objective of the present work is to describe a specific, linear, accurate, precise and a new reversed phase HPLC method using isocratic elution for the simultaneous determination of cinnarizine and piracetam in the combined formulation. The optimized mobile phase was determined as a mixture of acetonitrile:water (50:50 v/v), detection at 229 nm, pH 2.7 adjusted by orthophosphoric acid and a flow rate of 0.5 ml/min. Under these conditions, piracetam and cinnarizine eluted at 3.8, and 8.1 minutes respectively (Figures 1 and 2).

Experimental

Instrumentation

The HPLC system used was a computer based Agilent 1200 series instrument comprising of a quaternary pump, a UV detector (Thermo Scientific Co. USA) and Auto sampler (injector). The system equipped by Agilent chemstation PC program.

Materials and reagents

All reagents used were of analytical grade or HPLC grade. Ortho-phosphoric acid was supplied by (Germany), Acetonitrile HPLC grade and Water. Cinnarizine and piracetam standard working powders were kindly supplied by Egyptian sigma company (Egypt), and were used without further purification.




Figure 1: Cinnarizine structure IUPAC name: (E)-(Diphenylmethyl)-4-(3-phenylprop-2-enyl)piperazine.




Figure 2: Piracetam structure IUPAC name: 2-(2-oxopyrrolidin-1-yl)acetamide.

*Corresponding author: Sobhy M. El-Adl, Department of Medicinal Chemistry, Faculty of Pharmacy, Zagazig University, Zagazig, Egypt, Tel: 20 100 811 8736; E-mail: mar_hamdy87@yahoo.com

Received: June 04, 2016; **Accepted:** June 21, 2016; **Published:** June 27, 2016

Citation: El-Adl SM, El-sadek ME, Hasan MH (2016) Assessing the Quality of Household and Drinking Water in Tongi Industrial Zone of Bangladesh and Its Toxicological Impact on Healthy Sprague Dawley Rats. J App Pharm 8: 225. doi: 10.21065/1920-4159.1000225

Copyright © 2016 El-Adl SM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Chromatographic procedure

20 μ l from the sample solutions of the drugs were taken and detected at lambda 229 nm. Liquid chromatography was performed on a Hypersil gold C18 (10 μ m, 100x4.6mm) column. The used mobile phase was (acetonitrile: water 50:50 v/v) and pH at 2.7 using orthophosphoric acid. Mobile phase pumped at a flow rate equals to 0.5 ml/min at ambient temperature. Before all, the mobile phase was filtered by a 0.45 μ l Nylon membrane filter (USA,MA) under vacuum and degassed by ultra-sonication (USA, Vernon Hills).

Pharmaceutical preparation

Cinaretam capsules labeled to contain in each capsule 25 mg cinnarizine and 400 mg of piracetam and batch No. 805072.

Preparation of stock standard solutions

25 and 800 mg of cinnarizine and piracetam respectively weighted and dissolved in the least amount of methanol in 10 ml volumetric flask. Sonicating for 5 m and then completing the final volume with mobile phase. The resulted concentrations of solutions were of 5, 80 mg/ml of cinnarizine and piracetam respectively.

The solutions of calibration plot (working standard solutions)

To construct calibration plots, the working standard solutions have been prepared by dilution the stock standard solutions with mobile phase and preparing working solutions with concentration ranges (10-80 and 160-960 μ g/ml) for cinnarizine and piracetam respectively. Each solution (n=5) was injected in triplicate and chromatographed under the mentioned conditions above. Linear relationships were obtained when average drug standard peak area were plotted against the corresponding concentrations for each drug. Regression equation was computed.

Results and Discussion

Optimization of Chromatographic Conditions: All chromatographic conditions were illustrated in Table 1.

The chromatographic detection was performed at 229 nm after screening of absorption at UV spectrum using Surveyor photodiode array detector (PAD) (Thermo Scientific Co. USA) where we found that 229 is the best one that give more sensitivity to the method and also linearity and validation to the method. Detection at 190 nm was more sensitive but it was not valid as it did not give linear relationship between the concentration and the peak area. The method was performed on a Hypersil gold[®] C₁₈ (10 μ m, 100x4.6mm) column (Thermo Scientific Co. USA). Chromatographic conditions were optimized by changing the mobile phase composition, pH of the mobile phase, the flow rate and also studying the detection at different wave length. Different experiments were performed to optimize the mobile phase but adequate separation of drugs could be achieved by altering the composition of mobile phase from (methanol 60: water 40 v/v) [where both drugs appear at the same retention time] to (acetonitrile 60: water 40: v/v) [where cinnarizine peak is too broad and piracetam peak appear with tailing] at pH 2.7 by orthophosphoric acid, flow rate at 0.5 ml/min. and detection at 229 nm as shown in Figure 3.

The optimized mobile phase was determined as a mixture of acetonitrile:water (50:50, v/v) at a flow rate of 0.5 ml/min, pH 2.7 using orthophosphoric acid and detection at 229 nm as shown in Figure 4.

Method Validation

The developed methods were validated according to international conference on harmonization guidelines [26].

Specificity

Specificity is the ability to assess unequivocally the analyte in the presence of components which may be expected to be present. Typically these might include degradants, impurities, matrix, etc. A Bulk of Cinaretam capsules (solution contains excipients only) had been prepared by mixing its excipients like StarCap 1500, silicon , silicon dioxide, and calcium phosphate di or tri basic disodium edetate. Known concentration of studied excipients was added to the bulk




Figure 3: HPLC Chromatograms of authentic mixture (800 μ g. ml⁻¹ piracetam and 60 μ g ml⁻¹ of cinnarizine at different Mobile phase (A, B) at 229 nm, pH 2.7 and flow rate 0.5 ml/min.

(A) (acetonitrile 60: water 39: orthophosphoric acid 1 v/v) [where cinnarizine peak is too broad and piracetam peak appear with tailing].

(B) (methanol 60: water 39: orthophosphoric acid 1) v/v) [where both druges appear at the same retention time].




Figure 4: Typical HPLC chromatograms obtained from 20 μ l injections of piracetam at (3.888 min) and cinnarizine at (8.103 m) respectively under optimized chromatographic conditions.

Parameters	Conditions
Column	Hypersil gold C ₁₈ (10 μ m, 100x4.6mm) column
Mobile phase	Isocratic binary mobile phase of acetonitrile:water (50:50, v/v), filtered and degassed using 0.45 μ m membrane filter
Wave length nm	229
Flow rate, ml/min	0.5
Injected volume, μ l	10
Pressure, psig	11
Temperature	Ambient (25 \pm 5 $^{\circ}$ C)
pH	2.7 adjusted by orthophosphoric acid

Table 1: Chromatographic Conditions for the proposed method.

then was injected under the illustrated chromatographic conditions. Recovery results showed that the bulk has negligible effect Which Means That the bulk did not interfere with developed method.

Linearity and range

When the results of analytical procedure are directly proportional to injected concentration (in certain range) of analyte this mean that the ability of the method to give linearity in the specific range. For the conformation of linearity, a minimum of 5 concentrations is recommended [27]. Five Concentrations were chosen in the ranges (10-80 and 160-960 μ g/ml) for corresponding levels of 50-150% w/w of the nominal analytical concentration of cinnarizine and piracetam respectively. The linearity of peak area responses versus concentrations was demonstrated by linear least square regression analysis (Figures 5 and 6).

Limits of detection and Limits of quantitation

According to the ICH recommendations, determination of limits of detection and quantitation was based on the standard deviation of the y-intercepts of regression lines (n=3) and the slope of the calibration plots [26] as in Table 2.

Precision

The precision of the assay was investigated by measurement of both repeatability and Intermediate precision.

Repeatability: Repeatability was investigated by injecting a minimum of 6 determinations at 100% of the test concentration and percentage SD were calculated in Table 3.

Intermediate precision: In the inter-day studies, standard and sample solutions prepared as described above, were analyzed in triplicate on three consecutive days at 100% of the test concentration and percentage SD were calculated table 3.

Accuracy

Accuracy was assessed using 9 determinations over 3 concentration levels covering the specified range (80,100 and 120%). Accuracy was reported as percent recovery by the assay of known added amount of analyte in the sample Table 3.

Robustness

Robustness of an analytical procedure is a measure of its capacity to remain unaffected by small variations in method parameters and provides an indication of its reliability during normal usage [26]. Robustness was tested by studying the effect of changing mobile phase pH by \pm 0.5, the percentage of organic solvent (acetonitrile) in the mobile phase by \pm 5 %, temperature \pm 5 $^{\circ}$ C, wavelengths \pm 5 nm and flow rate \pm 0.1 ml/min had no significant effect on the chromatographic resolution of the method.

Stability of analytical solution

As a part of confirmation of robustness, the stability of standard stock solutions was evaluated. Analysis of the standard stock solutions after preparation 1, 2 and 3 days at room temperature and notification the change in peak area response over 3 days which was (0.56 and 1.09% for cinnarizine and piracetam respectively. Their solutions were found to be stable for 3 days at room temperature at least.

Applications on Pharmaceutical Preparation

The Pharmaceutical formulation containing stated drugs have been successfully analyzed by the proposed method. Results obtained were compared to those obtained by applying reported reference methods [27,28] for cinnarizine and piracetam respectively where Student's t-test and F-test were performed for comparison. Results are shown in table 4 where the calculated t and F values were less than




Figure 5: The linear regression equations of piracetam was $Y = 15328 X - 2775.1$ and regression coefficient (r^2)=0.9999, Where Y is the peak area of standard solution and X is the drug concentration.




Figure 6: The linear regression equations of cinnarizine was $Y = 651557 X - 2000000$, and regression coefficient (r^2)=0.9999, Where Y is the peak area of standard solution and X is the drug concentration.

Parameter	Cinnarizine	Piracetam
Mean	100.4	100.16
± SD	0.88	0.78
± RSD	0.87	0.78
± SE	0.39	0.35
Variance	0.77	0.62
Slope	651557	15328
L.D.	1.04	16
L.Q.	3.4	48

Average of three independent procedures.

Table 2: Results of the analysis for the proposed method.

	Cinnarizine		piracetam		
	AV ± SD mg/ml	AV ± SD %	AV ± SD mg/ml	AV ± SD %	
Repeatability	24.96 ± 0.01	99.86 ± 0.11%	400.16 ± 0.04	100.5 ± 1.1	
Intermediate precision	24.93 ± 0.03	100.9 ± 0.78%	399.93 ± 0.02	100.05 ± 0.78	
Accuracy & Recovery%	80%	19.96 ± 0.04	100.03 ± 0.63	319.75 ± 0.03	101.03 ± 0.63
	100%	25.11 ± 0.02	100.18 ± 0.98	400.08 ± 0.01	100.18 ± 0.98
	120%	30.51 ± 0.06	99.7 ± 1.30	479.97 ± 0.28	99.7 ± 1.30

Table 3: Repeatability and Intermediate precision and Accuracy (Recovery %) of cinnarizine and piracetam respectively.

Drug name	Recovery ± SD		Calculated t-values	Calculated F-values
	Proposed method	Reference method		
Cinnarizine [27]	99.74 ± 0.78	100.16 ± 1.07	0.264	3.49
Piracetam [28]	101.34 ± 0.79	100.04 ± 0.82	2.031	1.203

(Where the Tabulated t-values and F-ratios at p = 0.05 are 2.57 and 5.

Table 4: Statistical comparison of the proposed and published methods for determination of Cinnarizine and piracetam in the form of Cinaretam® capsules by reported method (T- student test) and (F-test for variance)

tabulated values for the both drugs which in turn indicate that there is no significant difference between proposed method and reference methods relative to accuracy and precision.

Conclusion

A novel RP-HPLC method for rapid simultaneous estimation of cinnarizine and piracetam within less than 10 minutes was developed and validated. Linearity was observed over a concentration range of 10 to 80 µg ml⁻¹ for cinnarizine and in the range of 160 to 960 µg. ml⁻¹ for piracetam. The results obtained indicate that the proposed method is very sensitive, rapid, accurate, selective, and reproducible. The method has been successfully applied for the analysis of marketed formulations. It can be used for the routine analysis of formulations containing any one of the above drugs or their combinations without any alteration in the assay. The most we can talk about, HPLC method is also being used in other drug formulation recently such as nanomedicine [29] which again shows the strength of the method.

Acknowledgments

The authors deeply thank Dr. Esslam Hamed (faculty of veterinary medicine, zagazig university) for his help during the trial conditions of HPLC work.

References

- Monographs Medicinal and Pharmaceutical Substances (2014) British Pharmacopoeia.
- Macnair AL (1983) Cinnarizine in the prophylaxis of car sickness in children. *Curr Med Res Opin* 8: 451-455.
- Kalyankar M, Wadher S (2014) Simultaneous Estimation and Development of UV Spectroscopic Method for Determination of Cinnarizine and Domperidone in Bulk and Pharmaceutical Formulation. *International Journal of PharmTech Research* 6: 323-329.
- Abdine H, Belal F, Zoman N (2002) Simple spectrophotometric determination of cinnarizine in its dosage forms. *Farmaco* 57: 267-271.
- Abdul-Azim M (2004) Spectroscopic and Spectrofluorimetric determination of cinnarizine and Flunarizine dihydrochloride in pure and dosage form. *Bull Fac Pharm Cairo Univ* 42: 27-37.
- Daabees H (1999) A New Sensitive Spectrophotometry Method for the Analysis of Some Antihistamines. *An International Journal for Rapid Communication* 32: 913-930.
- Mohamed IW, Fathalla B, Nahed EE, Amina A (2008) Second-Derivative Synchronous Fluorescence Spectroscopy for the Simultaneous Determination of Cinnarizine and Nicergoline in Pharmaceutical Preparations. *An international journal of analytical science* 91: 349-359.
- Sirisha M, Kumari A (2013) Validation of RP-HPLC for rapid simultaneous determination of cinnarizine and domperidone in bulk and pharmaceutical dosage form. *Journal of Pharmaceutical & Scientific Innovation* 2: 40-46.
- Dina S, Tarek B (2014) Validated HPTLC method for the simultaneous determination of cinnarizine and dimenhydrinate in their combined dosage form. *Journal of the Association of Arab Universities for Basic and Applied Sciences* 7: 46-51.
- Argekar AP, Shah SJ (1999) Simultaneous determination of cinnarizine and domperidone maleate from tablet dosage form by reverse phase ion pair high performance liquid chromatography. *J Pharm Biomed Anal* 19: 813-817.
- Monographs Medicinal and Pharmaceutical Substances (2013) British pharmacopoeia.
- Müller WE, Eckert GP, Eckert A (1999) Piracetam: novelty in a unique mode of action. *Pharmacopsychiatry* 32 Suppl 1: 2-9.
- Ullmann's Encyclopedia of Industrial Chemistry (2003) 6th (eds), Federal Republic of Germany: Wiley-VCH Verlag GmbH and Co.
- Bhowmick A, Khandelwal KR, Mhaske DV, Swapnil K (2003) Analytical Method Development and Validation for Piracetam as Bulk and in Pharmaceutical Formulation. *International Journal of PharmTech Research* 2: 201-204.
- Hadad GM, Abdel-Salam RA, Emara S (2012) Optimized and validated flow-injection spectrophotometric analysis of topiramate, piracetam and levetiracetam in pharmaceutical formulations. *Acta Pharmaceutica* 61: 377-389.
- Sivadas A, Sathi A, Sathi K, Rahate KP (2013) Development and validation of spectrophotometric methods for simultaneous estimation of citicolone and piracetam in tablet dosage form. *J Pharm Bioallied Sci* 5: 202-207.
- Ei-Saharty YS (2008) Simultaneous determination of piracetam and vincamine by spectrophotometric and high-performance liquid chromatographic methods. *J AOAC Int* 91: 311-321.
- Siddiqui FA, Sher N, Shafi N, Wafa Sial A, Ahmad M, et al. (2014) Development of new method for simultaneous analysis of piracetam and levetiracetam in pharmaceuticals and biological fluids: application in stability studies. *Biomed Res Int* 2014: 758283.
- Arayne MS, Sultana N, Siddiqui FA, Mirza AZ, Qureshi F, et al. (2010) Simultaneous determination of piracetam and its four impurities by RP-HPLC with UV detection. *J Chromatogr Sci* 48: 589-594.
- Ardhani DL, Andi TP, Tini P, Evi U, Mochammad Y, et al. (2005) HPLC Determination of Piracetam in Tablets. *Journal of Liquid Chromatography and Related Technologies* 28: 1407-1416.
- Curticapean A, Imre S (2007) New validated method for piracetam HPLC determination in human plasma. *J Biochem Biophys Methods* 69: 273-281.
- Louchahi K, Tod M, Bonnardel P, Petitjean O (1995) Determination of piracetam in human plasma and urine by liquid chromatography. *J Chromatogr B Biomed Appl* 663: 385-389.
- Blinkov IL, Kondratenko SN, Kukes VG, Starodubtsev AK (2013) Voltammetry on Solid Electrodes for Quantitative Determination of Fenotropil and Piracetam in the Drug Substances, Dosage Forms, and Blood Plasma. *Pharmaceutical Chemistry Journal* 46: 750-758.

24. Ola M E, Nagwan HZ, Mohammad AM (2013) Development and Validation of RP-LC Method for the Determination of Cinnarizine/Piracetam and Cinnarizine/Heptaminol Acefyllinate in Presence of Cinnarizine Reported Degradation Products. *Anal Chem Insights* 8: 99-106.
25. Guidance for Industry (1996) Q2B of Analytical Procedures: Methodology.
26. Heda A, Sonawane R, Naranje GH, Puranik PK (2010) A Rapid Determination of Cinnarizine in Bulk and Pharmaceutical Dosage Form by LC. *E-Journal of Chemistry* 7: 1080-1084.
27. Muthukumar S, Shanthi R, Selvakumar D, Banji D (2013) Development and validation of RP-HPLC method for the quantification of piracetam and Mecoblamine. *International Journal of Biological & Pharmaceutical Research* 4: 681-684.
28. Sun Y, Kaplan JA, Shieh A, Sun HL, Croce CM, et al. (2016) Self-assembly of a 5-fluorouracil-dipeptide hydrogel. *Chem Commun (Camb)* 52: 5254-5257.
29. Agrahari V, Youan BB (2012) Sensitive and rapid HPLC quantification of tenofovir from hyaluronic acid-based nanomedicine. *AAPS PharmSciTech* 13: 202-210.