

Exploring the Vital Role of Proteoglycans as Building Blocks of Connective Tissues

Caroline Gray*

Department of Laboratory Medicine, Institute of Biomedicine, Sahlgrenska Academy at the University of Gothenburg, Gothenburg, Sweden

DESCRIPTION

Connective tissues are a vital component of the human body, providing structural support and maintaining the integrity of various organs and tissues. Proteoglycans are a family of macromolecules that are integral to the functioning of these connective tissues. In this article, we had discussed, what proteoglycans are, their structure, their roles in the body, and some of the diseases associated with their dysfunction [1,2].

Proteoglycans are a type of glycoprotein that are found in the extracellular matrix of connective tissues, such as cartilage, bone and skin. They are composed of a core protein to which Glycosaminoglycans (GAGs) are attached. GAGs are long chains of complex sugars that give proteoglycans their unique properties. There are several types of GAGs, including chondroitin sulfate, keratan sulfate and heparan sulfate [3,4].

Structure of proteoglycans

Proteoglycans are large macromolecules that can range in size from a few hundred kilodaltons to several megadaltons. They consist of a core protein that is attached to one or more GAG chains. The core protein can be a transmembrane protein or a protein that is secreted into the extracellular matrix. The GAG chains are long, linear polymers that are composed of repeating disaccharide units. The GAG chains are covalently attached to the core protein at specific sites on the protein molecule [5,6].

Role of proteoglycans in the body

Proteoglycans play several important roles in the body. One of their primary functions is to provide structural support to tissues. In cartilage, for example, proteoglycans help to form a dense, gel-like matrix that cushions the joints and absorbs shock. In bone, proteoglycans are a critical component of the extracellular matrix that provides the strength and flexibility needed to support the weight of the body [7].

Proteoglycans also play a crucial role in regulating the activity of growth factors and cytokines. Many of these signaling molecules

are bound to proteoglycans in the extracellular matrix, which helps to localize their activity and prevent them from diffusing too far from their site of release. In this way, proteoglycans help to regulate cell behavior and ensure that tissues grow and develop properly [8].

Finally, proteoglycans are also important for maintaining the hydration of tissues. The GAG chains in proteoglycans are highly negatively charged, which attracts water molecules and helps to maintain the hydration of tissues. This is particularly important in tissues such as cartilage, which are under constant compression and need to be able to resist deformation [9].

Diseases associated with proteoglycan dysfunction

Dysfunction of proteoglycans can lead to several diseases and conditions. One of the most well-known is osteoarthritis, a degenerative joint disease that is characterized by the breakdown of cartilage in the joints. In osteoarthritis, the proteoglycans in the cartilage are degraded, leading to a loss of hydration and the breakdown of the extracellular matrix [10].

Other diseases that are associated with proteoglycan dysfunction include Ehlers-Danlos syndrome, a genetic disorder that affects the connective tissue in the body, and mucopolysaccharidoses, a group of metabolic disorders that are characterized by the accumulation of GAGs in the body [11].

CONCLUSION

In conclusion, proteoglycans are complex macromolecules that are essential for the proper functioning of connective tissues in the body. They provide structural support, regulate the activity of growth factors and cytokines, and maintain the hydration of tissues. Dysfunctions in proteoglycans can lead to several diseases and conditions, including osteoarthritis, Ehlers-Danlos syndrome, and mucopolysaccharidoses. Understanding the roles of proteoglycans in the body can help to identify potential therapeutic targets for these diseases and improve the overall understanding of the mechanisms that underlie connective tissue function.

Correspondence to: Caroline Gray, Department of Laboratory Medicine, Institute of Biomedicine, Sahlgrenska Academy at the University of Gothenburg, Gothenburg, Sweden, E-mail: carolinegray@yahoo.com

Received: 14-Mar-2025, Manuscript No. JGB-23-22928; **Editor assigned:** 17-Mar-2025, PreQC No. JGB-23-22928 (PQ); **Reviewed:** 31-Mar-2025, QC No. JGB-23-22928; **Revised:** 07-Apr-2025, Manuscript No. JGB-23-22928 (R); **Published:** 14-Apr-2025, DOI: 10.35841/2168-958X.25.14.307

Citation: Gray C (2025) Exploring the Vital Role of Proteoglycans as Building Blocks of Connective Tissues. J Glycobiol. 14:307

Copyright: © 2025 Gray C. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCES

1. Wight TN, Kinsella MG, Qwarnström EE. The role of proteoglycans in cell adhesion, migration and proliferation. *Curr Opin Cell Biol.* 1992;4(5):793-801.
2. Wight TN. A role for proteoglycans in vascular disease. *Matrix Biol.* 2018;71:396-420.
3. Yanagishita M. Function of proteoglycans in the extracellular matrix. *Pathol Int.* 1993;43(6):283-93.
4. Hardingham TE, Fosang AJ. Proteoglycans: Many forms and many functions. *FASEB J.* 1992;6(3):861-70.
5. Ruoslahti E. Structure and biology of proteoglycans. *Annu Rev Cell Biol.* 1988;4(1):229-55.
6. Kjellén L, Lindahl U. Proteoglycans: Structures and interactions. *Annu Rev Biochem.* 1991;60(1):443-75.
7. Roughley PJ. The structure and function of cartilage proteoglycans. *Eur Cell Mater.* 2006;12(9):92-101.
8. Heindryckx F, Li JP. Role of proteoglycans in neuro-inflammation and central nervous system fibrosis. *Matrix Biol.* 2018;68:589-601.
9. Song Y, Zhang F, Linhardt RJ. Analysis of the glycosaminoglycan chains of proteoglycans. *J Histochem Cytochem* 2021;69(2):121-35.
10. Pessentheiner AR, Ducasa GM, Gordts PL. Proteoglycans in obesity-associated metabolic dysfunction and meta-inflammation. *Front Immunol.* 2020;11:769.
11. Guo MH, Stoler J, Lui J, Nilsson O, Bianchi DW, Hirschhorn JN, et al. Redefining the progeroid form of Ehlers–Danlos syndrome: Report of the fourth patient with B4GALT7 deficiency and review of the literature. *Am J Med Genet A.* 2013;161(10):2519-27.