Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Commentary

Extracellular Matrix Remodeling and Immune Cell Dynamics in Solid Tumors

Rafael Moreno*

Department of Cellular Oncology, Iberian School of Medical Sciences, Valencia, Spain

DESCRIPTION

Cancer progression is influenced not only by the biological behavior of malignant cells but also by the surrounding tissue environment in which these cells exist. Every solid tumor develops within a highly organized structure composed of immune populations, fibroblasts, endothelial cells, adipocytes, extracellular proteins, signaling molecules, and vascular networks. Among these components, the extracellular matrix has become an important subject of investigation because it provides structural organization while simultaneously influencing cellular communication, migration, survival, and tissue architecture. Rather than serving solely as a physical scaffold, the extracellular matrix continuously exchanges biochemical and mechanical signals with neighboring cells. These interactions affect immune surveillance, tumor expansion, vascular organization, and responses to treatment, making extracellular matrix remodeling a major topic within cancer immuno-oncology.

The extracellular matrix consists of collagen fibers, elastin, fibronectin, laminins, proteoglycans, glycoproteins, and numerous associated molecules arranged in a complex three-dimensional structure. In healthy tissues, this matrix supports organ integrity, regulates cell adhesion, maintains mechanical stability, and participates in normal physiological repair. Matrix composition differs according to organ type because each tissue possesses unique structural and functional requirements. Continuous renewal of extracellular components occurs throughout life through balanced synthesis and degradation controlled by resident stromal cells and specialized enzymes.

During malignant transformation, this balance gradually changes. Cancer cells communicate extensively with surrounding fibroblasts, immune populations, endothelial cells, and perivascular cells, producing signals that modify matrix production and degradation. As disease advances, extracellular matrix architecture frequently becomes denser, less organized, and mechanically stiffer than normal tissue. These structural alterations influence cellular movement, nutrient diffusion, oxygen distribution, and immune cell infiltration. Consequently, the extracellular matrix becomes an active participant in disease progression rather than remaining an inactive structural framework.

Fibroblasts represent one of the principal contributors to extracellular matrix remodeling within tumors. Under physiological conditions, fibroblasts maintain connective tissue integrity by synthesizing collagen and other structural proteins while participating in wound repair following tissue injury. Within malignant tissue, prolonged exposure to signaling molecules released by cancer cells modifies fibroblast activity. These activated fibroblasts increase production of extracellular proteins, remodeling enzymes, cytokines, and growth factors that collectively reshape the surrounding microenvironment. Their biological behavior varies among tumor types, reflecting considerable functional diversity.

Mechanical properties of the tumor microenvironment have attracted increasing scientific attention. Healthy tissues maintain organ-specific elasticity that supports normal cellular behavior. Progressive accumulation of extracellular proteins within tumors increases tissue rigidity, altering mechanical forces experienced by resident cells. Malignant cells respond to these physical changes through mechanosensitive signaling pathways regulating proliferation, migration, and survival. Immune populations also detect alterations in tissue stiffness, influencing their movement and functional activity. Mechanical communication therefore complements biochemical signaling throughout the tumor microenvironment.

Matrix-remodeling enzymes participate extensively in maintaining extracellular architecture. These enzymes continuously degrade existing matrix proteins while allowing synthesis of new structural components. During cancer progression, altered enzyme activity changes tissue organization, creating pathways that facilitate cellular migration while modifying release of matrix-associated signaling molecules. Controlled degradation of extracellular components also generates biologically active fragments capable of influencing inflammation, vascular development, and immune regulation. The balance between matrix synthesis and degradation therefore remains essential for maintaining tissue organization.

Blood vessel development is closely associated with extracellular matrix remodeling. Endothelial cells require coordinated interaction with surrounding matrix proteins during formation of new vascular structures. Changes in matrix composition

Correspondence to: Rafael Moreno, Department of Cellular Oncology, Iberian School of Medical Sciences, Valencia, Spain, E-mail: rafael.moreno@isms-spain.edu

Received: 27-Feb-2026, Manuscript No. JCRIO-26-43980; **Editor assigned:** 02-Mar-2026, PreQC No. JCRIO-26-43980(PQ); **Reviewed:** 16-Mar-2026, QC No. JCRIO-26-43980; **Revised:** 23-Mar-2026, Manuscript No. JCRIO-26-43980 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2684-1266.26.12.279

Citation: Moreno R (2026). Extracellular Matrix Remodeling and Immune Cell Dynamics in Solid Tumors. J Cancer Res Immunooncol. 11:279.

Copyright: © 2026 Moreno R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

influence endothelial migration, adhesion, and organization into functional vessels. Tumor-associated vasculature frequently displays irregular branching patterns, variable diameter, and inconsistent permeability. These abnormalities contribute to unequal oxygen delivery, heterogeneous nutrient availability, and regional differences in therapeutic distribution throughout malignant tissue.

Macrophages contribute significantly to extracellular matrix remodeling through secretion of remodeling enzymes, cytokines, and growth factors. These immune cells participate in removal of damaged tissue, regulation of inflammatory responses, and coordination of repair processes under physiological conditions. Within tumors, macrophages interact continuously with fibroblasts and malignant cells, influencing collagen deposition, matrix degradation, and tissue organization. Their activities differ according to local environmental conditions, resulting in multiple macrophage populations with distinct biological functions.

Neutrophils also participate in modifying extracellular architecture. Traditionally recognized for their role in acute inflammatory responses, these cells release enzymes, extracellular DNA structures, and signaling molecules capable of influencing matrix composition. Their interactions with fibroblasts, macrophages, and endothelial cells contribute to dynamic remodeling processes occurring throughout cancer progression. Continued investigation seeks to clarify how neutrophil-mediated matrix alterations influence long-term disease behavior.

CONCLUSION

Future progress in cancer immuno-oncology will depend upon comprehensive understanding of structural and cellular interactions occurring within the tumor microenvironment. Extracellular matrix remodeling represents a dynamic biological process involving coordinated communication among fibroblasts, immune populations, endothelial cells, malignant cells, and numerous soluble mediators. Rather than functioning as passive connective tissue, the extracellular matrix actively influences immune surveillance, vascular organization, metastatic dissemination, and therapeutic response. Continued investigation into these interconnected biological systems may improve biomarker discovery, refine diagnostic assessment, and support development of innovative therapeutic strategies directed toward improving long-term outcomes for individuals affected by solid malignancies.