OPEN ACCESS Freely available online

Anatomy & Physiology: Current Research

Commentary

Functional Anatomy of Peripheral Nerves: Diagnostic and Therapeutic Advances

Kenji Sato*

Department of Experimental Physiology, Osaka Biomedical College, Osaka, Osaka Prefecture, Japan

DESCRIPTION

The peripheral nervous system serves as the essential communication network between the central nervous system and the rest of the body, transmitting sensory information, coordinating motor function and regulating autonomic activities. Peripheral nerves are highly organized anatomical structures composed of axons, Schwann cells, connective tissue layers and an extensive vascular supply, all of which contribute to their functional integrity. Advances in anatomy, neurophysiology, imaging and regenerative medicine have significantly improved our understanding of peripheral nerve function and pathology. These developments have transformed both the diagnosis and treatment of peripheral nerve disorders, offering new opportunities to restore function and improve patient outcomes. Functional anatomy provides the foundation for understanding peripheral nerve physiology. Each peripheral nerve contains bundles of myelinated and unmyelinated fibers enclosed by the endoneurium, perineurium and epineurium, which provide mechanical protection while maintaining the internal microenvironment required for efficient nerve conduction. The arrangement of motor, sensory and autonomic fibers within these structures allows peripheral nerves to coordinate complex physiological functions, including voluntary movement, tactile sensation, pain perception and autonomic regulation. Even minor anatomical disruptions can significantly impair nerve transmission, emphasizing the importance of detailed anatomical knowledge in clinical practice.

One of the most notable advances in recent years has been the development of high resolution imaging techniques capable of visualizing peripheral nerves in vivo. High frequency ultrasonography and Magnetic Resonance Neurography (MRN) now provide detailed assessments of nerve morphology, fascicular architecture, vascularity and surrounding soft tissues. These imaging modalities enable clinicians to identify nerve compression, traumatic injuries, inflammatory neuropathies and tumors with remarkable precision. Unlike conventional anatomical imaging, modern neuroimaging provides both structural and functional information, facilitating early diagnosis and individualized treatment planning. Electrophysiological evaluation remains an indispensable component of peripheral nerve assessment. Nerve conduction studies and electromyography provide objective measurements of nerve function by evaluating conduction velocity, signal amplitude and muscle activation patterns. When combined with advanced imaging, these techniques offer complementary information that improves diagnostic accuracy and distinguishes between axonal injury, demyelination and neuromuscular disorders. This integrated approach has become particularly valuable in the diagnosis of entrapment neuropathies, diabetic neuropathy, inflammatory neuropathies and traumatic nerve injuries.

The understanding of peripheral nerve regeneration has also expanded considerably. Unlike neurons within the central nervous system, peripheral nerves possess a remarkable capacity for regeneration following injury. Schwann cells play a central role by clearing cellular debris, secreting neurotrophic factors, guiding axonal regrowth and reconstructing myelin sheaths around regenerating axons. Nevertheless, successful regeneration depends on multiple factors, including the severity of injury, the distance required for axonal growth, vascular integrity and timely clinical intervention. Delayed repair often results in irreversible muscle atrophy and reduced functional recovery, underscoring the importance of early diagnosis and treatment. Regenerative medicine has introduced promising therapeutic approaches that extend beyond conventional microsurgical repair. Tissue engineered nerve conduits, biodegradable scaffolds, stem cell therapy, platelet rich plasma and growth factor delivery systems are being investigated to enhance nerve regeneration and functional recovery. These biologically inspired strategies seek to recreate the optimal microenvironment for axonal growth while minimizing scar formation and chronic inflammation. Although many of these approaches remain under clinical evaluation, they represent an important step toward improving outcomes for patients with complex nerve injuries. Artificial intelligence (AI) is emerging as another transformative tool in peripheral nerve research and clinical practice. Machine learning algorithms can analyze imaging datasets, electrophysiological recordings and clinical parameters to identify subtle pathological changes, predict recovery potential and assist in treatment selection.

Correspondence to: Kenji Sato, Department of Experimental Physiology, Osaka Biomedical College, Osaka, Osaka Prefecture, Japan, E-mail: kenji.sato@obc.jp

Received: 03-Mar-2025, Manuscript No. APCR-25-43568; **Editor assigned:** 05-Mar-2025, PreQC No. APCR-25-43568 (PQ); **Reviewed:** 19-Mar-2025, QC No. APCR-25-43568; **Revised:** 25-Mar-2025, Manuscript No. APCR-25-43568 (R); **Published:** 02-Apr-2025, DOI: 10.35248/2161-0940.25.15.532

Citation: Sato K (2025). Functional Anatomy of Peripheral Nerves: Diagnostic and Therapeutic Advances. Anat Physiol. 15:532.

Copyright: © 2025 Sato K. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.